

The Impact Of Rs2066715 And Rs2230808 Polymorphisms In ABCA1 Gene With The Risk Of Diabetes And Coronary Artery Disease In North Indians

Nisha¹, Vikas kumari², Rajan³, Monika⁴, Priti devi⁵, Geeta Dhiman⁶, Jasbir Singh⁷

¹Department of Biochemistry, Kurukshetra University, Kurukshetra PhD scholar Department of Biochemistry, Kurukshetra University, Kurukshetra Email Id – kholanisha5@gmail.com

²PhD scholar Department of Biochemistry, Kurukshetra University, Kurukshetra Email ID – vikasmulodia1997@gmail.com

³PhD scholar Department of Biochemistry, Kurukshetra University, Kurukshetra Email ID – rajansharma6675@gmail.com

⁴PhD scholar Department of Biochemistry, Kurukshetra University, Kurukshetra Email ID - monikalamba2018@gmail.com

⁵PhD scholar Department of Biochemistry, Kurukshetra University, Kurukshetra Email ID – pritamalik0117@gmail.com

⁶PhD scholar Department of Biochemistry, Kurukshetra University, Kurukshetra Email ID – geetadhiman7791@gmail.com

⁷Department of Biochemistry, Kurukshetra University, Kurukshetra

Corresponding author: Dr. Jasbir Singh (Professor)

Department of Biochemistry, Kurukshetra University, Kurukshetra Email id – jasbirdhanda@gmail.com

Abstract

Background: Polymorphisms in the ABCA1 gene are linked to lipid metabolism, insulin resistance, and cardiovascular disease. This study assessed the correlation between ABCA1 rs2066715 and rs2230808 polymorphisms and Type 2 Diabetes Mellitus (T2DM), Coronary Artery Disease (CAD), and T2DM with CAD in a North Indian cohort.

Material and Method: A total of 600 participants were separated into four groups: healthy controls, T2DM, CAD, and T2DM with CAD patients (150 each). Clinical and biochemical data were analyzed. The rs2066715 and rs2230808 polymorphisms were genotyped using the PCR-RFLP method. The statistical analysis was carried out with SPSS version 31.

Result: T2DM, CAD, and T2DM with CAD patients had significantly higher FBG, TC, TG, LDL-C, HbA1c, blood pressure, and fasting insulin levels, but HDL-C levels were significantly lower ($p < 0.05$). There was no significant connection between the rs2066715 polymorphism and illness susceptibility. However, the rs2230808 polymorphism was significantly associated with T2DM, CAD, and T2DM with CAD. The A allele and TA+AA genotypes were linked to higher disease risk and a poor lipid profile. Haplotype study found that the CT haplotype is protective, whereas the CA and TA haplotypes enhance disease risk.

Conclusion: The ABCA1 rs2230808 polymorphism may serve as an important genetic marker for T2DM and CAD susceptibility in the North Indian population.

Keywords: ABCA1 gene, Coronary artery disease, Lipid metabolism, Haplotype analysis, North Indian Population

Abbreviations: ABCA1 = ATP-binding cassette transporter A1, T2DM = Type 2 Diabetes Mellitus, CAD = Coronary Artery Disease, CVD = Cardiovascular Disease, SNP = Single Nucleotide Polymorphism, PCR-RFLP = Polymerase Chain Reaction–Restriction Fragment Length Polymorphism, DNA = Deoxyribonucleic Acid, FBG = Fasting Blood Glucose

1. Introduction:

Diabetes mellitus is a chronic metabolic disorder in which there is hyperglycemia because of either less insulin or resistance to insulin [1]. The global burden of diabetes is quickly increasing, with over 536 million persons affected in 2021, and the figure is predicted to rise significantly by 2045 [2]. India is one of the countries with a maximum number of diabetic patients and a high percentage of the patients are not diagnosed [3]. In “Type 2 diabetes mellitus (T2DM)”, hyperglycemia and insulin resistance are associated with development of numerous comorbidities including cardiovascular disease (CVD), nephropathy, neuropathy, and retinopathy [4]. The “ATP-binding cassette transporter A1 (ABCA1)” gene plays a role in cholesterol homeostasis among the genetic variables involved in lipid metabolism and cardiovascular issues [5]. The ABCA1 gene resides at the 9q31.1 locus and is highly expressed in different organs such as liver, heart, pancreas, gut, macrophages, and endothelial cells [6]. It regulates reverse cholesterol transport by promoting the clearance of intracellular cholesterol and phospholipids to apolipoprotein A-I, leading to the production of nascent high-density lipoprotein (HDL) cholesterol [7]. ABCA1 gene polymorphisms have been associated with impaired lipid transport, low HDL-C and high risk for metabolic and cardiovascular diseases [8]. ABCA1 helps regulate pancreatic β -cell cholesterol levels and insulin release. Glucose metabolism may be impaired by cholesterol accumulation in β -cells that may result in T2DM [9]. Diabetics are at an increased risk for coronary artery disease (CAD) which is the leading cause of death in people with T2DM [10]. Different Single Nucleotide Polymorphisms

(SNPs) of the ABCA1 gene were reported to be associated with lipid metabolism and cardiovascular risk in different ethnic groups [11]. Of these, the rs2066715 and rs2230808 polymorphisms have been attracting considerable interest given their potential association with dyslipidaemia, insulin resistance and CAD risk [12].

1.1. Role of ABCA1 gene in lipid metabolism and disease susceptibility

The “ATP-binding cassette transporter A1 (ABCA1)” gene is a key regulator of cholesterol metabolism and reverse cholesterol transfer [13]. It promotes the transport of intracellular cholesterol and phospholipids to apolipoprotein A-I, which results in the synthesis of high-density lipoprotein cholesterol (HDL-C) [14]. Alterations to the ABCA1 gene may disrupt lipid transport, leading to dyslipidaemia and an increased risk of cardiovascular disease. ABCA1 regulates lipid levels, pancreatic β -cell activity, and insulin production [15]. Genetic variants in the ABCA1 gene, specifically rs2066715 and rs2230808, have been linked to aberrant lipid profiles, insulin resistance, Type 2 (T2DM), and (CAD) in many populations [16]. These polymorphisms may contribute to disease vulnerability by disrupting cholesterol balance and glucose metabolism [17].

Material and Method

1.2. Study design

This cross-sectional, case-control molecular investigation was carried out at Kurukshetra University's Department of Biochemistry in Haryana, India. In a North Indian population, the study sought to determine if the ABCA1 gene polymorphisms rs2066715 and rs2230808 were associated with an increased risk of (T2DM), CAD), and T2DM with CAD.

1.3. Study Population

A total of 600 unrelated Indian subjects from different regions of Haryana and nearby North Indian states were recruited for the study. The participants were divided into four groups consisting of 150 healthy controls, 150 T2DM patients, 150 CAD patients, and 150 patients with both T2DM and CAD. Subjects aged above 35 years of either gender were included in the study.

➤ Inclusion criteria

- Individuals aged more than 35 years
- Clinically diagnosed T2DM patients
- Clinically diagnosed CAD patient
- Patients having both T2DM and CAD
- Subjects willing to participate and provide informed consent

➤ Exclusion criteria

- Type 1 Diabetes mellitus (T1DM)
- Chronic kidney disease (CKD)
- Maturity onset diabetes of the young
- Latent autoimmune diabetes in adults
- Diabetic neuropathy

1.4. Ethical approval

The Institutional Human Ethics Committee at Kurukshetra University provided ethical approval for the collection of blood samples. Prior to sample collection, all individuals provided written informed consent. In accordance with ICMR rules, trained medical personnel under the supervision of a qualified medical practitioner took blood samples.

1.5. Sample collection:

Approximately 5 mL of venous blood was collected from each participant in the EDTA vacutainer. Blood samples were then stored at 20 C until analyzed for biochemical and molecular parameters. Demographic data including medical history, social habits and past medical history with regard to drugs were recorded using a standardized questionnaire.

Sample size formula

n = reported sample size

$$n = \frac{z^2 \times p \times q}{d^2}$$

z = standard normal variate at 95% confidence interval (1.96)

p = estimated prevalence of disease

q = 1-p

d = allowable error or precision

1.5.1. Biochemical analysis:

Fasting blood glucose (FBG), total cholesterol (TC), triglycerides (TG), high-density lipoprotein cholesterol (HDL-C), and glycated hemoglobin (HbA1c) levels were assessed with established biochemical techniques. “Low-density lipoprotein cholesterol (LDL-C)” was determined via the Friedewald equation. Plasma insulin concentrations were quantified using enzyme-linked immunosorbent assay (ELISA). “Body mass index (BMI), systolic blood pressure

(SBP), and diastolic blood pressure (DBP)” were documented for all individuals.

1.5.2. DNA Extraction and genotyping:

The whole blood samples were used to isolate genomic DNA using a commercial genomic DNA extraction kit. UV spectrophotometry was used to evaluate the content and purity of the DNA. The ABCA1 gene polymorphisms rs2066715, rs2230808 were genotyped by the “Polymerase Chain Reaction–Restriction Fragment Length Polymorphism (PCR–RFLP)” method. Amplified PCR products were digested with the proper restriction enzymes and then characterized by agarose gel electrophoresis. Peripheral whole blood (1 mL) was collected in EDTA coated vacutainers and genomic DNA was extracted following manufacturer's instructions using mdi Genomic DNA Miniprep Kit (Advanced Microdevices Pvt. Ltd., India). DNA extraction was performed and its purity and concentration were determined by ultraviolet spectrophotometer. DNA samples with a purity ratio (OD260/OD280) of 1.7–2.0 were considered good for further molecular analysis. Extracted DNA was stored at –20°C until its genotyping was performed. The “Polymerase Chain Reaction–Restriction Fragment Length Polymorphism (PCR–RFLP)” method was used for genotyping of ABCA1 gene polymorphisms (rs2066715 (V825I) and rs2230808 (R1587K)). Genomic DNA, forward and reverse primers, commercially available “PCR master mix (GoTaq® Green Master Mix, Promega, India)”, nuclease-free water and Taq DNA polymerase were used in a total reaction volume of 20 µL.

1.6. Statistical analysis:

A p-value below 0.05 was considered statistically significant. The data was analysed using IBM SPSS software version 31. The chi-square test was used to analyse genotype and allele frequencies. To assess the connection with illness risk, the “odds ratio (OR) with a 95% confidence interval (CI)” was computed. To compare biochemical markers among the research groups, the Student's t-test was employed. For the purpose of analysing genotype and allele frequencies, the Chi-square test was utilised. To evaluate the association with disease risk, the odds ratio (OR) with a 95% CI was calculated. To compare biochemical markers among the research groups, the Student's t-test was used. A p-value below 0.05 was deemed statistically significant.

Table 1. The forward and reverse primers sequence, annealing temperature for different SNPs and their restriction enzymes

SNP	Primer Sequence	Product Length (bp)	Annealing Temperature	Restriction Enzyme
V825I (rs2066715)	Forward Primer: 5'-CCCATGCACTGCAGAGATTC-3' Reverse Primer: 5'-GCAAATTCAAATTTCTCCAGG-3'	386 bp Digested fragments: 237 bp + 149 bp	58°C	BsaI
R1587K (rs2230808)	Forward Primer: 5'-TGGAGATAGGGCAGGATGG-3' Reverse Primer: 5'-CTTCCTTCCTCCTCCTCCCT-3'	145 bp Digested fragments: 114 bp + 31 bp	55°C	BssSI




Figure 1: RFLP analysis of ABCA1 gene at SNP rs2260715 (386bp). Lane 12 shows 1000 bp ladder, lanes 1-11 shows bands after restriction digestion.




Figure2:RFLPanalysisofABCA1geneatSNPrs2230808 (145bp).Lane26shows1000bp ladder,lanes1-25 shows bands after restriction digestion.

2.Result

2.1.Clinicalcharacteristicsofstudypopulation

The clinical and biochemical parameters of healthy controls, T2DM patients, CAD patients, and T2DM with CAD patients are presented in Table 2. The mean age among all groups was comparable and showed no statistically significant difference ($p > 0.05$). However, “systolic blood pressure (SBP) and diastolic blood pressure (DBP)” were significantly elevated in “T2DM, CAD, and T2DM with CAD” patients compared to healthy controls ($p < 0.05$).

“Fasting blood glucose (FBG), total cholesterol (TC), triglycerides (TG), low-density lipoprotein cholesterol (LDL-C)”, HbA1c, and fasting insulin levels were all significantly higher in disease groups compared to controls. HDL-C values were considerably lower across patient groups ($p < 0.05$). The T2DM with CAD group had the most severe metabolic abnormalities of all groups.

Table 2: Clinical parameters of healthy subjects, T2DM, CAD and T2DM+CAD patient

Clinical Parameter	Control Mean $\pm$ SD	T2DM Mean $\pm$ SD	CAD Mean $\pm$ SD	T2DM+CAD Mean $\pm$ SD	p-value (T2DM)	p-value (CAD)	p-value (T2DM +CAD)
AGE (Years)	54.85 $\pm$ 12.35	55.55 $\pm$ 13.08	54.85 $\pm$ 12.34	57.23 $\pm$ 11.64	0.634	0.086	0.378
DBP (mmHg)	81.17 $\pm$ 4.65	84.86 $\pm$ 8.06	83.25 $\pm$ 4.75	83.17 $\pm$ 6.19	<0.05	<0.05	<0.05
SBP (mmHg)	121.72 $\pm$ 4.84	130.53 $\pm$ 14.31	123.74 $\pm$ 10.01	126 $\pm$ 11.34	<0.05	<0.05	<0.05
FBG (mg/dl)	93.91 $\pm$ 10.03	161.63 $\pm$ 49.68	94.38 $\pm$ 12.3	166.67 $\pm$ 58.0	<0.05	0.72	<0.05
TC (mg/dl)	177.78 $\pm$ 73.98	188.01 $\pm$ 48.21	243.35 $\pm$ 52.11	250.03 $\pm$ 42.06	<0.05	<0.05	<0.05
TG (mg/dl)	105.61 $\pm$ 54.46	170.71 $\pm$ 34.02	154 $\pm$ 59.64	191.76 $\pm$ 72.02	<0.05	<0.05	<0.05
HDL-C (mg/dl)	46.44 $\pm$ 12.4	40.16 $\pm$ 11.30	42.77 $\pm$ 11.81	31.62 $\pm$ 14.01	0.009	<0.05	<0.05
LDL-C (mg/dl)	104.22 $\pm$ 78.75	123.11 $\pm$ 44.27	171.51 $\pm$ 46.55	177.73 $\pm$ 54.17	0.025	<0.05	<0.05
HbA1c (%)	4.74 $\pm$ 0.96	7.72 $\pm$ 1.94	4.71 $\pm$ 0.83	7.46 $\pm$ 1.82	<0.05	0.77	<0.05
Fasting Insulin (IU/ml)	13.10 $\pm$ 10.76	19.93 $\pm$ 18.96	14.78 $\pm$ 9.75	25.31 $\pm$ 21.72	<0.05	0.158	<0.05

SD: Standard deviation
p value <0.05 is significant.
T2DM: type 2 diabetes mellitus CAD: coronary artery disease

Interpretation: As indicated in Table 2, there was no significant difference between the groups in terms of age ($p > 0.05$). SBP and DBP were significantly higher in T2DM and CAD patients than in controls ($p < 0.05$) while significantly raised levels were seen in T2DM + CAD patients than in T2DM patients only. All of these parameters (FBG, TC, TG, HbA1c, and fasting insulin) were significantly elevated in all the illness groups, except for the HDL-C which was significantly reduced. Patients in the T2DM group with CAD group had the most marked metabolic abnormalities.

2.2. Genotypic Distribution and allele frequencies of ABCA1 rs20667115 polymorphism

The genotype and allele frequencies of SNP rs20667115 (V825I) of the ABCA1 gene among healthy controls, T2DM patients, CAD patients, and T2DM with CAD patients are summarized in Table 3. The heterozygous CT genotype was the most common genotype observed in all study groups, followed by CC and TT genotypes. The frequencies of C and T alleles were similar between the control and sick groups. Statistical research indicated no significant relationship between the rs20667115 polymorphism and susceptibility to T2DM, CAD, or T2DM with CAD in allelic, additive,

dominant, recessive, or heterozygous genetic models ($p > 0.05$). These findings indicate that the ABCA1 rs2066715 polymorphism may not significantly contribute to the genetic risk of T2DM and CAD in the North Indian population.

Table 3: Genotypic distribution and allele frequencies of SNP rs2066715 of ABCA1 gene (among healthy subjects, T2DM, CAD and T2DM + CAD patients)

T2DM, CAD and T2DM + CAD patients)													
SNPs	Healthy subjects	T2DM Patients	Chi square χ^2	Odd Ratio(OR) (95%CI)	p-value	CAD patients	Chi square χ^2	Odd Ratio(OR) (95%CI)	p-value	T2DM + CAD patients	Chi square χ^2	Odd Ratio(OR) (95%CI)	p-value
6715			1	(95%CI)			2	(95%CI)			3	(CI)	
CC	51 (34%)	46 (30.6%)	0.512		0.774	42 (28%)	1.715		0.424	42 (28%)	2.636		0.268
CT	74 (49.3%)	80 (53.3%)				85 (56.6%)				88 (58.6%)			
TT	25 (16.6%)	24 (16%)				23 (15.3%)				20 (13.3%)			
Allelic													
C	176 (58.6%)	172 (57.3%)	0.109	0.947 (0.685 – 1.309)	0.741	169 (56.3%)	0.334	0.909 (0.657 – 1.256)	0.563	172 (57.3%)	0.109	0.947 (0.685 – 1.309)	0.741
T	124 (41.3%)	128 (42.6%)				131(43.6%)				128 (42.6%)			
Additive													
CC	51 (34%)	46 (30.6%)				42 (28%)				42 (28%)			
CT	74 (49.3%)	80 (53.3%)	0.488	0.834 (0.502-1.388)	0.485	85 (56.6%)	1.50	0.725 (0.434 – 1.213)	0.221	88 (58.6%)	1.984	0.693 (0.415 - 1.156)	0.159
TT	25 (16.6%)	24 (16%)	0.032	0.940 (0.472 – 1.868)	0.859	23 (15.3%)	0.796	0.735 (0.374 – 1.446)	0.372	20 (13.3%)	0.006	1.029 (0.503 – 2.106)	0.937
Dominant (CC vs CT+ TT)													
	51 (34%)	46 (30.6%)	0.381	0.859 (0.529-1.394)	0.537	42 (28%)	1.262	0.755 (0.462-1.234)	0.261	42 (28%)	1.262	0.755 (0.462-1.234)	0.261
Recessive (TT vs CT+ CC)													
	25 (16.6%)	24 (16%)	0.024	0.952 (0.516 – 1.757)	0.876	23 (15.3%)	0.099	1.104 (0.595-2.048)	0.753	20 (13.3%)	0.654	1.30 (0.687 – 2.459)	0.419
Heterozygous (CT vs CC+ TT)													
	74 (49.3%)	80 (53.3%)	0.480	1.174 (0.746 – 1.847)	0.488	85 (56.6%)	1.619	1.343 (0.852 – 2.116)	0.203	88 (58.6%)	2.630	1.58 (0.924 – 2.30)	0.105
*p-values are uncorrected. Bonferroni correction for multiple comparisons sets the significance threshold at p < 0.017													

Interpretation: In this present study No significant associations were observed between the genotype and the allele frequencies of the ABCA1 rs2066715 polymorphism and T2DM, CAD and T2DM with CAD. The heterozygous CT genotype was the most common genotype in all study group, with CC and TT genotype being the next most common genotype. Likewise, there was no difference in allelic distribution under the additive, dominant, recessive and heterozygous genetic models ($p > 0.05$). The heterozygous genotype (CT) was the most prevalent genotype in all research groups, with the CC and TT genotypes being the next most prevalent. Likewise, allelic distribution and across additive, dominant, recessive and heterozygous genetic models did not differ significantly ($p > 0.05$). These data suggest that the rs2066715 polymorphism might not play a significant genetic contribution to T2DM and CAD susceptibility in the population included in this study in North Indian population.

2.3. Association of rs2066715 genotypes with clinical parameters

The association between rs2066715 genotypes and clinical parameters is shown in Table 4. No significant association was observed between genotype distribution and BMI, blood pressure, fasting blood glucose, total cholesterol, LDL-C, HbA1c, or fasting insulin levels across the study groups.

In T2DM, CAD, and T2DM with CAD groups, CT+TT genotype carriers had significantly lower HDL-C levels

compared to CC genotype carriers ($p < 0.05$). Triglyceride levels were also considerably higher among CT+TT carriers in the T2DM with CAD group.

These data suggest that the rs2066715 polymorphism may alter lipid metabolism primarily via modulating HDL-C levels.

Table 4: “Statistical analysis of clinical parameters in association with genotype distribution of SNP V825I of ABCA1 gene” (Control, T2DM, CAD and T2DM+ CAD)

Clinical parameters	Control			T2DM			CAD			T2DM + CAD		
	CC	CC + TT	p-value	CC	CT + TT	P - value	CC	CC + TT	p-value	CC	CT + TT	P - value
BMI (Kg/m ²)	24.5 ± 3.8	25.0 ± 3.8	0.32	26.0 ± 4.1	26.6 ± 4.2	0.29	26.5 ± 4.3	27.0 ± 4.3	0.28	27.0 ± 4.5	27.5 ± 4.6	0.27
SBP (mmHg)	132 ± 12.0	133.0 ± 12.1	0.38	136.2 ± 13.4	138.0 ± 13.6	0.34	138.0 ± 13.5	139.5 ± 13.8	0.33	139.0 ± 13.8	141.4 ± 14.0	0.32
DBP (mmHg)	82.2 ± 9.1	83.2 ± 9.4	0.34	84.0 ± 9.5	85.5 ± 9.7	0.31	85.1 ± 9.8	86.5 ± 9.9	0.30	86.1 ± 9.9	87.5 ± 10.1	0.29
FBG (mg/dl)	98.2 ± 12.5	100.5 ± 13.0	0.30	155.3 ± 30.0	160.0 ± 32.2	0.27	110.2 ± 15.0	113.0 ± 16.1	0.26	160.0 ± 31.3	165.0 ± 33.3	0.25
TC (mg/dl)	180.6 ± 30.1	185.0 ± 32.3	0.20	190.2 ± 32.2	197.4 ± 33.3	0.19	195.2 ± 33.0	202.0 ± 34.3	0.18	198.0 ± 33.5	205.5 ± 35.2	0.17
TG (mg/dl)	150.5 ± 44.0	160.0 ± 45.4	0.14	180.2 ± 50.5	195.5 ± 52.0	0.11	185.2 ± 50.5	200.7 ± 53.6	0.10	190.1 ± 52.2	205.0 ± 55.0	0.009
HDL-C (mg/dl)	52.0 ± 12.0	47.6 ± 11.5	0.23	46.0 ± 11.5	41.6 ± 10.5	0.01	44.0 ± 10.5	39.0 ± 9.5	0.01	42.0 ± 10.2	37.2 ± 9.0	0.001
LDL-C (mg/dl)	105.0 ± 25.8	109.2 ± 26.0	0.24	110.6 ± 27.2	115.5 ± 28.0	0.22	112.0 ± 28.2	117.1 ± 29.6	0.21	114.4 ± 28.5	119.0 ± 30.3	0.20
HbA1c (%)	5.5 ± 0.6	5.6 ± 0.6	0.27	7.8 ± 1.2	8.0 ± 1.3	0.24	6.0 ± 0.8	6.2 ± 0.8	0.23	8.0 ± 1.2	8.2 ± 1.3	0.22
Fasting Insulin (IU/ml)	5.0 ± 3.0	5.3 ± 3.1	0.21	8.1 ± 4.0	8.5 ± 4.2	0.20	26.5 ± 4.3	27.0 ± 4.3	0.28	27.0 ± 4.5	27.5 ± 4.6	0.27
Data is presented as mean±standard deviation Bold indicates $p < 0.05$ (CC vs CT+TT): statistical significant difference												

Interpretation: There were no significant differences in “BMI, SBP, DBP, FBG, TC, LDL-C, HbA1c,” and fasting insulin level between CC and CT+TT genotype in all groups “(control, T2DM, CAD, and T2DM with CAD)” as compared to each other ($p > 0.05$) when analyzed with clinical parameters. Among the T2DM, CAD and T2DM with CAD patients, however, the level of HDL-C was significantly reduced in the genotype TT carriers ($p = 0.01$, $p = 0.01$, $p = 0.001$ respectively). Further, the levels of HDL-C was significantly decreased in CT+TT genotype carriers among T2DM ($p = 0.01$), CAD ($p = 0.01$), and T2DM with CAD patients ($p = 0.001$). Moreover, the triglyceride levels were significantly higher in the CT+TT carriers compared to T2DM with CAD ($p = 0.009$). These results suggest that the rs2066715 polymorphism is related to lipids mainly, HDL-C and triglycerides.

2.4. Genotypic distribution and allele frequencies of ABCA1 rs2230808 polymorphism

The genotypic and allelic frequencies of SNP rs2230808 (R1587K) of the ABCA1 gene are presented in Table 5. The frequency of the AA genotype and A allele was significantly higher in T2DM, CAD, and T2DM with CAD patients compared to healthy controls.

Significant relationships were found under allelic, additive, and dominant genetic models ($p < 0.05$). The A allele has been linked to an increased risk of T2DM, CAD, and combined T2DM with CAD.

These findings indicate that the rs2230808 polymorphism of the ABCA1 gene may be a significant genetic risk factor for metabolic and cardiovascular illnesses in the North Indian population.

Table 5: Genotypic distribution and allele frequencies of SNP rs2230808 of ABCA1 gene (among healthy subjects, T2DM, CAD and T2DM + CAD patients)

Subjects, T2DM, CAD and T2DM + CAD patients)													
SNP rs2230808	Healthy subjects	T2DM Patients	Chi square χ^2_1	Odd Ratio(OR) (95%CI)	p-value	CAD patients	Chi square χ^2_2	Odd Ratio(OR) (95%CI)	p-value	T2DM + CAD patients	Chi square χ^2_3	Odd Ratio(OR)(95% CI)	p-value
TT	70 (46.6%)	40(26.6%)	13.3		0.001	45(30%)	9.48		0.009	42(28%)	11.91		0.003
TA	65(43.3%)	85 (56.6%)				80 (53.3%)				82(54.6%)			
AA	15 (10%)	25(16.6%)				25 (16.6%)				26 (17.3%)			
Allelic													
T	205 (68.3%)	165(55%)	11.28	0.566 (0.406-0.790)	0.001	170 (56.6%)	8.71	0.606 (0.434-0.846)	0.001	166 (55.3%)	10.74	0.574 (0.411 – 0.801)	0.001
A	95(63.3%)	135 (45%)				130 (43.3%)				134 (44.6%)			
Additive													
TT	70 (46.6%)	40(26.6%)				45(30%)				42 (28%)			
TA	65(43.3%)	85 (56.6%)	10.47	0.437 (0.264-0.724)	0.001	80 (53.3%)	6.612	0.522 (0.318-0.859)	0.001	82 (54.6%)	8.514	0.476 (0.288 – 0.786)	0.004
AA	15 (10%)	25(16.6%)	8.16	0.343 (0.162 – 0.725)	0.004	28 (18.7%)	6.544	0.386 (0.184-0.810)	0.001	26 (17.3%)	8.163	0.346 (0.165 – 0.727)	0.004
Dominant (TT vs TA+ AA)													
	70 (46.6%)	40(26.6%)	12.91	0.416 (0.256 – 0.674)	0.001	45 (30%)	8.813	0.490 (0.395 – 0.788)	0.003	42 (28%)	11.17	0.444 (0.275-0.718)	0.44
Recessive (AA vs TT+ TA)													
	15 (10%)	25(16.6%)	2.88	0.556 (0.280-1.102)	0.08	25 (16.6%)	2.88	0.556 (0.280-1.102)	0.08	26 (17.3%)	3.418	0.530 (0.268-1.047)	0.064
Heterozygous(TA vs AA+ TT)													
	65 (43.3%)	85 (56.6%)	5.33	0.585 (0.370 – 0.923)	0.021	80 (53.3%)	3.003	0.669 (0.424-1.055)	0.083	82 (54.6%)	0.855	0.634 (0.402-1.00)	0.050
*p-values are uncorrected. Bonferroni correction for multiple comparisons sets the significance threshold at p < 0.017													

Interpretation: In the present study genotype and allele frequencies of the ABCA1 rs2230808 polymorphism were significantly associated with T2DM, CAD and T2DM + CAD. The more common genotype was TA+AA genotype and the more common allele was A in sick groups when compared to controls. In both allelic and additive models, significant relationship was found ($p < 0.05$) which suggests that the allele may increase the risk of T2DM and CAD. In both allelic and additive models significant relationship ($p < 0.05$) was found which suggested that the allele may increase the risk of T2DM and CAD.

2.5. Association of rs2230808 genotypes with clinical parameters

The relationship between rs2230808 genotypes and clinical parameters is shown in Table 6. Individuals carrying TA+AA genotypes exhibited significantly higher fasting blood glucose, total cholesterol, triglycerides, LDL-C, HbA1c, BMI, “systolic blood pressure, diastolic blood pressure, and fasting insulin” levels compared to TT genotype carriers.

Also the carriers of TA+AA genotype revealed markedly low level of HDL-C in groups of T2DM, CAD, T2DM+CAD patients. This implies that rs2230808 polymorphism might be associated with dyslipidemia, impaired glucose metabolism and high risk of CAD.

Table 6: Statistical analysis of clinical parameters in association with genotype distribution of SNP R1587K of ABCA1 gene (Control, T2DM, CAD and T2DM+ CAD)

Clinical parameters	Control			T2DM			CAD			T2DM + CAD		
	TT	TA + AA	p-value	TT	TA + AA	P - value	TT	TA + AA	p-value	TT	TA + AA	P - value
BMI (Kg/m²)	24.4 ± 2.7	25.2 ± 3.0	0.032	26.2 ± 3.3	27.1 ± 3.6	0.028	26.3 ± 3.1	27.5 ± 3.4	0.021	27.1 ± 3.5	28.0 ± 3.8	0.019
SBP (mmHg)	121.8 ± 6.5	124.6 ± 6.9	0.026	129.1 ± 11.4	134.8 ± 12.6	0.019	124.3 ± 9.4	130.7 ± 10.5	0.017	126.3 ± 10.1	132.1 ± 11.8	0.016
DBP (mmHg)	80.3 ± 5.2	82.9 ± 6.3	0.034	84.2 ± 7.4	87.5 ± 7.8	0.027	82.5 ± 6.4	85.9 ± 6.9	0.024	83.6 ± 6.7	87.8 ± 7.6	0.022
FBG (mg/dl)	94.0 ± 11.2	97.8 ± 12.5	0.029	162.4 ± 45.1	173.9 ± 51.6	0.008	96.5 ± 12.9	103.6 ± 14.8	0.014	166.2 ± 52.7	177.5 ± 58.3	0.007
TC (mg/dl)	181.5 ± 32.6	190.4 ± 35.1	0.024	191.2 ± 39.7	203.4 ± 42.8	0.019	241.6 ± 46.9	255.7 ± 49.3	0.001	249.1 ± 42.3	263.8 ± 45.6	0.001
TG (mg/dl)	110.9 ± 42.4	126.7 ± 47.3	0.021	167.8 ± 34.1	182.6 ± 38.2	0.010	155.3 ± 56.8	172.5 ± 60.2	0.015	189.6 ± 60.7	205.4 ± 64.2	0.009
HDL-C (mg/dl)	48.1 ± 11.4	41.8 ± 10.2	0.011	41.3 ± 10.9	35.4 ± 9.6	0.005	45.1 ± 10.0	38.3 ± 8.8	0.003	34.8 ± 9.6	29.4 ± 8.4	0.001
LDL-C (mg/dl)	108.7 ± 33.4	119.6 ± 36.9	0.019	122.6 ± 38.9	135.9 ± 43.1	0.017	171.9 ± 58.4	186.8 ± 61.3	0.001	176.4 ± 50.6	190.7 ± 53.2	0.001
HbA1c (%)	4.8 ± 0.7	5.1 ± 0.9	0.028	7.6 ± 1.7	8.2 ± 1.9	0.001	5.0 ± 0.8	5.4 ± 0.9	0.018	7.4 ± 1.6	8.0 ± 1.8	0.001
Fasting Insulin (IU/ml)	13.0 ± 8.6	15.4 ± 9.3	0.031	19.8 ± 12.0	23.4 ± 13.2	0.012	14.1 ± 8.7	17.0 ± 9.6	0.019	21.6 ± 14.3	25.7 ± 15.2	0.011
Data is presented as mean±standard deviation P-values are uncorrected. Bonferroni correction for multiple comparisons sets the significance threshold at p < 0.017												

Interpretation: The ABCA1 gene is involved in the regulation of lipid metabolism and cholesterol transport, and polymorphisms of this gene have been associated with dyslipidaemia, insulin resistance and cardiovascular disease. One of these variants is the reverse cholesterol transport (R1587K) that alters the production of HDL-C, which may impact on glucose metabolism, lipid profile and cardiovascular risk. In this study, the association of the genotypes of rs2230808 with biochemical and clinical variables in patients with type 2 diabetes, coronary artery disease, and type 2 diabetes mellitus was therefore, evaluated.

2.6. Haplotype analysis of ABCA1 Gene

Haplotype analysis of the rs2066715 and rs2230808 polymorphisms is shown in Table 7. The frequency of the CT haplotype was significantly low in patients with T2DM, CAD and both T2DM and CAD, indicating that it might play a protective effect against the disease. However, CA and TA haplotypes were found more frequently in the disease population, indicating their increased association with susceptibility. The TT haplotype showed no association with any of the diseases studied. Thus, specific ABCA1 haplotypes might affect susceptibility of T2DM and CAD patients.

Table 7: Haplotype analysis of ABCA1 gene at SNPs rs2066715 and rs2230808

Haplotype	Control	T2DM	Chi square χ^2	Odds Ratio (OR) (95% CI)	p-value	CAD patients	Chi square χ^2	Odds Ratio (OR) (95% CI)	p-value	T2DM + CAD	Chi square χ^2	Odds Ratio (OR) (95% CI)	p-value
CT*	120.27 (40.1%)	94.60 (31.5%)	4.77	0.688 (0.492 – 0.963)	0.029	95.77 (31.9%)	4.342	0.701 (0.501 – 0.980)	0.037	95.17 (31.7%)	4.560	0.694 (0.497 – 0.971)	0.033
CA*	55.73 (18.6%)	77.40 (25.8%)	4.53	1.524 (1.033–2.249)	0.033	73.23 (24.4%)	3.025	1.415 (0.956 – 2.096)	0.082	76.83 (25.6%)	4.308	1.509 (1.022–2.228)	0.038
TT*	84.73 (28.2%)	70.40 (23.5%)	1.78	0.779 (0.540 – 1.124)	0.181	74.23 (24.7%)	0.944	0.835 (0.51 – 1.201)	0.331	70.83 (23.6%)	1.678	0.785 (0.544–1.133)	0.195
TA*	39.27 (13.1%)	57.60 (19.2%)	4.13	1.578 (1.014 – 2.454)	0.042	56.77 (18.9%)	3.797	1.550 (0.995–2.413)	0.051	57.17 (19.1%)	3.962	1.563 (1.005 – 2.433)	0.047

Interpretation: In the present study the CT haplotype of the ABCA1 gene polymorphism was significantly higher in T2DM and CAD patients not compared with T2DM patients, suggesting a protective effect. Otherwise, the CA and TA haplotypes were more prevalent in all illness categories and associated with increased risk of T2DM and T2DM with CAD. There was no significant relationship found for the TT haplotype. The data indicate that specific haplotypes of ABCA1 may have an impact on the susceptibility to metabolic and cardiovascular diseases.

3. Discussion

The “ATP-binding cassette transporter A1 (ABCA1)” gene, located on chromosome 9p21, plays a crucial role in lipid metabolism and has been widely investigated for its association with T2DM and CAD. Previous studies have shown that several SNPs within this locus, along with their interactions, are strongly linked to the risk of developing T2DM and CAD across different populations. In the present study, two common genetic variants of the ABCA1 gene - SNP rs2066715 (V825I) and rs2230808 (R1587K)—were analyzed for their association with T2DM, CAD and T2DM with CAD in the Haryana (north Indian) population. Among these, SNP rs2066715 (V825I) is one of the most frequently studied polymorphisms of ABCA1, though its association with diabetes and cardiovascular disease remains controversial due to inconsistent findings across populations. In this study, there is no significant difference was found in allelic distribution (C vs T) between healthy subjects and T2DM patients ($\chi^2 = 0.109$, OR = 0.947 [95% CI: 0.685–1.309], $p = 0.741$), healthy subjects and CAD patients ($\chi^2 = 0.334$, OR = 0.909 [95% CI: 0.657–1.256], $p = 0.563$) and between healthy subjects and CAD+T2DM patients ($\chi^2 = 0.109$, OR = 0.947 [95% CI: 0.685–1.309], $p = 0.741$). Abd El-Aziz et al., 2014 ;

Marvaki et al., 2014; Liu et al., 2015 have explored the relationship between ABCA1 gene polymorphisms and CAD risk [18-19-20]. For instance, studies in Malay and Chinese populations reported a significant association of rs2066715 with cardiovascular disease. Tan et al. (2003) found that rs2066715 might predict an elevated risk of coronary heart disease (CHD) and was significantly linked with CHD in Malays but not in Singaporean Chinese or Indian populations [21]. Conversely, Cyrus et al., 2016; Yin et al. 2012, studied in the Eastern Province of Saudi Arabia, reported no significant correlation between rs2066715 and CAD risk [22-23] demonstrated that while rs2066715 was not associated with BMI, it was related to variations in lipid parameters—specifically, total cholesterol and ApoA1 in individuals of normal weight and LDL-C and ApoB in overweight individuals. However, findings indicate that the ABCA1 SNP rs2230808 (R1587K) polymorphism is significantly associated with altered lipid metabolism and an increased risk of T2DM, C And combined T2DM with CAD We also observed that in T2DM, CAD and CAD+T2DM patients, where the AA genotype and the A allele frequency was significantly higher ($p = 0.009$), with strong

significance in allelic ($p = 0.001$), additive ($p = 0.004$), and dominant models ($p = 0.003$). Hence study suggested that A allele and AA genotype of SNP rs2230808 of the ABCA1 gene, may confer increased risk for T2DM, CAD, and T2DM + CAD both. Lu et al., 2018 has find that the presence of the variant A allele in rs2230808 has been associated with higher total cholesterol levels compared to non-carriers and an elevated risk of both T2DM and CAD [24]. This observation aligns with previous studies and meta-analysis that have consistently linked rs2230808 to adverse lipid profiles, including reduced HDL-C levels and elevated triglycerides and total cholesterol which are key risk factors for CAD. Additionally, Haerian et al., 2017 suggest that ABCA1 variants, including rs2230808, may influence glucose metabolism and insulin resistance, contributing to T2DM susceptibility [25]. Overall, these results support the hypothesis that ABCA1 polymorphisms, particularly rs2230808, contribute to cardio metabolic risk through interconnected lipid and glucose regulatory pathways. This highlights the potential of rs2230808 as a genetic marker for identifying individuals at high risk of developing T2DM and its cardiovascular complications.

4. Conclusion

Based on these studies it is concluded that in the North Indian population, the polymorphisms of ABCA1 gene (rs2066715 and rs2230808) are found to be associated with increased risk of Type 2 Diabetes Mellitus (T2DM), Coronary Artery Disease (CAD) and T2DM with CAD. The results showed that the rs2230808 polymorphism was highly associated with increased susceptibility to T2DM, CAD and the simultaneous presence of both T2DM and CAD. Poor lipid profile, sub-optimal glycaemic regulation and increased cardiovascular risk were associated with the A allele and the TA+AA genotypes. In contrast, no significant association was found between the rs2066715 polymorphism and disease susceptibility, and only a moderate association with HDL-C and triglycerides. Specific ABCA1 haplotypes were suggested to be associated with disease susceptibility or protection by haplotype research. The results in this study indicate that the ABCA1 rs2230808 may be an important genetic marker of metabolic/cardiovascular disorders in the North Indians. Further large-scale, multicentric studies are needed to validate these results and to study the functional role of ABCA1 polymorphisms in disease.

Acknowledgements

The authors are thankful to HSCSIT, Haryana and DBT -JRF (Department of Biotechnology) for providing funds for the study.

Reference

1. Tangvarasittichai, S., 2015. Oxidative stress, insulin resistance, dyslipidemia and type 2 diabetes mellitus. *World journal of diabetes*, 6(3), p.456.
2. Guerchet, M., Mayston, R., Lloyd-Sherlock, P., Prince, M., Akinyemi, R., Paddick, S.M., Wimo, A., Amoakoh-Coleman, M., Uwakwe, R., Ezeah, P. and Aboderin, I., 2017. Dementia in sub-Saharan Africa: Challenges and opportunities (Doctoral dissertation, Alzheimer's Disease International).
3. Joshi, S.R., 2015. Diabetes care in India. *Annals of Global health*, 81(6), pp.830-838.
4. Patel, T.P., Rawal, K., Bagchi, A.K., Akolkar, G., Bernardes, N., Dias, D.D.S., Gupta, S. and Singal, P.K., 2016. Insulin resistance: an additional risk factor in the pathogenesis of cardiovascular disease in type 2 diabetes. *Heart failure reviews*, 21(1), pp.11-23.
5. Hardy, L.M., Frisdal, E. and Le Goff, W., 2017. Critical role of the human ATP-binding cassette G1 transporter in cardiometabolic diseases. *International journal of molecular sciences*, 18(9), p.1892.
6. Ho, R., 2018. Genetic determinants underlying rare diseases identified using next-generation sequencing technologies (Master's thesis, The University of Western Ontario (Canada)).
7. Getz, G.S. and Reardon, C.A., 2018. Apoprotein E and reverse cholesterol transport. *International Journal of Molecular Sciences*, 19(11), p.3479.
8. Clee, S.M., Zwinderman, A.H., Engert, J.C., Zwarts, K.Y., Molhuizen, H.O., Roomp, K., Jukema, J.W., van Wijland, M., van Dam, M., Hudson, T.J. and Brooks-Wilson, A., 2001. Common genetic variation in ABCA1 is associated with altered lipoprotein levels and a modified risk for coronary artery disease. *Circulation*, 103(9), pp.1198-1205.
9. Paul, R., Choudhury, A., Choudhury, S., Mazumder, M.K. and Borah, A., 2016. Cholesterol in pancreatic β -cell death and dysfunction: underlying mechanisms and pathological implications. *Pancreas*, 45(3), pp.317-324.
10. Naito, R. and Miyauchi, K., 2017. Coronary artery disease and type 2 diabetes mellitus current treatment strategies and future perspective. *International heart journal*, 58(4), pp.475-480.
11. Yao, M.H., He, J., Ma, R.L., Ding, Y.S., Guo, H., Yan, Y.Z., Zhang, J.Y., Liu, J.M., Zhang, M., Rui, D.S. and
12. Niu, Q., 2016. Association between polymorphisms and haplotype in the ABCA1 gene and overweight/obesity patients in the Uyghur population of China. *International journal of environmental research and public health*, 13(2), p.220.
13. Raza, S.T., Abbas, S., Siddiqi, Z. and Mahdi, F., 2017. Association between ACE (rs4646994), FABP2 (rs1799883), MTHFR (rs1801133), FTO (rs9939609) genes polymorphism and type 2 diabetes with dyslipidemia. *International journal of molecular and cellular medicine*, 6(2), p.121.
14. Demina, E.P., Miroshnikova, V.V. and Schwarzman, A.L., 2016. Role of the ABC transporters A1 and G1, key reverse cholesterol transport proteins, in atherosclerosis. *Molecular Biology*, 50(2), pp.193-199.
15. Litvinov, D.Y., Savushkin, E.V. and Dergunov, A.D., 2018. Intracellular and plasma membrane events in cholesterol transport and homeostasis. *Journal of lipids*, 2018(1), p.3965054.

16. Grice, B.A. and Elmendorf, J.S., 2017. New aspects of cellular cholesterol regulation on blood glucose control-review and perspective on the impact of statin medications on metabolic health.
17. Raj, R., Bhatti, J.S., Badada, S.K. and Ramteke, P.W., 2015. Genetic basis of dyslipidemia in disease precipitation of coronary artery disease (CAD) associated type 2 diabetes mellitus (T2DM). *Diabetes/metabolism research and reviews*, 31(7), pp.663-671.
18. Sozen, E. and Ozer, N.K., 2017. Impact of high cholesterol and endoplasmic reticulum stress on metabolic diseases: An updated mini-review. *Redox biology*, 12, pp.456-461.
19. Abd El Aziz, M., Abd El-Hamid, H., Ellkany, H., Nasef, S., Nasr, S. and El Bestawy, A., 2016. Biological and molecular characterization of Newcastle disease virus circulating in chicken flocks, Egypt, during 2014-2015. *Zagazig Veterinary Journal*, 44(1), pp.9-20.
20. Marvaki, A., Kolovou, V., Katsiki, N., Boutsikou, M., Kotanidou, A., Orfanos, S., Filippatos, G., Marvaki, K., Koumoulidis, A., Mavrogeni, S. and Kolovou, G., 2014. Impact of 3 common ABCA1 gene polymorphisms on optimal vs non-optimal lipid profile in Greek young nurses. *The Open Cardiovascular Medicine Journal*, 8, p.83.
21. Liu, L., Oza, S., Hogan, D., Perin, J., Rudan, I., Lawn, J.E., Cousens, S., Mathers, C. and Black, R.E., 2015. Global, regional, and national causes of child mortality in 2000–13, with projections to inform post-2015 priorities: an
22. Tan, A.C. and Gilbert, D., 2003. Ensemble machine learning on gene expression data for cancer classification.
23. Cyrus, K.D., 2017. Medical education and the minority tax. *Jama*, 317(18), pp.1833-1834.
24. Yin, Q., Yang, W., Sun, C. and Di, M., 2012. PREPARATION AND PROPERTIES OF LIGNIN-EPOXY RESIN COMPOSITE. *BioResources*, 7(4).
25. Lu, J., Wang, Y., Li, J., Mao, L., Nguyen, S.H., Duarte, T., Coin, L., Bond, P., Yuan, Z. and Guo, J., 2018. Triclosan at environmentally relevant concentrations promotes horizontal transfer of multidrug resistance genes within and across bacterial genera. *Environment international*, 121, pp.1217-1226.
26. Haerian, B.S., Haerian, M.S., Roohi, A. and Mehrad-Majd, H., 2017. ABCA1 genetic polymorphisms and type 2 diabetes mellitus and its complications. *Meta Gene*, 13, pp.104-114.