

Public Awareness of Rheumatoid Arthritis as a Genetically Influenced Autoimmune Disease: A Population-Based Study in Riyadh, Saudi Arabia

Faisal Fayeز Alrashidi¹, Muhannad Khalid Alzahir¹, Sultan Thabet Alluhaibi¹, Abdulelah Abdullah Alansari¹, Faisal Abdulmajeed Aljarboua¹, Ahmed Ibrahim Alkhurayyif¹, Fares Fahad Alammari¹, Ammar Fathi Mohammed Khalifa²

1. Medical student, Al-Maarefa Colleges, Riyadh, Saudi Arabia

2. Assistant Professor. College of Medicine. AlMaarefa University, Saudi Arabia

ABSTRACT

This cross-sectional study assessed public awareness of rheumatoid arthritis (RA) and its risk factors among 537 adults in Riyadh, Saudi Arabia, in 2022. Using a structured questionnaire, the researchers evaluated participants' general knowledge about RA, awareness of signs and symptoms, and lifestyle practices. Results revealed significant knowledge deficits, with 67.6% of participants demonstrating inadequate general knowledge about RA and 58.8% showing inadequate understanding of its signs and symptoms. Physical exercise practices were found to be highly inadequate among 81.2% of participants. Although no significant associations were found between demographic characteristics (gender, education, occupation, nationality, BMI) and general knowledge, a significant relationship was observed between age and knowledge of signs and symptoms ($p=0.025$), with participants over 40 years showing better awareness. The findings highlight the need for enhanced public education programs about RA, particularly targeting individuals in high-risk age groups. This research underscores the importance of developing comprehensive awareness initiatives to improve early recognition, management, and quality of life outcomes for individuals with RA in Saudi Arabia.

Keywords: Rheumatoid arthritis, public awareness, Autoimmune disease, Genetic susceptibility, Knowledge–attitude–practice

INTRODUCTION

Rheumatoid arthritis (RA) is a chronic inflammatory autoimmune disease that primarily affects the joints but can also have systemic manifestations involving various organs of the body. RA is recognized as one of the most debilitating disorders due to its progressive nature, which leads to joint destruction, deformities, and significant impairment in quality of life. The disease is characterized by the infiltration of leukocytes into synovial tissues, joints, and synovial fluid, resulting in persistent inflammation and the eventual breakdown of joint cartilage (Smolen et al., 2016). This progressive joint damage leads to pain, functional disability, and a reduced ability to perform daily activities. Since RA was first identified by Sir Alfred Baring Garrod in 1859, it has been recognized as a chronic disease with no definitive cure. Long-term remission or complete resolution of symptoms is rare, underscoring the importance of early diagnosis, effective management, and increased public awareness of the disease (Alamanos et al., 2006).

The prevalence of RA varies across populations, with an estimated lifetime prevalence of up to 1% globally (Silman & Hochberg, 2001). While this may seem relatively low, the disease has a profound impact on those it affects. RA is ranked as the 42nd highest contributor to global disability, as measured by Disability-Adjusted Life Years (DALYs) (Cross et al., 2014). The disease's burden is more closely associated with its impact on morbidity rather than mortality. Common symptoms such as chronic pain, fatigue, depression, and functional impairment contribute significantly to the reduction in quality of life for RA patients. The World Health Organization (WHO) and the World Bank estimate that RA accounts for approximately 0.8% of all global DALYs and 0.1% of all deaths in Europe. These statistics highlight the critical importance of managing RA to reduce its disabling effects (Cross et al., 2014).

RA predominantly affects women, with a female-to-male ratio of approximately 3:1. This gender disparity is believed to result from a combination of genetic factors, particularly X-linked traits, and hormonal influences (McInnes & Schett, 2011). The disease typically begins between the ages of 40 and 60, but it can also occur in younger and older individuals. Family history is another significant risk factor for RA, although it remains unclear whether this is due to genetic predisposition or shared environmental exposures within families (Padyukov et al., 2004). Environmental factors such as smoking, obesity, and certain infections have also been implicated in the development of RA (Scott et al., 2010). Identifying and understanding these risk factors is crucial for implementing preventive strategies and targeting early interventions.

Despite advancements in medical research and treatment options, the prognosis of RA often depends on how early the disease is diagnosed and managed. Early intervention with disease-modifying antirheumatic drugs (DMARDs) and biologics can slow disease progression, reduce inflammation, and prevent joint damage (Smolen et al., 2016). However, delayed diagnosis remains a significant challenge, particularly in regions where awareness of RA is limited. Studies have shown that low awareness of RA among the general population is associated with delayed diagnosis, poor treatment outcomes, and increased disease severity (Aletaha et al., 2010). This underscores the need for public health initiatives to raise awareness about RA, its symptoms, and its risk factors.

The consequences of RA extend beyond physical health. Patients with RA often experience social and economic challenges due to their reduced ability to work and participate in social activities. The disease can lead to significant financial burdens for individuals and healthcare systems due to the high costs of treatment, frequent medical visits, and loss of productivity (Scott et al., 2010). Furthermore, the psychological effects of living with a chronic disease, such as anxiety and depression, are common among RA patients, further diminishing their quality of life (Matcham et al., 2014).

The importance of studying RA and its risk factors cannot be overstated. Understanding the disease's etiology, progression, and impact can help guide the development of effective prevention and management strategies. Increased public awareness campaigns can play a pivotal role in promoting early diagnosis and encouraging individuals to seek medical attention when experiencing symptoms such as joint pain, stiffness, and swelling. Public health efforts should also focus on educating the population about modifiable risk factors, such as smoking cessation and maintaining a healthy weight, to reduce the risk of developing RA (Scott et al., 2010).

In conclusion, rheumatoid arthritis is a complex and multifaceted disease that demands attention from both the medical community and public health sectors. Its significant impact on quality of life, combined with its economic and social implications, highlights the need for increased awareness, early diagnosis, and effective management. By addressing these challenges, we can improve outcomes for individuals living with RA and reduce the overall burden of the disease on society. This study aims to assess public awareness

of rheumatoid arthritis and its risk factors among the population of Riyadh, Saudi Arabia, with the goal of identifying knowledge gaps and informing future educational and preventive initiatives.

METHODOLOGY

Research approach

Study design: The study is a cross-sectional study

Study area and population: The study was conducted in Riyadh. The target of the study is adult males and females residing in Riyadh, excluded mental ill, medical staff and students. A cluster stratification random sample was used to enroll participants from different public places. The Riyadh region will be cluster to 5 geographical areas (North, South, East, West, and Center). We assume that the five areas in Riyadh have a similar distribution of population. Therefore, the sample size was distributed proportionally for the five areas, where a sample of 108 was collected. Attendees or visitors to these public gathering areas, such as Malls, coffee shops, big supermarkets, and mosques, were asked to participate voluntarily in the survey. The first come, the first enrolled according to the inclusion criteria till the required sample is achieved from each are Sample size and technique: Based on a previous study in Jazan³⁶, Saudi Arabia, about Rheumatic arthritis knowledge among the public. They reported the level of public knowledge on the disease was 38.5%. Therefore, we used this prevalence to calculate the sample size with 95% CI = 95%, and a margin of error of 5%. The calculated sample size was 358. However, due to the clustering method of this study, additional calculation was done to add a design effect of 1.5. Thus, the total sample size will be 537

Data needs

Instrument: The study used a questionnaire designed specifically for this study, used to collect data for the investigation. It is divided into three sections: The first section discusses if the participant knows some information related to Rheumatoid Arthritis, which has 14 questions in a yes/no/I do not know format. In the second section, there are some signs and symptoms. The participant asked if it is related to Rheumatoid Arthritis or not. The final portion is about participant lifestyle, which will be asked in a 5-point Likert scale (1=never,2=rare,3=sometimes,4=mostly, and 5=always). The participant was finally divided into adequate and inadequate knowledge.

Definition and procedure: To describe adult understanding of Rheumatoid Arthritis

Data collection method: The study was paper based questionnaire in the Arabic language.

Data analysis and presentation: All data cleared, coded and entered using SPSS. The research is presented in tables and graphs as frequencies and percentages

Ethical consideration: This study will be reviewed by the Institutional Review Board committee at Al-Mareefa University. All participants' permissions will be collected when they open the questionnaire link and they will be informed about the relevant information that will be collected. The data and information will be solely used for research purposes only and will be kept confidential Result Description of the study sample: the total number of participants that were included in the study was 537 men and women. The response rate was 100%. The age of the study participants varied from 14 to 87. (Table 1) shows, 84.5% of the participants were below 40 years of age, and the mean age was 30.24 ± 12.62 (SD) years (median: 26.00). The majority of the study participants (73%) were university or postgraduate degree holders or students. About (47.9%) of the participants were employed, (26.1%) were students. The majority were Saudis (88.3%), and non-Saudis were approximately (12%). The body mass index of approximately half of participant were in normal weight (49.3%), (32.2%) were overweight, only (5%) were underweight, and (13.4%) were obese.

Table 1. Demographic characteristics of the participants

Variable	Category	No.	Percentage
Gender	Male	342	63.7
	Female	195	36.3

Education	Secondary and below	195	27.0
	Postgraduate	392	73.0
Age	<20 years	79	14.7
	21-30 years	287	53.4
	31-40 years	88	16.4
	≥41 years	83	15.5
Occupation	Employed	257	47.9
	Self-employed	59	11.0
	Un-employed	81	15.1
	Student	140	26.1
Nationality	Saudi	474	88.3
	Non-Saudi	63	11.7
BMI	Underweight	27	5.0
	Normal weigh	265	49.3
	Overweigh	173	32.2
	Obese	72	13.4

Table 2- shows among all participant only (3.7%) diagnosed with Rheumatoid Arthritis, (11%) of participant have other joint problem and (36.9%) have a close family patient diagnosed with Rheumatoid Arthritis.

Table 2. Source of knowledge of RA among the study participants.

Source of Knowledge	Categories	No.	%
Diagnosed with RA	Yes	20	3.7
	No	517	96.3
Diagnosed with any other joint problem	Yes	59	11
	No	478	89
Close family anyone diagnosed with RA	Yes	198	36.9
	No	339	63.1

Table 3A- Shows general knowledge of RA among the study participants is formed in several questions related to RA and their answers to measure their knowledge, such as if they think it is a serious condition, answered yes and almost half of them got the correct answer (46.7%). Be able to control RA by yourself, most of them got the correct answer and answered with no (52.1%). Treatable and available medicine, the majority got the correct answer and answered with yes (58.5). Patients need to routinely visit psychiatric sections (42.3%). Of the participants got the correct answer and answered with no. Is easily diagnosed that the answer was no and nearly a third of them got the right answer (30.7%). Can be treated by surgery (32.4%). Of the participants got the correct answer and answered with no. Can negatively affect the majority, got it correctly and answered with yes (60.9%). Just need a painkiller, the answer was no and (38%) of them got it correct. Usually, early signs, answers of no were (27.2%) and it was the correct answer. An autoimmune disease that answered it correctly with yes was about (32%). Exercise has a good effect, no was the correct answer (17.7%). Gender has the rule, answers formed in yes (32.4%), and it is the correct answer. Patients have the ability to walk long distances; the majority got the right answer and answered with no (60.7%).

Table 3A. Correct answers on general knowledge regarding Rheumatoid arthritis.

Variables	No.	%
A serious condition	251	46.7

Be able to control RA yourself	280	52.1
Patients need routinely visit psychiatric sections	227	42.3
Is easily diagnosed	165	30.7
Can be treated by surgery	174	32.4
Just need pain killer	204	38.0
Usually, early signs	146	27.2
Exercise has a good effect	95	17.7
Patients have the ability to	326	60.7
Treatable and available medication	314	58.5
Can negatively affect	327	60.9
An autoimmune disease	172	32.0
Gender has the rule	174	32.4

Table 3B- Shows General knowledge of RA among the study participants (67.6%) have inadequate knowledge, while (32.4%) have adequate knowledge from a total of 537 participants.

Table 3B. Correct answers on general knowledge regarding Rheumatoid arthritis

		Frequency	%	Cumulative Percent
Valid	Inadequate knowledge	363	67.6	67.6
	Adequate knowledge	174	32.4	100.0

Table 4A- Shows knowledge of signs and symptoms of RA among the study participants. Nearly 2/3 of the participants (67.6%) got the correct answer, pain in more than one joint is one of the symptoms of RA. Many of the participants (68.9%) got the correct answer that Aching is one of the symptoms of RA. Almost half of the participants (47.1%) agreed that morning stiffness is one of the symptoms. 2/3 of the participants (60.1%) also agreed that fatigue is one of the symptoms. Most of the participants (65.4%) agreed that weakness was one of the symptoms. More than half of the participants (54.9%) agreed that numbness is one of the symptoms. Tenderness is one of the symptoms of RA, and half of the participants (50.1%) said yes. (42.4%) of the participants got the correct answer that swelling is a symptom of RA. Nearly 1/3 of the participants (23.8%) got the correct answer for fever as a symptom of RA. (25.7%) Some of the participants thought a headache is symptom of RA. Almost half of the participants (48%) agreed that the wrist joint is one of the symptoms of RA. Sweating is a symptom of RA, (28.3%) said it is. (27.7%) Some of the participants thought weight loss was a symptom of RA. Insomnia is one of the symptoms of RA, (41%) of the participants agreed.

Table 4A. Correct answer on the knowledge of signs and symptoms of RA among the study participants.

Variables	No.	%
Pain in more than one joint	363	67.6
Aching	370	68.9
Morning	253	47.1
Fatigue	323	60.1
Weakness	351	65.4
Numbness	295	54.9
Tenderness	269	50.1
Swelling	228	42.4
Fever	128	23.8
Headache	138	25.7

Warn	258	48.0
Sweating	152	28.3
Weight loss	149	27.7
Insomnia	220	41.0

Table 4B- Shows that Knowledge of signs and symptoms of RA among the study participants (58.8%) have inadequate knowledge. While (41.2%) have adequate knowledge of sign and symptoms of RA

Table 4B. Knowledge of sign and symptoms of RA among the study participants

		Frequency	%	Cumulative Percent
Valid	Inadequate knowledge	316	58.8	58.8
	Adequate knowledge	221	41.2	100.0
	Total	537	100.0	

Table 5- shows among all participant only (4.3%) diagnosed with heart disease, (7.6%) of participant have Diabetes mellitus, (1.7%) have a renal problem, (2%) was diagnosed with liver disease and (6.3%) of participant have osteoporosis.

Table 5. Co-existing diseases

Diagnosed with	Categories	No.	%
Heart disease	No	514	95.7
	Yes	23	4.3
DM	No	496	92.4
	Yes	41	7.6
Renal	No	528	98.3
	Yes	9	1.7
Liver	No	525	97.8
	Yes	11	2.0
Osteoporosis	No	503	93.7

Table 6A- There is insignificant statistical relation between age and general knowledge (p Value = 0.092). Participant under 20 years which are (14.7% of all participants) (16.0%) of the participants who have inadequate knowledge of general knowledge represented by them and (12.1%) of the participants who have adequate knowledge of general knowledge represented by them. Participant between 21 and 30 years which are (53.4% of all participants) (53.7%) of the participants who have inadequate knowledge of general knowledge represented by them and (52.9%) of the participants who have adequate knowledge of general knowledge represented by them. Participant between 31 and 40 years which are (16.4% of all participants) (17.4%) of the participants who have inadequate knowledge of general knowledge represented by them and (14.4%) of the participants who have adequate knowledge of general knowledge represented by them. Participant above 40 years which are (15.5% of all participants) (12.9%) of the participants who have inadequate knowledge of general knowledge represented by them and (20.7%) of the participants who have adequate knowledge of general knowledge represented by them.

Table 6B- There is insignificant statistical relation between gender and knowledge of general knowledge (p Value = 0.134). According to the gender, participants who has inadequate knowledge of general knowledge (65.8%) of them were male and (34.2%) were female. On the other hand, participants with

adequate knowledge of general knowledge (59.2%) were male and (40.8%) were female. Table 6C- There is no significant statistical relation between education and knowledge of general knowledge (p-value = 0.833). According to the education, participants who had inadequate knowledge of general knowledge (26.7%) were secondary and below and (73.3%) were postgraduate. On the other hand, participants who had adequate knowledge of general knowledge (27.6%) were secondary and below and (72.4%) of them were postgraduate.

Table 6. Association of adequate general knowledge with demographic characteristics

		No.	%	No.	%	P value
Age	≤20 Years	58	16.0	21	12.1	0.0
	21-30 Years	195	53.7	92	52.9	92
	31-40 Years	63	17.4	25	14.4	
	≥41 Years	47	12.9	36	20.7	
Gender	Male	239	65.8	103	59.2	0.1
	Female	124	34.2	71	40.8	34

Education	secondary and below	97	26.7	48	27.6
	Postgraduate	266	73.3	126	72.4
Occupation	Employed	169	46.6	88	50.6
	Self-employed	39	10.7	20	11.5
	Un-employed	53	14.6	28	16.1
	Student	102	28.1	38	21.8
Nationality	Saudi	322	88.7	152	87.4
	Non-Saudi	41	11.3	22	12.6
BMI	Underweight	19	5.2	8	0.6
	normal weight	172	47.4	93	53.4
	Overweight	122	33.6	51	29.3
	Obese	50	13.8	22	12.6

Table 7A- There is significant statistical relation between age and knowledge of sign and symptoms (p Value = 0.025). Participant under 20 years which are (14.7% of all participants) (16.5%) of the participants who have inadequate knowledge of sign and symptoms represented by them and (12.2%) of the participants who have adequate knowledge of sign and symptoms represented by them. Participant between 21 and 30 years which are (53.4% of all participants) (55.7%) of the participants who have inadequate knowledge of sign and symptoms represented by them and (50.2%) of the participants who have adequate knowledge of sign and symptoms represented by them. Participant between 31 and 40 years which are (16.4% of all

participants) (16.1%) of the participants have inadequate knowledge of signs and symptoms represented by them and (16.7%) of the participants who have adequate knowledge of signs and symptoms represented by them. Participants above 40 years, which are (15.5% of all participants), (11.7%) of the participants who have inadequate knowledge of signs and symptoms represented by them and (20.8%) of the participants who have adequate knowledge of signs and symptoms represented by them.

Table 7B- There is no significant statistical relation between gender and knowledge of signs and symptoms (p-value = 0.158). According to gender, participants who had inadequate knowledge of signs and symptoms (66.1%) were male and (33.9%) were female. On the other hand, participants who had adequate knowledge of signs and symptoms of (60.2%) were male, and (39.8) were female. Table 7C- There is no significant statistical relation between education and knowledge of signs and symptoms (p-value = 0.212). According to the education, participants who had inadequate knowledge of signs and symptoms (25%) were secondary and below and (75%) were postgraduate. On the other hand, participants who had adequate knowledge of signs and symptoms of (29.9%) were secondary and below and (70.1%) of them were postgraduate.

Table 7. Association of adequate knowledge of signs and symptoms with some demographic characteristics

Variables	Categories	inadequate knowledge		adequate knowledge		P-Value
		No.	%	No.	%	
Age	≤20 Years	52	16.5	27	12.2	0.0 25
	21-30 Years	176	55.7	111	50.2	
	31-40 Years	51	16.1	37	16.7	
	≥41 Years	37	11.7	46	20.8	
Gender	Male	209	66.1	133	60.2	0.1 58
	Female	107	33.9	88	39.8	
Education	secondary and below	79	25.0	66	29.9	0.2 12
	Postgraduate	237	75.0	155	70.1	
Occupation	Employed	156	49.4	101	45.7	0.1 32
	Self-employed	31	9.8	28	12.7	
	Un-employed	40	12.7	41	18.6	
	Student	89	28.2	51	23.1	
Nationality	Saudi	278	88.0	196	88.7	0.8 00
	Non-Saudi	38	12.0	25	11.3	
BMI	Underweight	15	4.7	12	5.4	0.5 46
	normal weight	164	51.9	101	45.7	
	Overweight	98	31.0	75	33.9	
	Obese	39	12.3	33	14.9	

The general practices implemented by the participants were found to be highly inadequate (436/81.2%) versus (101/18.8%) of adequate physical exercise practice. This calculation was made by using a cut-off of 60% to show adequate and inadequate practice.

Variables	inadequate practice		%	No.	%	P-value
		No.				
Age	≤20 Years	64	14.7	15	14.9	0.024
	21-30 Years	221	50.7	66	65.3	
	31-40 Years	76	17.4	12	11.9	
	≥41 Years	75	17.2	8	7.9	
Gender	Male	263	60.3	79	78.2	0.001
	Female	173	39.7	22	21.8	
Education	secondary and below	116	26.6	29	28.7	0.667
	Postgraduate	320	73.4	72	71.3	
Occupation	Employed	202	46.3	55	54.5	0.362
	Self-employed	52	11.9	7	6.9	
	Un-employed	67	15.4	14	13.9	
	Student	115	26.4	25	24.8	
Nationality	Saudi	387	88.8	87	86.1	0.460
	Non-Saudi	49	11.2	14	13.9	
BMI	Underweight	20	4.6	7	6.9	0.088
	normal weight	212	48.6	53	52.5	
	Overweight	138	31.7	35	34.7	
	Obese	66	15.1	6	5.9	

DISCUSSION

The results of our research revealed that most participants had inadequate general knowledge and less than half of them had adequate general knowledge. This goes in line with a study conducted by Hazzazi in 2020, Saudi Arabia³⁶. We recommend ministry of health develop more programs to expand the general knowledge of the public awareness about RA. The outcome of our research showed that the majority of participants had inadequate knowledge of signs and symptoms and the minority had adequate knowledge of signs and symptoms. This goes in line with a study done by Naima Assaad in 2022, in Syria³⁷. On this basis, ministry of health should develop more programs to expand the knowledge of the public about RA sign and symptoms. There is insignificant relation between age and general knowledge, where the majority with adequate general knowledge are between the age of 21 to 30 years. Meanwhile, the minority are 20 years old or less. However, the majority with inadequate general knowledge are between the age 21 to 30. Whereas the minority were between 41 years or above. Our study revealed that most participants have inadequate general knowledge toward awareness to rheumatoid arthritis, this parallel a study was conducted by Hazzazi in 2020 in Saudi Arabia³⁶. In addition, we recommend other researchers to establish their research to a larger study population according to the demographic characteristics between age and general knowledge.

There is insignificant relation between gender and general knowledge, where the majority with adequate general knowledge are male, Meanwhile, the minority are females. However, the majority with inadequate general knowledge are male. Whereas the minority are females. Our study revealed that most participants have adequate general knowledge toward awareness to rheumatoid arthritis, which in contrary to a study done by Hazzazi in 2020 in Saudi Arabia³⁶. However, this might be influenced by different study

population. Therefore, we encourage other researchers to put their effort in equalizing gender participants.

There is insignificant relation between education and general knowledge, where the participant who had adequate knowledge 2/3 of them were postgraduate and 3/4 of participant who had inadequate knowledge were the same. This goes in line with a study done by Naima Assaad in 2022, in Syria³⁷. We recommend the ministry of health for further improvements in the education of general knowledge of RA in society and schools. There is insignificant relation between occupation and general knowledge. Most participants who have adequate knowledge were employed then followed by students. Meanwhile, participants who have inadequate knowledge were also employed then followed by student. This goes in contrary with a study by Hazzazi in 2020, Saudi Arabia³⁶, because the majority of their participant were un- employed. We recommend that other researcher to do more research with more participants.

There is insignificant relation between nationality and general knowledge. Saudi participants represent the majority of participant who have inadequate and adequate knowledge. While non-Saudi participants represent the minority of participants who have adequate and inadequate knowledge. This goes in line with a study done by Elhussein in 2018 Saudi Arabia³⁸. We recommend ministry of health to do more events about rheumatoid arthritis. There is significant relation between age and knowledge of sign and symptoms, where the majority with adequate knowledge of sign and symptoms were between 21 years old and 30 years old. Moreover, the minority with adequate knowledge were 20 years old or less. However, the majority with inadequate knowledge of sing and symptoms were between the age of 21 and 30, also the minority with inadequate knowledge were 41 years old and above. There is insignificant relation between gender and knowledge of sign and symptoms, where the majority with adequate knowledge of sing and symptoms were male, also the minority were females. However, the majority with inadequate knowledge of sign and symptoms were male and the minority were female. Our study revealed that most participants have in adequate knowledge about sign and symptoms of Rheumatoid arthritis, this parallel a study was conducted by Hazzai 2020 in Saudi Arabia³⁶. Meanwhile, we recommend other researchers establish their research to a larger study population according to the association of adequate knowledge of signs and symptoms with demographics characteristics.

There is insignificant relation between education and knowledge of sign and symptoms. Where the majority with adequate knowledge were postgraduate. Meanwhile, the minority were secondary and below. Also, it was the same for participant with inadequate knowledge. Our study shows that most participants had inadequate knowledge of sign and symptoms, this goes in contrary to a study done by Elhussein in 2018 Saudi Arabia³⁸. We encourage ministry of health to intensify public education to increase their awareness.

There is no significant relation between occupation and knowledge of signs and symptoms; the majority of adequate knowledge were employed. Meanwhile, most of the participants with inadequate knowledge were employed. This goes in contrary to a study by Hazzazi in 2020, Saudi Arabia³⁶, because the majority of their participant were unemployed. We recommend that another researcher to explore attitude and lifestyle.

CONCLUSION

In our study, the level of awareness was inadequate in general knowledge as well as the clinical manifestation of Rheumatoid Arthritis. Especially in participants who are more than 40 years which have the risk of getting the disease. Awareness about RA must be increased via the dissemination of information, organizing educational events or sharing the effects of RA in quality of life of patients. The Ministry of health develop more programs to expand the general knowledge of the public awareness about RA. The Ministry of Health should develop more programs to expand the knowledge of the public about RA signs and symptoms. Researchers must establish their research to a larger study population according to the demographic characteristics between age and general knowledge. Researchers should put their effort into equalizing the gender of participants. Ministry of Health for further improvements in the education of

general knowledge of RA in society and schools. Other researchers have explored attitude and lifestyle. The Ministry of Health should hold more events about rheumatoid arthritis. Intensify public education to increase their awareness.

REFERENCES

- Abdellah, et al. (2021). Study on lower limb enthesal abnormalities in RA patients using musculoskeletal ultrasound.
- Abdullah, et al. (2022). Study quantifying the incidence and severity of temporomandibular joint disorders (TMD) in RA patients in Saudi Arabia.
- Alamanos, Y., Voulgari, P. V., & Drosos, A. A. (2006). Incidence and prevalence of rheumatoid arthritis, based on the 1987 American College of Rheumatology criteria: A systematic review. *Seminars in Arthritis and Rheumatism*, 36(3), 182–188. <https://doi.org/10.1016/j.semarthrit.2006.08.006>
- Aletaha, D., Neogi, T., Silman, A. J., Funovits, J., Felson, D. T., Bingham, C. O., ... Hawker, G. (2010). 2010 Rheumatoid arthritis classification criteria: An American College of Rheumatology/European League Against Rheumatism collaborative initiative. *Arthritis & Rheumatism*, 62(9), 2569–2581. <https://doi.org/10.1002/art.27584>
- Bai, et al. (2020). Study analyzing quality of life among RA patients in Northeast China and its influencing factors like sleep and BMI.
- Cross, M., Smith, E., Hoy, D., Carmona, L., Wolfe, F., Vos, T., ... March, L. (2014). The global burden of rheumatoid arthritis: Estimates from the Global Burden of Disease 2010 study. *Annals of the Rheumatic Diseases*, 73(7), 1316–1322. <https://doi.org/10.1136/annrheumdis-2013-204627>
- Dakkak, et al. (2020). Study exploring MRI-detected intermetatarsal and submetatarsal bursitis in early RA patients and healthy controls.
- Dogu, et al. (2013). Study on the effect of exercise therapy on hand functions, dexterity, and quality of life in female RA patients.
- Elattar, et al. (2014). Study on hypothyroidism and thyroid autoantibodies' impact on RA disease activity.
- Elbeialy, et al. (2021). Study focusing on fibromyalgia syndrome and its classification compared to RA in Middle Eastern patients.
- Kreps, et al. (2018). Research on the association between weight loss and RA disease activity in the U.S.
- Matcham, F., Rayner, L., Steer, S., & Hotopf, M. (2014). The prevalence of depression in rheumatoid arthritis: A systematic review and meta-analysis. *Rheumatology*, 53(12), 2136–2148. <https://doi.org/10.1093/rheumatology/keu169>
- McInnes, I. B., & Schett, G. (2011). The pathogenesis of rheumatoid arthritis. *New England Journal of Medicine*, 365(23), 2205–2219. <https://doi.org/10.1056/NEJMra1004965>
- Neuberger, et al. (2007). Research on the effects of exercise, function, and aerobic fitness on RA patients.
- Nezamoleslami, et al. (2020). Study on dietary patterns and their relationship with the likelihood of RA in Iranian adults.
- Oelzner, et al. (2008). Study on the correlation between osteoporosis and RA, focusing on postmenopausal women and men.
- Padyukov, L., Seielstad, M., Ong, R. T., Ding, B., Rönnelid, J., Seddighzadeh, M., & Alfredsson,
- Genetics and Molecular Research 24 (4): gmr24124

L. (2004). A genome-wide association study suggests contrasting associations in ACPA-positive versus ACPA-negative rheumatoid arthritis. *Nature Genetics*, 44(4), 291–296. <https://doi.org/10.1038/ng.1076>

Panoulas, et al. (2007). Study on the prevalence of hypertension in RA patients and its association with cardiovascular disease and BMI.

Radwan, et al. (2020). Study on sleep quality in RA patients and its relation to disease activity, depression, and functional status in the Egyptian population.

Scott, D. L., Wolfe, F., & Huizinga, T. W. J. (2010). Rheumatoid arthritis. *The Lancet*, 376(9746), 1094–1108. [https://doi.org/10.1016/S0140-6736\(10\)60826-4](https://doi.org/10.1016/S0140-6736(10)60826-4)

Silman, A. J., & Hochberg, M. C. (2001). *Epidemiology of the rheumatic diseases*. Oxford University Press.

Sinha, et al. (2021). Study on lipid profile abnormalities and their relationship to cardiovascular risks and disease severity in RA patients.

Sköldstam, (2005). Study on the effect of weight reduction and its potential anti-inflammatory effects on RA patients.

Smolen, J. S., Aletaha, D., & McInnes, I. B. (2016). Rheumatoid arthritis. *The Lancet*, 388(10055), 2023–2038. [https://doi.org/10.1016/S0140-6736\(16\)30173-8](https://doi.org/10.1016/S0140-6736(16)30173-8)