

# MELATONIN GUMMY SELF-MEDICATION AMONG INDIAN YOUTH: A CROSS-SECTIONAL STUDY OF SLEEP QUALITY, LIFESTYLE FACTORS, AND THE IMPACT OF THE COVID-19 PANDEMIC

Sunny Babber<sup>1</sup>, Chandni Gupta<sup>2</sup>, Sandeepa Bailam<sup>3</sup>, Aakriti Khajuria<sup>4</sup>, Rakesh Khajuria<sup>5</sup>

<sup>1</sup>Associate Professor, Department of Community Medicine, Acharya Shri Chander College of Medical Sciences and Hospital, Jammu and Kashmir, Email- sunnybabber85@gmail.com, Orcid Id: 0009-0009-4263-8217

<sup>2</sup>Associate Professor, Department of Community Medicine, Acharya Shri Chander College of Medical Sciences and Hospital, Jammu and Kashmir, Email- Tkg2102@gmail.com, Orcid Id: 0000-0001-5335-7781

<sup>3</sup>Associate Professor, Department of Community Medicine, Acharya Shri Chander College of Medical Sciences and Hospital, Jammu and Kashmir, Email- sandeepahkumar@gmail.com, Orcid Id: 0009-0005-7379-2613

<sup>4</sup>PG Resident, Department of Psychiatry, Acharya Shri Chander College of Medical Sciences and Hospital, Jammu and Kashmir, Email- dr.aakritikhajuria@gmail.com, Orcid Id: 0009-0007-6613-6698

<sup>5</sup>Professor and HOD, Department of Community Medicine, Acharya Shri Chander College of Medical Sciences and Hospital, Jammu and Kashmir, Email- rakeshkhajuria61@gmail.com, Orcid Id: 0009-0008-4595-5819

## ABSTRACT

**Background:** Sleep disturbances have become increasingly common among young people, especially after the COVID-19 pandemic, and the use of over-the-counter sleep aids has increased, especially melatonin. The palatable nature of melatonin gummies and their convenient accessibility have increased their popularity among the young, where the use of these substances is not regulated by a doctor.

**Aim:** The aim of the current research is to determine the rates and trends of self-medication using melatonin gummies among Indian youth and to determine the relationship between these factors and their sleep quality, lifestyle habits, and influencing factors of COVID-19.

**Methods:** An online Google Form using social media platforms administered a cross-sectional, questionnaire-based study to 1450 participants aged 18 years to 30 years. Sleep quality was assessed using a self-developed structured sleep assessment questionnaire including subjective sleep quality, sleep duration, sleep latency, sleep disturbances, and daytime dysfunction. Melatonin use, self-medication habits, lifestyle, and adverse effects were gathered. Statistical procedures were done by the use of SPSS where descriptive statistics and chi-square tests were applied with  $p < 0.05$  taken to be significant.

**Results:** The prevalence of melatonin gummy use was 54.4%. Almost half (49.6%) of the respondents had mild sleep disturbance with 34.2% having poor sleep quality. There was a significant association between poor sleep quality and melatonin gummy use ( $p < 0.001$ ). Most users started taking melatonin during (42.0%), or following (38.0%), the COVID-19 pandemic. It was found that moderate to high levels of self-medication were used by 73% of the participants, with 48.5% saying that they experienced adverse effects. The most prevalent reasons of use were sleep disturbance (65.9%), stress (57.7%), and irregular schedules (50.9%) reasons. Stress and screen time were found to be significantly associated with poor sleep quality.

**Conclusion:** Melatonin gummy use was highly prevalent among Indian youth and was significantly associated with poor sleep quality, lifestyle factors, and post-pandemic behavioral changes. The high prevalence of self-medication and adverse effects highlights the need for greater awareness, regulation, and emphasis on non-pharmacological management of sleep disturbances.

**KEYWORDS:** COVID-19, Lifestyle Factors, Melatonin Gummies, Sleep Quality, Self-medication, Youth.

## INTRODUCTION

Sleep is a biological process that is essential in the physical and psychological well-being. In recent years, the problem of sleep disorders has also been a significant concern in terms of population health particularly among young adults due to their academic demands, excessive exposure to screens and unbalanced ways of living. Coupled with the COVID-19 pandemic, these issues were further aggravated as people have been more uprooted concerning their daily routine, more psychologically affected, and more signs of insomnia, or what is often called a coronasomnia [1]. Being an over-the-counter hormone and being considered a safe and non-toxic compound, Melatonin has become a widely used sleep aid since its discovery as an endogenous hormone that controls the circadian rhythm. Melatonin is also being actively consumed in India in the form of gummies, a product being marketed as non-toxic and as a palatable sleep aid. It is this formulation that particularly appeals to the young generation, since it is like confectionery and being actively promoted on the internet [2]. Melatonin is a neurohormone that is endogenous in nature and is primarily synthesized in the pineal

gland and plays significant roles in the control of circadian rhythm as well as in the stimulation of sleep. Its exogenous form to be used typically in the synthetic forms is normally used as a chrono biotic agent to prevent jet lag, shift work disorder, or certain primary sleep onset insomnias [3]. However, its increased over-the-counter availability has also led to the increase of its unofficial usage particularly among the younger populations who are likely to seek easy solutions to sleeping disturbances [4]. The growing popularity of melatonin is becoming a cause of concern on the likelihood of misuse and adverse effects of melatonin, particularly due to the lack of stringent regulatory control of dietary supplements [5]. Another notable development which has defined the post-pandemic period is the shift in the attitude of health-seeking behavior whereby majority of the populace are more inclined towards self-medication, especially among the youth segments of the population [6]. The easy access, the heavy advertisement, and the influence of social media have prompted the non-prescription use of dietary supplements, and melatonin is one of them. Despite the fact that people tend to consider these gummies as harmless, the inappropriate use of melatonin, in particular its use in the absence of medical protection and in inappropriate sleep aid doses, may lead to certain side effects, daytime sleepiness, hormonal disorder, and post-use sleep improvement and the impact of the COVID-19 pandemic on the consumption of such gummies [7,8]. The given research will assist in identifying the popularity and tendencies of melatonin gummies self-medication in young Indians, its impact on their sleep and explore how COVID-19 pandemic influences this new practice.

**Aim:** To determine the frequency and trends of self-prescription of melatonin gummies in Indian youth and to examine its association with the quality of sleep, lifestyle variables, and the COVID-19 pandemic influence.

### **Objectives:**

1. To identify the prevalence and frequency of using melatonin gummies in young people aged 18-30 years.
2. To evaluate sleep quality among respondents using a structured sleep assessment questionnaire.
3. To assess the relation between sleep quality and the use of melatonin gummies.
4. To determine the time of starting melatonin, use in connection with COVID-19 pandemic.
5. To determine the typical causes of melatonin gummies use.
6. To assess self-medication behaviors and how it correlates with negative outcome.
7. To examine how the lifestyle factors (stress levels and screen time before bedtime) relate to the quality of sleep.

### **MATERIAL AND METHODS**

This was a cross-sectional and observational study that involved young people aged 18-30 years in India to determine the use of melatonin gummies, the quality of sleep, and other related variables. The research was conducted by the department of community medicine in a tertiary care hospital a 3-month period from November 2025 to January 2026 after getting approval from the Institutional Ethical Committee with reference no. ASCOMS/IEC/2025/Meeting-II/FM/29 dated 18/09/25 on the basis of online survey with the use of the Google Forms system, and the questionnaire was spread over a three-month period in a social network (WhatsApp, Instagram, email, etc.). Participants were recruited by means of convenience sampling. Participants with diagnosed sleep disorders undergoing treatment, those with severe psychiatric or neurological disorders, and those who did not complete the questionnaire were excluded from the study. The survey on Google Forms received a total of 1511 responses out of which 1450 were complete and met the inclusion criteria; the rest was incomplete and inconsistent, and was not included in the final analysis.

The self-administered, structured questionnaire was created by using literature previously published on sleep quality assessment, and some of the most commonly assessed domains were included: subjective sleep quality, sleep duration, sleep latency, sleep disturbances and daytime dysfunction. The questionnaire was pretested with 30 subjects for clarity and feasibility, and The questionnaire demonstrated satisfactory internal consistency (Cronbach's  $\alpha = 0.84$ ). Changes were made based on feedback obtained during pilot testing. The results of the pilot study were not used in the final analysis. The survey form was comprised of various sections. Electronic informed consent was obtained in the first part after which the socio-demographic information comprised of age, gender, education, and occupation. Sleep quality was assessed using a structured sleep assessment questionnaire that included subjective sleep quality, sleep duration, sleep latency, sleep disturbances, and daytime dysfunction. Each item was scored on a 4-point scale, and a composite sleep score ranging from 0–15 was calculated. Follow-up parts evaluated the use of melatonin gummies, prevalence, frequency, length, and time of initiation concerning the COVID-19 pandemic. A risk score was created by evaluating self-medication behavior using consultation practices, sources of information and awareness of dosage. Other sections addressed use reasons, self-reported adverse effects and lifestyle e.g., stress levels and screen time before sleep that were compared in relation to quality of sleep.

### **Statistical Analysis**

Statistical analysis was performed by IBM SPSS 21. Categorical variables like age, gender, education, occupation, sleep quality, melatonin use, and self-medication practices were summarized using descriptive statistics which were summarized in frequencies and percentages and presented in tables and figures. Sleep quality was classified into good sleep, mild sleep disturbance, and poor sleep categories according to composite sleep scores obtained from the structured sleep assessment questionnaire. The Chi-square test was used in inferential analysis to identify the relationship between categorical variables, such as sleep quality and melatonin use, risk of self-medication and adverse effects, and lifestyle factors, i.e. the level of stress and screen time in relation to sleep quality. A p-value of below 0.05 was taken as statistically significant and results were interpreted accordingly without assuming any causality because the design of the study was cross-sectional.


## RESULTS

A total of 1,450 participants were included in the study with overall mean age of  $24 \pm 2.63$ .

**Table 1. Socio-demographic Profile and Sleep Quality of Participants (N = 1450)**


| Variable                                              | Category               | N   | %    |
|-------------------------------------------------------|------------------------|-----|------|
| Age (years)                                           | 18–21                  | 492 | 33.9 |
|                                                       | 22–25                  | 479 | 33.0 |
|                                                       | 26–30                  | 479 | 33.0 |
| Gender                                                | Male                   | 643 | 44.3 |
|                                                       | Female                 | 803 | 55.4 |
|                                                       | Prefer not to say      | 04  | 0.3  |
| Education                                             | Undergraduate          | 369 | 25.4 |
|                                                       | Graduate               | 361 | 24.9 |
|                                                       | Postgraduate           | 356 | 24.6 |
|                                                       | Other                  | 364 | 25.1 |
| Occupation                                            | Student                | 366 | 25.2 |
|                                                       | Employed               | 354 | 24.4 |
|                                                       | Unemployed             | 370 | 25.5 |
|                                                       | Other                  | 360 | 24.8 |
| Sleep Quality (Self-developed sleep assessment scale) | Good sleep (0–4)       | 235 | 16.2 |
|                                                       | Mild disturbance (5–9) | 719 | 49.6 |
|                                                       | Poor sleep (10–15)     | 496 | 34.2 |
| Melatonin Gummy Use                                   | Yes                    | 789 | 54.4 |
|                                                       | No                     | 661 | 45.6 |

Table 1. The age distribution was evenly divided among three groups: 18–21 years ( $n = 492$ , 33.9%), 22–25 years ( $n = 479$ , 33.0%), and 26–30 years ( $n = 479$ , 33.0%). The sample comprised males ( $n = 643$ , 44.3%) and females ( $n = 803$ , 55.4%), while a small proportion preferred not to disclose their gender ( $n = 4$ , 0.3%). Educational attainment was evenly distributed across undergraduate ( $n = 369$ , 25.4%), graduate ( $n = 361$ , 24.9%), postgraduate ( $n = 356$ , 24.6%), and other categories ( $n = 364$ , 25.1%). Similarly, occupational status included students ( $n = 366$ , 25.2%), employed individuals ( $n = 354$ , 24.4%), unemployed individuals ( $n = 370$ , 25.5%), and other categories ( $n = 360$ , 24.8%). Regarding sleep quality measured using the self-developed sleep assessment scale, the respondents rated their sleep as good ( $n = 235$ , 16.2%), mildly disturbed ( $n = 719$ , 49.6%), and poor ( $n = 496$ , 34.2%). Moreover, more than half of the participants ( $n = 789$ , 54.4%), and the rest ( $n = 661$ , 45.6%) did not use melatonin gummies.




**Figure 1. Frequency of Melatonin Gummy Use among Participants (n = 789)**

The frequency of the use of melatonin gummies among the participants is shown in figure 1. Among users, the most common pattern was daily use ( $n = 368$ , 46.6%), followed by use 3–5 times per week ( $n = 201$ , 25.5%) and 1–2 times per week ( $n = 164$ , 20.8%). A lower percentage of the participants ( $n = 56$ , 7.1%) reported occasional use.



**Figure 2. Timing of Initiation of Melatonin Gummy Use among Users (n = 789)**

Figure 2 shows the initiation date of the use of melatonin gummies among the respondents who reported the use (n = 789). Most of the users started using melatonin during the COVID-19 period (n = 331, 42.0%), after COVID-19 (n = 300, 38.0%), and only a smaller percentage started using it before the COVID-19 period (n = 158, 20.0%).




**Figure 3. Reasons for melatonin gummy use (n = 789)**

Figure 3 shows the reasons for melatonin gummy use among those respondents who reported the use (n = 789). The most commonly reported reason was sleep disturbance (n = 520, 65.9%), followed by stress or anxiety (n = 455, 57.7%), and irregular schedules (n = 402, 50.9%). A significant percentage of participants (n = 298, 37.8%), reported social media influence, which was the least reported reason (n = 165, 20.9%).

**Table 2. Association between Sleep Quality and Melatonin Gummy Use (N = 1450)**


| Sleep Quality    | Non-Users (n = 661) | Users (n = 789) | Total |
|------------------|---------------------|-----------------|-------|
| Good sleep       | 210 (31.8%)         | 25 (3.2%)       | 235   |
| Mild disturbance | 365 (55.2%)         | 354 (44.9%)     | 719   |
| Poor sleep       | 86 (13.0%)          | 410 (52.0%)     | 496   |
| Chi-square (df)  | 412.6 (2)           |                 |       |
| p-value          | <0.001*             |                 |       |

Table 2 shows the association between sleep quality and use of melatonin gummy among the participants (N = 1,450). Among non-users (n = 661), the majority reported mild sleep disturbance (n = 365, 55.2%), followed by good sleep (n = 210, 31.8%) and poor sleep (n = 86, 13.0%). Conversely, among the users (n = 789), more than half reported having poor sleep quality (n = 410, 52.0%), then the mild disturbance (n = 354, 44.9%), with only a small percentage reporting the good sleep (n = 25, 3.2%). A chi-square test of independence revealed a statistically significant relationship between the quality of sleep and the use of melatonin gummies, ( $\chi^2$  (2) = 412.6,  $p < 0.001^*$ ), which showed that poor sleep quality had a statistically significant relationship with increased melatonin gummy use.



**Figure 4. Distribution of Self-Medication Risk among Melatonin Users (n = 789)**

Figure 4 shows the trend of self-medication risk among the users of melatonin gummies (n = 789). The highest percentage of respondents was in the moderate risk group (n = 349, 44.3%), then the high-risk group (n = 227, 28.7%), and lastly the low-risk group (n = 213, 27.0%). It shows that a significant percentage of the users were moderate to high-risk self-medication.



**Figure 5. Prevalence of Adverse Effects among Melatonin Users (n = 789)**

Figure 5 presents the prevalence of adverse effects among melatonin gummy users (n = 789). Nearly half of the participants reported experiencing adverse effects (n = 383, 48.5%), while a slightly higher proportion reported no adverse effects (n = 406, 51.5%).

**Table 3. Association between Self-Medication Risk and Adverse Effects Among Melatonin Users (n = 789)**

| Self-Medication Risk | No Adverse Effects (n = 406) | Adverse Effects (n = 383) | Total |
|----------------------|------------------------------|---------------------------|-------|
| Low risk             | 150 (36.9%)                  | 63 (16.4%)                | 213   |
| Moderate risk        | 180 (44.3%)                  | 169 (44.1%)               | 349   |
| High risk            | 76 (18.7%)                   | 151 (39.4%)               | 227   |
| Chi-square (df)      | 137.5 (2)                    |                           |       |
| p-value              | <0.001*                      |                           |       |

Table 3 includes the association between the risk of self-medication and adverse effects of melatonin gummies users (n = 789). Among participants reporting no adverse effects (n = 406), the majority were in the moderate risk category (n = 180, 44.3%), followed by low risk (n = 150, 36.9%) and high risk (n = 76, 18.7%). Conversely, in the cases of those reporting adverse effects (n = 383, 48.5%) the highest percentage of the population was in the moderate risk group (n = 169, 44.1%), with the next highest proportion in the high-risk group (n = 151, 39.4%), and the lowest percentage was in the low-risk category (n = 63, 16.4%). A chi-square test of independence showed that self-medication risk and the prevalence of adverse effects were statistically significantly related, ( $\chi^2$  (2) = 137.5,  $p < 0.001^*$ ), and so the higher the self-medication risk, the higher the probability of experiencing adverse effects.

**Table 4. Comparison of Lifestyle Factors between Melatonin Users and Non-Users (N = 1450)**

| Table 4. Comparison of Lifestyle Factors between Melatonin Users and Non-Users (N = 1456) |                     |                 |                 |         |
|-------------------------------------------------------------------------------------------|---------------------|-----------------|-----------------|---------|
| General Stress Level                                                                      | Melatonin           |                 | Chi-square (df) | p-value |
|                                                                                           | Non-Users (n = 661) | Users (n = 789) |                 |         |
| Low                                                                                       | 210 (31.8%)         | 113 (14.3%)     | 96.4 (2)        | <0.001* |
| Moderate                                                                                  | 250 (37.8%)         | 261 (33.1%)     |                 |         |
| High                                                                                      | 201 (30.4%)         | 415 (52.6%)     |                 |         |
| Screen time before bed                                                                    |                     |                 |                 |         |
| <1 hour                                                                                   | 210 (31.8%)         | 117 (14.8%)     | 121.7 (2)       | <0.001* |
| 1–2 hours                                                                                 | 250 (37.8%)         | 230 (29.1%)     |                 |         |

|          |             |             |  |  |
|----------|-------------|-------------|--|--|
| >2 hours | 201 (30.4%) | 442 (56.0%) |  |  |
|----------|-------------|-------------|--|--|

Table 4 shows the comparison of the lifestyle factors between melatonin users and non-users (N = 1,450). With respect to general stress levels, a greater percentage of melatonin users (n = 415, 52.6%) than non-users (n = 201, 30.4%) reported high levels of stress, and more non-users (n = 210, 31.8%) than users (n = 113, 14.3%) reported low levels of stress. The relative moderate levels of stress were not much different between the two groups (non-users: n = 250, 37.8%, users: n = 261, 33.1%). The chi-square test showed that the level of stress and the use of melatonin were statistically related, ( $\chi^2$  (2) = 96.4,  $p < 0.001^*$ ).

Likewise, there were very strong variations in pre-bed screen time. A greater proportion of melatonin users reported more than 2 hours of screen time (n = 442, 56.0%) compared to non-users (n = 201, 30.4%), whereas shorter screen time (<1 hour) was more common among non-users (n = 210, 31.8%) than users (n = 117, 14.8%). The prevalence of participants with 1-2 hours of screen time also was higher among non-users (n = 250, 37.8%) as compared to users (n = 230, 29.1%). This was statistically significant, ( $\chi^2$  (2) = 121.7,  $p < 0.001^*$ ), which means that increased stress levels and increased screen time before bed were significantly related with the use of melatonin gummies.

## DISCUSSION

The present study has discussed the frequency and trend of melatonin gummy consumption among Indian youth and how it correlates with the sleep quality, self-medication, and lifestyle among participants in the present study, the socio-demographic profile was well balanced in terms of age, gender, education, and occupation. The balanced sociodemographic distribution improved the representativeness and generalizability of the findings while minimizing demographic bias of the findings. The same trend is reflected in other research works involving online surveys among the young adults [9]. The present research study discovered a high prevalence of sleep disturbances where approximately half of the participants reported mild disturbance and more than one-third of the participants reported poor sleep quality. It is in line with the literature that documented more insomnia and disturbed sleep patterns among young adults during and after the pandemic [10]. An important result of the current research is that over fifty percent of the interviewees said that they used melatonin gummies, indicating a high prevalence of use. The frequency distribution with a significant percentage of daily users indicates that people do not use it occasionally but frequently. This is in line with results by Al-Naggar RA, Chen R. who reported that over-the-counter supplements were used more by the youth during the post-pandemic period [11]. The issue of self-medication is also worrying as noted in the present study, with most of the respondents being in moderate and high-risk groups. Almost fifty percent of the participants had adverse effects. Even though melatonin is usually regarded as relatively safe, these results indicate that its use ought not to be uncontrolled as it could lead to an amplification of side effects. This finding is supported by previous research, which showed variability in the content of melatonin supplement and dangers of its misuse [12]. Moreover, the current research illustrates that there is a strong association between the quality of sleep and the use of melatonin, where participants with poor sleep quality are more inclined to use melatonin. This result is not surprising since patients with sleeping problems tend to resort to drugs or supplementary treatment [13]. The time when melatonin was initiated as revealed in the present study also highlights the effect of the pandemic where most participants indicated that they started using melatonin during or after the COVID-19 time. This finding supports the concept of coronasomnia, in which stress, anxiety, and the interruption of normal activities contributed to sleep disturbances and eventual consumption of sleep aids [14]. Regarding the causes of use, sleep disturbance, stress/anxiety and irregular schedules were the most widespread ones, with the impact of social media playing a significant role. This shows that the clinical requirement and the behavioral factors also contribute to the usage of melatonin. The increasing influence of social media on health-related behaviors, especially in the youth, has been a highly recognized issue in the recent studies [15]. Another finding of the current research was a strong relationship between increased risk of self-medication and the presence of adverse effects. The participants of the high-risk group also reported considerably more side effects than those of the low-risk group, which indicates that wrongful or unmonitored use leads to adverse outcomes [16]. Also, other lifestyle changes, including stress levels and time spent in front of the screen before sleeping, had a great effect on the quality of sleep. Poor sleep outcomes were associated with the increase in the levels of stress and long screen time. These results align with what has been found in past studies in which an association between screen time and sleep disturbance was established and literature reports stressful experiences as a major determinant of sleep quality [17]. Altogether, the results of the current research point at the presence of a complicated interconnection between sleeping problems, lifestyle habits and melatonin consumption in young people. The rise in the use of melatonin gummies, especially in the wake of the COVID-19 pandemic, highlights the fact that self-medication is becoming increasingly more of a trend that is driven by both social and psychological issues. Although the use of melatonin can have temporary positive effects, its unsupervised administration raises important concerns in terms of safety and necessity of raising awareness and regulation levels. [18,19]. A high prevalence of taking melatonin gummies without a professional medical recommendation requires a critical analysis of the labeling accuracy of the product and the possible consequences of taking it over the long term in this group of people [20]. All in all, this evidence supports the idea of the dire necessity of balanced awareness, responsible melatonin usage, and greater emphasis on underlying lifestyle factors that lead to sleep disorders in youth.

## CONCLUSION

The current study showed a high prevalence of melatonin gummy use among youth in India and a significant number of them using these products for self-medication without any medical advice. Increased melatonin use was significantly associated with poor sleep quality, indicating that melatonin supplements are more likely to be used by persons with

existing sleep issues. The COVID-19 pandemic seemed to have had a significant effect as a catalyst for starting and boosting melatonin intake, which is consistent with the overall effect on sleep and mental health. Important safety concerns were identified, given the size of the self-medication groups of moderate and high risk, as well as the high proportion of participants stating that they had experienced adverse effects. Additionally, lifestyle factors like high levels of stress and extended hours of screen use prior to bedtime were significantly linked with melatonin use and poor sleep quality. The results highlight the importance of increasing public awareness of the potential risks of melatonin supplements and their appropriate use, as well as effective regulation of OTC supplements and the promotion of non-pharmaceutical interventions (NPIs) for sleep health that include better sleep hygiene, stress management and less screen time exposure for youth.

### Limitations

The findings of the present study are subject to several limitations: First, the cross-sectional nature of the study does not permit the establishment of cause-and-effect relations among the use of melatonin gummies, sleep quality, various lifestyle factors, and self-medication behaviors. Second, the data was self-administered online questionnaire and may be vulnerable to recall bias, reporting bias and social desirability bias. Thirdly, the use of convenience sampling using social media could have skewed towards selection bias and restricted the findings to Indian youth population and not the wider population, especially those with limited internet access and digital literacy.

Although the questionnaire demonstrated acceptable internal consistency and was pilot tested prior to the study, it was not a formally validated psychometric instrument, which may limit comparability with standardized sleep assessment tools. In addition, information on precise melatonin concentration, differences in formulation, long-term usage and the potential for simultaneous administration of other sleep drugs was not evaluated. Lastly, psychological factors (anxiety, depression and academic stress) were not assessed separately, and this can also affect sleep quality and melatonin consumption.

### Declarations

#### Ethics approval and consent to participate

This study was approved by Acharya Shri Chander College of Medical Sciences and Hospital, Jammu, India, Institutional Independent Ethics Committee, ASCOMS/IEC/2025/Meeting-II/FM/29 dated 18/09/2025. All participants gave informed consent electronically before participating in the study.

#### Consent for publication

Not applicable.

#### Availability of data and materials

The datasets used and/or analyzed in the current study are available, upon reasonable request, from the corresponding author.

#### Competing interests

The authors declare that there are no competing interests.

#### Funding

The authors did not receive financial support for the research, for the authorship or for the publication of this article.

#### Acknowledgements

The authors would like to thank all the participants who willingly participated in the study and provided their time and responses.

#### List of Abbreviations

| Abbreviation | Definition                                  |
|--------------|---------------------------------------------|
| COVID-19     | Coronavirus Disease 2019                    |
| IEC          | Institutional Ethics Committee              |
| IBM          | International Business Machines             |
| NPI          | Non-Pharmacological Intervention            |
| OTC          | Over-the-Counter                            |
| SPSS         | Statistical Package for the Social Sciences |
| WHO          | World Health Organization                   |

#### REFERENCES

- Selak Š, Šorgo A, Crnković N, Gabrovec B, Cesar K, Žmavc M. Three measures of internet use, social media use and video game playing as predictors of insomnia during the pandemic among students. *Scientific Reports*. 2024;14(1). doi:10.1038/s41598-024-53351-2
- Cohen PA, Avula B, Wang Y, Katragunta K, Khan I. Quantity of Melatonin and CBD in Melatonin Gummies Sold in the US. *JAMA*. 2023;329(16):1401. doi:10.1001/jama.2023.2296
- Arendt J, Skene DJ. Melatonin as a chronobiotic. *Sleep Medicine Reviews*. 2004;9(1):25. doi:10.1016/j.smr.2004.05.002

4. Pagel JF, Pandi-Perumal SR, Monti JM. Treating insomnia with medications. *Sleep Science and Practice*. 2018;2(1). doi:10.1186/s41606-018-0025-z
5. Lelak K, Vohra V, Neuman MI, Toce MS, Sethuraman U. Pediatric Melatonin Ingestions — United States, 2012–2021. *MMWR Morbidity and Mortality Weekly Report*. 2022;71(22):725. doi:10.15585/mmwr.mm7122a1
6. Khajuria A, Gupta U, Gupta RK, Babber S: Behavioural and psychological impact of COVID-19 and lockdown on children and their parents. *Glob J Res Anal*. 2023, 12:45-9. 10.36106/gjra
7. Costello RB, Lentino CV, Boyd C, et al. The effectiveness of melatonin for promoting healthy sleep: a rapid evidence assessment of the literature. *Nutrition Journal*. 2014;13(1). doi:10.1186/1475-2891-13-106
8. Dutta R, Raja D, Anuradha R, D’Cruze L, Jain T, Sivaprakasam P. Self-medication practices versus health of the community. *International Journal of Community Medicine and Public Health*. 2017;4(8):2757. doi:10.18203/2394-6040.ijcmph20173169
9. Kalimeri K, Beiró MG, Bonanomi A, Rosina A, Cattuto C. Traditional versus Facebook-based surveys: Evaluation of biases in self-reported demographic and psychometric information. *Demographic Research*. 2020;42:133. doi:10.4054/demres.2020.42.5
10. Riva E, Terraneo M, Lucchini M, Gerosa T. The prevalence of insomnia in different COVID-19 policy phases: Longitudinal evidence from ITA.LI – Italian Lives. *BMC Public Health*. 2022;22(1). doi:10.1186/s12889-022-14048-1
11. Al-Naggar RA, Chen R. Prevalence of vitamin-mineral supplements use and associated factors among young Malaysians. *Asian Pac J Cancer Prev*. 2011;12(4):1023-9.
12. Tuft C, Matar E, Schrire ZM, Grunstein RR, Yee BJ, Hoyos CM. Current Insights into the Risks of Using Melatonin as a Treatment for Sleep Disorders in Older Adults. *Clinical Interventions in Aging*. Published online January 1, 2023:49. doi:10.2147/cia.s361519
13. Koopman-Verhoeff ME, Dries MA van den, Seters JJ van, Luijk M, Tiemeier H, Luik AI. Association of Sleep Problems and Melatonin Use in School-aged Children. *JAMA Pediatrics*. 2019;173(9):883. doi:10.1001/jamapediatrics.2019.2084
14. Chen S, Cheng C. Unveiling Coronasomnia: Pandemic Stress and Sleep Problems During the COVID-19 Outbreak. *Nature and Science of Sleep*. Published online May 1, 2024:543. doi:10.2147/nss.s459945
15. Wesselhöft R, Rasmussen L, Jensen PB, et al. Use of hypnotic drugs among children, adolescents, and young adults in Scandinavia. *Acta Psychiatrica Scandinavica*. 2021;144(2):100. doi:10.1111/acps.13329
16. Lee CH, Chang F, Hsu S, Chi HY, Huang LJ, Yeh M. Inappropriate self-medication among adolescents and its association with lower medication literacy and substance use. *PLoS ONE*. 2017;12(12). doi:10.1371/journal.pone.0189199
17. Zhong C, Masters M, Donzella SM, Diver WR, Patel AV. Electronic Screen Use and Sleep Duration and Timing in Adults. *JAMA Network Open*. 2025;8(3). doi:10.1001/jamanetworkopen.2025.2493.
18. Erland LAE, Saxena PK. Melatonin natural health products and supplements: Presence of serotonin and significant variability of melatonin content. *J Clin Sleep Med*. 2017;13(2):275-81. doi:10.5664/jcsm.6462.
19. Saqib W, Gazerani P. Social Media Use and Consumption of Prescription-Free Medications for Anxiety, Sleep, and Pain among Norwegian University Students. *European Journal of Investigation in Health Psychology and Education*. 2024;14(8):2205. doi:10.3390/ejihpe14080147
20. Hartstein LE, Garrison MM, Lewin D, Boergers J, LeBourgeois MK. Characteristics of Melatonin Use Among US Children and Adolescents. *JAMA Pediatrics*. 2023; 178(1):91. doi:10.1001/jamapediatrics.2023.4749

## Questionnaire with Scoring System

### Title: Self-medication with Melatonin Gummies among Indian Youth

| Section                                           | Question/Domain             | Response Options                                        | Scoring                        |
|---------------------------------------------------|-----------------------------|---------------------------------------------------------|--------------------------------|
| Section 1: Consent                                | Participation consent       | Yes / No                                                | Non-scoring                    |
| Section 2: Socio-demographic Details              | Age                         | 18–21 / 22–25 / 26–30                                   | Non-scoring                    |
|                                                   | Gender                      | Male / Female / Prefer not to say                       | Non-scoring                    |
|                                                   | Education                   | Undergraduate / Graduate / Postgraduate / Other         | Non-scoring                    |
|                                                   | Occupation                  | Student / Employed / Unemployed / Other                 | Non-scoring                    |
| Section 3: Sleep Quality Assessment Questionnaire | Subjective sleep quality    | Very good / Fairly good / Fairly bad / Very bad         | 0–3                            |
|                                                   | Sleep duration              | ≥7 h / 6–7 h / 5–6 h / <5 h                             | 0–3                            |
|                                                   | Sleep latency               | <15 min / 15–30 min / 30–60 min / >60 min               | 0–3                            |
|                                                   | Sleep disturbance           | None / Once weekly / 2–3 times weekly / ≥4 times weekly | 0–3                            |
|                                                   | Daytime dysfunction         | None / Mild / Moderate / Severe                         | 0–3                            |
|                                                   | Total Sleep Score           | Range: 0–15                                             | 0–4 Good, 5–9 Mild, 10–15 Poor |
|                                                   | Ever used melatonin gummies | Yes / No                                                | 1 / 0                          |

|                                  |                                                         |                                                                                        |                                     |
|----------------------------------|---------------------------------------------------------|----------------------------------------------------------------------------------------|-------------------------------------|
| Section 4: Melatonin Gummy Use   | Initiation of use                                       | Before COVID-19 / During COVID-19 / After COVID-19                                     | 0–2                                 |
|                                  | Frequency of use                                        | Occasional / 1–2 times weekly / 3–5 times weekly / Daily                               | 0–3                                 |
|                                  | Duration of use                                         | <1 month / 1–3 months / 3–6 months / >6 months                                         | 0–3                                 |
| Section 5: Self-Medication Score | Consulted a doctor before using melatonin               | Yes / No                                                                               | 0 / 1                               |
|                                  | Source of information                                   | Doctor / Social media / Friends / Pharmacy / Internet                                  | 0–4                                 |
|                                  | Awareness of dosage                                     | Yes / No                                                                               | 0 / 1                               |
|                                  | Total Self-Medication Score                             | Range: 0–6                                                                             | 0–2 Low, 3–4 Moderate, 5–6 High     |
| Section 6: Reasons for Use       | Main reasons                                            | Sleep disturbance / Stress / Irregular schedule / Social media / Peer pressure / Other | Non-scoring                         |
| Section 7: Adverse Effects Score | Any side effects                                        | Yes / No                                                                               | 1 / 0                               |
|                                  | Number of reported adverse effects                      | Drowsiness / Headache / Dizziness / Nausea                                             | 0–4                                 |
|                                  | Adverse Effect Score                                    | Range: 0–4                                                                             | Higher score = more adverse effects |
| Section 8: COVID-19 Impact       | Sleep pattern after COVID-19                            | Improved / No change / Worsened                                                        | 0–2                                 |
|                                  | Melatonin initiated due to COVID-related sleep problems | Yes / No                                                                               | 1 / 0                               |

#### Final Scoring Summary

| Domain              | Score Range | Interpretation             |
|---------------------|-------------|----------------------------|
| Sleep Quality Score | 0–15        | Higher = worse sleep       |
| Self-medication     | 0–6         | Higher = risky behavior    |
| Adverse effects     | 0–4         | Higher = more side effects |