

ASSESSMENT OF GENETIC DIVERGENCE IN TOMATO (*Solanum lycopersicum* L.)

Samima Yeasmin^{1†}, Jyoti Saini^{1†}, Alen Phillip Mathew¹, Shivangi Dubey¹, Ravi Pratap Singh², Deepak Maurya^{1*}

¹School of Agricultural Sciences, GD Goenka University, Sohna-122 103, Gurugram, Haryana, India

²Faculty of Agriculture, Maharishi Markandeshwar (Deemed to be) University, Mullana, Ambala-133 207, Haryana

[†] Both authors contributed equally to this work and share first authorship.

*Corresponding author: mauryadeepak0077@gmail.com, ORCID ID: <https://orcid.org/0000-0001-5469-1749>

Genet. Mol. Res. 25(17s) gmr4196

Received November 14, 2025

Accepted August 24, 2026

Published September 06, 2026

DOI: <https://doi.org/10.4238/jx684559>

ABSTRACT

The present investigation evaluated 25 tomato genotypes during the *Rabi* 2025 season under a randomized block design with three replications. Eighteen quantitative traits related to flowering, fruit development, yield, quality, and plant growth were recorded to estimate genetic diversity using Mahalanobis D² statistics and Tocher's clustering method. The analysis grouped the genotypes into seven distinct clusters, indicating substantial genetic variability. The highest inter-cluster divergence was observed between Clusters II and VII, suggesting that hybridization between these groups may generate superior recombinants with greater heterotic potential. Cluster V exhibited superior vegetative growth, while Cluster III was characterized by desirable fruit quality attributes, including thicker pericarp. Cluster VI recorded the highest fruit yield per plant and per hectare, whereas Cluster VII produced the largest fruits. Genotypes such as Pusa Sheetal, Kashi Amrit, Rosemary, Oaklyan, Punjab Chhuhara, Black Cherry, Pusa Uphar, Money Maker, NF-54, Kashi Hemant, Kashi Anupam, Kashi Sharad, Pusa Rohini, Pusa Sadabahar, Kashi Adarsh, EC 620444, CO-3 and ZS21 were identified as genetically diverse and promising parental lines for hybridization. The findings demonstrate considerable scope for exploiting genetic divergence in tomato breeding programmes aimed at improving fruit yield, quality and overall productivity through heterosis and selection.

KEYWORDS: Tomato, genetic divergence, Mahalanobis D² analysis, Tocher clustering, heterosis breeding

INTRODUCTION

Tomato (*Solanum lycopersicum* L.) is one of the most important vegetable crops cultivated worldwide and belongs to the family Solanaceae. It is a diploid species with a chromosome number of $2n = 24$ (Maurya et al., 2020; Saini et al., 2026). The crop is believed to have originated in the Andean region of South America, where its wild relatives are naturally distributed across Peru, Ecuador, Bolivia, Chile and Colombia. Tomato has evolved from being an ornamental plant once considered poisonous to becoming one of the most extensively cultivated and consumed vegetable crops across the globe (Dorais et al., 2008). Owing to its exceptional nutritional value, it is often described as a "protective food" (Sepat et al., 2013). The fruits are consumed in fresh as well as processed forms, including juice, puree, ketchup, sauces, canned products and several value-added food preparations such as pizza and soups (Maurya et al., 2022; Maurya et al., 2023). It is an excellent source of vitamin C, provitamin A, potassium, dietary fibre, carotenoids, phenolic compounds and other natural antioxidants that contribute significantly to human nutrition (Perveen et al., 2015). Regular dietary intake of tomato has been associated with reduced risks of cardiovascular disorders, type 2 diabetes, certain cancers and neurodegenerative diseases owing to its antioxidant and health-promoting bioactive compounds (Ganesan et al., 2012; Ali et al., 2020). In India, tomato is cultivated throughout the year under diverse agro-climatic conditions and plays a vital role in nutritional security as well as the horticultural economy. In Haryana, the crop occupies an important position among vegetable crops. According to the Third Advance Estimates (2024 - 25), tomato was cultivated over 22.27 thousand hectares, producing 484.37 thousand tonnes with an average productivity

of 21.74 t ha⁻¹ (Department of Agriculture & Farmers Welfare, 2025). Although the state is not among the leading tomato-producing regions of the country, its productivity remains competitive under the agro-climatic conditions of the northern Indo-Gangetic Plains. Adoption of improved cultivars, efficient irrigation management and integrated crop protection practices is expected to further enhance productivity and strengthen horticultural development within the state. Wild tomato species exhibit remarkable adaptation to diverse ecological environments ranging from coastal areas to elevations of nearly 3,600 m in western South America and the Galápagos Islands. Such broad geographical distribution has resulted in substantial genetic and phenotypic diversity among wild taxa, providing valuable genetic resources for crop improvement (Peralta et al., 2008; Tieman et al., 2017). In contrast, cultivated tomato possesses a relatively narrow genetic base because domestication and continuous selection have retained only a limited proportion of the diversity available in wild relatives (Bai et al., 2007; Aflitos et al., 2014; Caramante et al., 2021). Nevertheless, sustained breeding efforts have generated numerous cultivars and hybrids exhibiting considerable variation in fruit size, shape, colour and agronomic performance (Bai et al., 2007; Schouten et al., 2019). The success of any crop improvement programme largely depends on the availability of sufficient genetic variability and the identification of genetically diverse parents possessing desirable traits. The systematic evaluation of germplasm enables plant breeders to understand the genetic potential of available material and to select suitable parents for hybridization, thereby facilitating the development of superior cultivars (Singh et al., 2002). In the context of increasing population pressure and changing climatic conditions, broadening the genetic base of cultivated tomato has become essential for enhancing yield potential, resilience and long-term productivity (Agele et al., 2008; Arena et al., 2020; Francesca et al., 2022; Asare-Addo et al., 2022). Assessment of genetic diversity through multivariate statistical techniques provides an effective approach for identifying divergent genotypes. Among these, the Mahalanobis D² statistic (Mahalanobis, 1936) has been extensively used to estimate genetic divergence by considering multiple quantitative traits simultaneously. The approach was introduced into biological research by Rao (1952) and subsequently applied to plant populations by Nair and Mukharji (1960), followed by its wider application in crop classification and breeding studies (Murthy and Pavate, 1962). Cluster analysis using Tocher's method further facilitates the grouping of genetically similar genotypes, thereby enabling the selection of diverse parents for hybridization programmes. Previous studies have shown that crosses involving genetically divergent parents generally produce higher heterosis and greater breeding gains than those involving closely related genotypes (Ram and Panwar, 1970; Moll and Stuber, 1974). Moreover, characterization of genetic diversity provides a strong foundation for developing improved breeding populations and achieving sustained genetic improvement (Van Ginkel and Ortiz, 2018). Keeping these considerations in view, the present investigation was undertaken to evaluate the genetic divergence among twenty-five tomato genotypes using seventeen important growth, yield and quality traits. The study also aimed to identify genetically diverse and agronomically superior parental lines and to examine the relationships among important traits, thereby providing valuable information for future hybridization programmes and the development of high-yielding, adaptable tomato cultivars.

MATERIAL AND METHODS

The experiment was performed at the Research Farm, GD Goenka University, Sohna, Gurugram, Haryana (28°15' N, 77°04' E; 210-230 m above MSL). A total of twenty-five tomato genotypes were collected from different sources, were evaluated using a Randomized Block Design (RBD) during the *Rabi* season of 2024-2025, as presented in Table 1. The soil of the experimental field was characterized as sandy loam that has a pH range of 6.5-7. Seeds were first sown in seedbeds and subsequently treated with Bavistin for five minutes to protect seedlings from fungal and soil-borne diseases, ensuring better germination and seedling vigor. After 30 days, seedlings were transplanted into plots measuring 2.5 m × 2.5 m, maintaining a spacing of 50 × 50 cm between rows and plants. All recommended agronomic practices were followed to achieve optimum crop performance. Observations were recorded from five randomly selected plants from each genotype in each replication. Data were collected on twelve different characters, namely: number of primary branches, plant height (cm), number of fruits per cluster, number of clusters per plant, number of fruits per plant, number of locules per fruit, harvest duration (days), average fruit weight (g), fruit yield per plant (g), fruit yield per hectare (q).

Statistical Analysis

Genetic divergence among the 25 genotypes was assessed using Mahalanobis D² statistics as described by Rao (1952). The collected data were analyzed using both univariate and multivariate statistical methods with software packages including the Agri Analyze tool [Online tool]. www.agrianalyze.com (Popat et al., 2024) and JMP® Version 19 (<https://www.jmp.com/getstarted>)

RESULTS AND DISCUSSION

Cluster analysis of genotypes based on genetic divergence

Genetic divergence among the twenty-five tomato genotypes was evaluated using Mahalanobis D² analysis, which classified the genotypes into seven distinct clusters, as presented in Table 2 and illustrated by the dendrogram in Fig. 1. The clustering pattern indicated the presence of substantial genetic variability, providing a strong foundation for selecting genetically diverse parents in tomato improvement programmes. Similar observations that genotype within the same cluster are genetically more similar than those belonging to different clusters have been reported by Khapte and Jansirani (2014) and Lekshmi and Celine (2016). Among the seven clusters, Cluster III was the largest, comprising six genotypes (Pusa Uphar, Money Maker, NF-54, Kashi Hemant, Kashi Anupam and Kashi Sharad), followed by Cluster VI with five genotypes (Kashi Aman, EC-620424, Kashi Vishesh, ND-04 and Rishi). Clusters I (Pusa Sheetal, Kashi Amrit, Rosemarry and Oaklyan) and IV (Pusa Rohini, Pusa Sadabahar, Kashi Adarsh and EC-620444) each contained four genotypes, whereas Clusters II (Punjab Chhuhara and Black Cherry), V (ND-02 and ND-08) and VII (CO-3 and ZS21) consisted of two genotypes each. The distribution of genotypes across clusters demonstrated that genetic divergence was independent of geographical origin, indicating that geographical distance alone is not a reliable measure of genetic diversity. The presence of both large and small clusters further reflects the broad genetic variability available within the experimental material. Therefore, the selection of parents for hybridization should be based primarily on genetic divergence rather than geographical distribution to maximize heterosis and broaden the genetic base of breeding populations. Similar findings have been reported by Mulge et al. (2012), Reddy et al. (2013), Kiran et al. (2017), Maurya et al. (2019), Anuradha et al. (2020) and Kumar et al. (2021).

Table 1: List of genotypes (Experimental Materials)

Sr. No.	Genotype	Sources
01	Pusa Sheetal	IIVR, Varanasi
02	Pusa Rohini	IIVR, Varanasi
03	Pusa Uphar	IIVR, Varanasi
04	Pusa Sadabahar	IIVR, Varanasi
05	Kashi Hemant	IIVR, Varanasi
06	Kashi Aman	IIVR, Varanasi
07	Kashi Anupam	IIVR, Varanasi
08	Kashi Amrit	IIVR, Varanasi
09	Kashi Adarsh	IIVR, Varanasi
10	Kashi Sharad	IIVR, Varanasi
11	Kashi Vishesh	IIVR, Varanasi
12	ND-02	ANDUAT, Ayodhya
13	ND-04	ANDUAT, Ayodhya
14	ND-08	ANDUAT, Ayodhya
15	EC-620424	IIVR, Varanasi
16	EC 620444	IIVR, Varanasi
17	Punjab Chhuhara	IIVR, Varanasi
18	Rishi	Local collection from MP
19	CO-3	IIVR, Varanasi
20	Money Maker	IIVR, Varanasi
21	NF-54	Local Collection from MP
22	ZS21	Local Collection from MP
23	Black cherry	Collected from All that Grow
24	Rosemarry	Collected from All that Grow Pvt.
25	Oaklyan	Collected from All that Grow Pvt.

Table 2: Clustering Pattern of 25 tomato genotypes

No. of clusters	Number of genotypes included in the cluster	Name of the genotypes
I	4	Pusa Sheetal, Kashi Amrit, Rosemarry and Oaklyan
II	2	Punjab Chhuhara and Black cherry
III	6	Pusa Uphar, Money Maker, NF-54, Kashi Hemant,

Kashi Anupam and Kashi Sharad		
IV	4	Pusa Rohini, Pusa Sadabahar, Kashi Adarsh and EC 620444
V	2	ND-02 and ND-08
VI	5	Kashi Aman, EC-620424, Kashi Vishesh, ND-04 and Rishi
VII	2	CO-3 and ZS21




Fig. 1: Dendrogram representing the grouping of 25 genotypes of tomato generated using D² cluster analysis

Intra and inter-cluster distances

The intra and inter-cluster distances estimated through Mahalanobis D² analysis provide an effective measure of genetic diversity among the tomato genotypes and assist in identifying suitable parents for hybridization (Table 3). Higher inter-cluster distances indicate greater genetic divergence, whereas lower intra-cluster distances reflect genetic similarity among genotypes within the same cluster. The average intra-cluster D² values ranged from 2.967 to 6.633 (Table 3). Cluster II exhibited the highest intra-cluster distance (6.633), followed by Cluster VI (5.816), Cluster V (5.516), Cluster III (5.416), Cluster IV (3.563) and Cluster I (3.390), while Cluster VII recorded the lowest value (2.967). The higher intra-cluster distance in Cluster II suggests the presence of relatively greater variability among its constituent genotypes. Among the inter-cluster distances, the minimum genetic divergence was observed between Clusters I and IV (3.320), indicating close genetic similarity. In contrast, the maximum inter-cluster distance was recorded between Clusters II and VII (10.829), followed by Clusters IV and VII (10.131), III and VII (9.590) and I and VII (8.849). These highly divergent clusters are expected to provide superior parental combinations for hybridization, as wider genetic divergence enhances the probability of obtaining transgressive segregants through increased recombination and crossing over (Thoday, 1969). Similar observations have been reported by Singh et al. (2008) and Sudeepthi et al. (2020). Therefore, the combined assessment of inter- and intra-cluster distances provides a reliable basis for selecting genetically diverse parents to improve breeding efficiency. Comparable findings have also been reported by Narolia and Reddy (2012), Reddy et al. (2013), Pedapati et al. (2014), Kiran et al. (2017), Maurya et al. (2019), Anuradha et al. (2020) and Kumar et al. (2021).

Table 3: Intra and inter-cluster distances

Cluster	I	II	III	IV	V	VI	VII
I	3.390	4.824	4.198	3.320	4.390	4.369	8.849
II		6.633	4.820	5.257	6.412	6.884	10.829
III			5.416	5.148	4.165	5.060	9.590
IV				3.563	4.434	6.051	10.131
V					5.516	5.243	8.808
VI						5.816	6.122
VII							2.967

(Bold marked values → Intra cluster differences)

Cluster means of eighteen characters and characterization of different clusters of tomato genotypes

The cluster means for eighteen quantitative traits are presented in Table 4. The seven clusters exhibited substantial differences for growth, flowering, fruit, yield and quality attributes, confirming considerable genetic variability among the evaluated tomato genotypes. Cluster mean analysis was useful for identifying clusters with complementary traits that could serve as potential parents in future breeding programmes. Cluster I consisted mainly of early-flowering genotypes, showing the minimum days to first flowering (18.40 days), first fruit setting (13.02 days), and 50% flowering (16.79 days). However, these genotypes had comparatively low TSS, suggesting a possible association between earliness and reduced fruit quality. In contrast, Cluster II comprised late and relatively weak-growing genotypes, with maximum days to first flowering (69.76 days) and 50% flowering (48.87 days), along with poor performance for plant height, branching, locule number, clusters per plant, fruits per plant, and yield. Nevertheless, this cluster recorded the highest TSS, indicating its potential for improving fruit quality. Cluster III was notable for delayed fruit maturity, higher polar diameter (94.39 mm), and greater pericarp thickness (76.08 mm), making it useful for improving fruit firmness and processing attributes. Cluster IV recorded the maximum flowers per cluster (101.68) and generally intermediate performance for other traits, indicating a balanced combination of characteristics. Cluster V showed vigorous growth and prolific fruiting, but smaller fruits and thinner pericarp. Cluster VI recorded the highest fruit yield per plant (128.28 kg) and per hectare (128.28 t ha⁻¹), together with maximum equatorial diameter and locule number, highlighting its value for yield and fruit-size improvement. Cluster VII produced the highest average fruit weight (126.08 g), but fewer fruits per cluster. The contrasting cluster profiles provide opportunities for selecting complementary parents to combine earliness, productivity, fruit size, and quality. The effectiveness of Mahalanobis' D² analysis in identifying genetically divergent parents has also been reported by Kandhola and Panwar (1999), Kuchanur et al. (2009), Bhati et al. (2015), Vennela et al. (2017), Mounika et al. (2020), Roy et al. (2022) and Yeasmin et al. (2023).

Table 4: Cluster means of eighteen characters and showing cluster characteristics of different clusters of tomato genotypes

Characters	Clusters							Mean	Max.	Min
	I	II	III	IV	V	VI	VII			
Days to first flowering	18.40	69.76	34.78	30.68	67.52	50.11	29.57	42.97	69.76	18.40
Days to first fruit setting	13.02	80.23	46.21	44.14	82.72	58.83	37.91	51.87	82.72	13.02
Days to 50% flowering	16.79	48.87	35.03	41.32	41.32	42.08	22.45	35.41	48.87	16.79
Days to first fruit harvest	93.98	92.40	96.60	64.00	29.29	59.58	70.31	72.31	96.60	29.29
Average fruit weight (g)	76.40	63.52	122.40	87.44	59.84	123.87	126.08	94.22	126.08	59.84
Fruit equatorial diameter (mm)	52.67	54.48	117.17	96.08	47.25	121.03	87.03	82.25	121.03	47.25
Polar diameter (mm)	66.10	77.31	94.39	77.31	29.26	74.10	67.70	69.45	94.39	29.26
Locule number	54.75	39.17	114.17	70.33	84.17	123.63	111.86	85.44	123.63	39.17
Pericarp thickness (mm)	41.35	54.19	76.08	62.76	19.94	62.19	48.48	52.14	76.08	19.94
Number of flowers/clusters	75.41	64.59	49.14	101.68	36.78	53.46	67.68	64.10	101.68	36.78
Number of clusters/plants	82.28	22.04	65.14	58.77	101.38	91.39	54.36	67.91	101.38	22.04
Number of fruits /clusters	46.05	39.55	30.88	41.72	54.72	34.78	20.04	38.25	54.72	20.04
Number of fruits/plants	126.08	48.80	85.60	94.80	155.52	119.45	52.48	97.53	155.52	48.80
Number of primary branches/plants	25.16	9.97	49.75	44.69	62.05	46.86	42.52	40.14	62.05	9.97
Plant height (cm)	105.12	40.02	89.45	70.76	134.05	113.80	61.72	87.84	134.05	40.02
Fruit yield/plant (kg)	96.64	48.80	113.81	92.96	89.28	128.28	74.56	92.05	128.28	48.80
Fruit yield/ha (t/ha)	96.64	48.80	113.81	92.96	89.28	128.28	74.56	92.05	128.28	48.80
TSS (Total Soluble Solids)	18.40	69.76	34.78	30.68	67.52	50.11	29.57	42.97	69.76	18.40

CONCLUSION

According to present investigation demonstrated considerable genetic diversity among the twenty-five tomato genotypes, as shown by their distribution into seven distinct clusters using Mahalanobis D² analysis. The clustering pattern was independent of geographical origin, highlighting the importance of genetic divergence over geographical distance for selecting superior breeding materials. Wide inter-cluster distances indicated the availability of highly diverse parental combinations, while cluster mean analysis identified complementary traits associated with earliness, fruit yield, fruit size, and quality. Cluster VI was distinguished by its superior yield potential, Cluster VII by high fruit weight, Cluster I by earliness and Cluster II by enhanced soluble solids content. These findings suggest that hybridization between genetically divergent clusters possessing desirable attributes would enhance the probability of obtaining superior recombinants with improved productivity and fruit quality. The identified diverse genotypes provide valuable genetic resources for developing high-yielding and quality tomato cultivars in future breeding programmes.

Acknowledgment

The authors thank GD Goenka University for providing facilities. No financial support or funding was received for this research.

Author contribution

SY and DM conceptualized and drafted the manuscript. JS, APM, SD and RPS assisted in preparing the manuscript. All authors read the final version of the manuscript, provided necessary suggestions and approved it for publication.

Conflict of interest:

On behalf of all authors, the corresponding author states that there is no conflict of interest. The corresponding author confirms that the submitted manuscript is original, unpublished and not under simultaneous consideration by another journal.

REFERENCES

1. Aflitos, S., Schijlen, E., de Jong, H., de Ridder, D., Smit, S., Finkers, R., Wang, J., Zhang, G., Li, N. and Mao, L., 2014. Exploring genetic variation in the tomato (*Solanum* section *Lycopersicon*) clade by whole-genome sequencing. *The Plant Journal*, 80(1), pp.136-148. <https://doi.org/10.1111/tpl.12616>
2. Agele, S.O., Adeniji, I.A., Alabi, E.O. and Olabomi, A., 2008. Responses of growth, yield and N use efficiency of selected tomato cultivars to variations in hydrothermal regimes of the cropping seasons in a tropical rainforest zone of Nigeria. *Journal of Plant Interactions*, 3(4), pp.273-285. <https://doi.org/10.1080/17429140802112>.
3. Ali, M.Y., Sina, A.A.I., Khandker, S.S., Neesa, L., Tanvir, E.M., Kabir, A., Khalil, M.I. and Gan, S.H., 2020. Nutritional composition and bioactive compounds in tomatoes and their impact on human health and disease: A review. *Foods*, 10(1), p.45. <https://doi.org/10.3390/foods10010045>
4. Anuradha, B., Reddy, P.S.K.R., Sudini, H.A.R.I.K.I.S.H.A.N. and Geetha, A., 2020. Genetic divergence for yield and yield attributes in tomato (*Solanum lycopersicum* L.). *Green Farming*, 11(4&5), pp.293-298.
5. Arena, C., Conti, S., Francesca, S., Melchionna, G., Hájek, J., Barták, M., Barone, A. and Rigano, M.M., 2020. Eco-physiological screening of different tomato genotypes in response to high temperatures: A combined field-to-laboratory approach. *Plants*, 9(4), p.508. <https://doi.org/10.3390/plants9040508>
6. Asare-Addo, D.C., Amisah, J.N., Ofori, P.A., Owusu-Nketia, S., Opoku-Agyemang, F. and Nkansah, G.O., 2022. Evaluation of agronomic performances and fruit quality of improved tomato (*Solanum lycopersicum* L.) lines under greenhouse conditions. *Journal of Agriculture and Food Research*, 9, p.100360. <https://doi.org/10.1016/j.jafr.2022.100360>
7. Bai, Y. and Lindhout, P., 2007. Domestication and breeding of tomatoes: what have we gained and what can we gain in the future?. *Annals of Botany*, 100(5), pp.1085-1094. <https://doi.org/10.1093/aob/mcm150>
8. Bhati, P.K., Singh, S.K., Dhurai, S.Y. and Sharma, A., 2015. Genetic divergence for quantitative traits in rice germplasm. *Electronic Journal of Plant Breeding*, 6(2), pp.528-534.
9. Caramante, M., Rouphael, Y. and Corrado, G., 2021. The genetic diversity and structure of tomato landraces from the Campania region (Southern Italy) uncover a distinct population identity. *Agronomy*, 11(3), p.564. <https://doi.org/10.3390/agronomy11030564>
10. Comstock, R.E. and Robinson, H.F., 1952. Genetic parameters, their estimation and significance. In *Proceedings of the 6th International Grassland Congress*.
11. Department of Agriculture & Farmers Welfare, Government of Haryana. 2025. Third advance

- estimates of area and production of horticultural crops for 2024-25. Government of Haryana, Chandigarh, India
12. Dorais, M., Ehret, D.L. and Papadopoulos, A.P., 2008. Tomato (*Solanum lycopersicum*) health components: from the seed to the consumer. *Phytochemistry reviews*, 7(2), pp.231-250. <https://doi.org/10.1007/s11101-007-9085-x>
 13. Francesca, S., Vitale, L., Arena, C., Raimondi, G., Olivieri, F., Cirillo, V., Paradiso, A., de Pinto, M.C., Maggio, A., Barone, A. and Rigano, M.M., 2022. The efficient physiological strategy of a novel tomato genotype to adapt to chronic combined water and heat stress. *Plant Biology*, 24(1), pp.62-74. <https://doi.org/10.1111/plb.13339>
 14. Ganesan, M., Rajesh, M., Solairaj, P. and Senthilkumar, T., 2012. Tomato as a pioneer in health management. *Int. J. Pharm. Chem. Biol. Sci*, 2(3), pp.210-217.
 15. JMP Statistical Discovery LLC, 2025-2026. *JMP® 19 Automation Reference*. Cary, NC: JMP Statistical Discovery LLC (<https://www.jmp.com/getstarted>)
 16. Kandhola, S.S. and Panwar, D.V.S., 1999. Genetic divergence in rice. *Annals of Biology*, 15(1), 35-39.
 17. Khapte, P.S. and Jansirani, P., 2014. Genetic diversity studies in tomato (*Solanum lycopersicum* L.) genotypes. *Trends in Biosciences*, 7(15), pp.1958-1961.
 18. Kiran, M.L., Mishra, H.N., Mishra, A., Nandi, A., Tripathy, P., Senapati, N. and Pradhan, K., 2017. Genetic divergence and cluster analysis in tomato (*Solanum lycopersicum* L.). *Environment and Ecology*, 35(4B), pp.3167-3171.
 19. Kuchanur, P.H., Naresh Dronavalli, N.D. and Vijayakumar, A.G., 2009. Genetic variability and divergence in 'New Plant Type rice genotypes. *Crop Improvement*, 36, 20-24.
 20. Kumar, J., Dixit, P.S., Yadav, L., Kumar, M., Singh, B.P. and Tomar, S., 2021. Studies about genetic divergence in 40 genotypes of Tomato (*Solanum lycopersicum* L.). *Journal of Pharmacognosy and Phytochemistry*, 10(1), pp.2290-2292.
 21. Lekshmi, S.L. and Celine, V.A., 2016. Genetic diversity studies in tomato (*Solanum lycopersicum* L.) under protected conditions. *International Journal of Current Microbiology and Applied Science*, 5(4), pp.212-217.
 22. Mahalanobis, P. C., 1936. On the generalized distance in statistics. *Proceedings of the National Academy of Sciences, India*, 2, 49-55.
 23. Maurya, D., Akhtar, S., Chattopadhyay, T., Kumar, R., Sahay, S., Sangam, S., Kumari, N. and Siddiqui, M.W., 2022. Genetic variability and character association in tomato (*Solanum lycopersicum* L.). *Bangladesh Journal of Botany*, 51(4), pp.747-757. <https://doi.org/10.3329/bjb.v51i4.63494>
 24. Maurya, S., Singh, A.K., Singh, S.K. and Kumawat, O.P., 2019. Study of genetic divergence in tomato (*Solanum lycopersicum* L.). *Int. J. Curr. Microbiol. App. Sci*, 8(8), pp.2429-2436.
 25. Maurya, D., A. Mukherjee, Bhagyashree, S. Sangam, R. Kumar, S. Akhtar, T. Chattopadhyay, 2023. Marker-assisted stacking of *Ty3*, *Mi1.2* and *Ph3* resistance alleles for leaf curl, root knot and late blight diseases in tomato. *Physiol. Mol. Biol. Plants*, 29(1):121-129. <https://doi.org/10.1007/s12298-022-01277-2>
 26. Maurya, D., B. Shree, S. Akhtar, T. Chattopadhyay, 2020. Exploring allelic status of selected disease resistance genes in a set of tomato genotypes using gene-linked molecular markers. *J. Crop Weed*, 16(1): 236-241. <https://doi.org/10.22271/09746315.2020.v16.i1.1299>
 27. Moll, R. H. and Stuber, C. W., 1974, Quantitative genetics empirical results relevant to plant breeding. *Adv. Agron.*, 26: 277-313.
 28. Mounika, K., Singh, S.K., Pandey, V., Singh, D.K., Khaire, A.R., Habde, S. and Majhi, P.K., 2020. Study of genetic divergence in rice (*Oryza sativa* L.) genotypes with high grain zinc using Mahalanobis' D² analysis. *Electronic Journal of Plant Breeding*, 11(02), pp.367-372.
 29. Mulge, R., Mahendrakar, P. and Prashanth, S.J., 2012. Prediction of heterosis based on genetic divergence in tomato. *Indian Journal of Horticulture*, 69(4), pp.530-535.
 30. Murty, G.S. and Pavate, M.V., 1962. Studies on quantitative inheritance in *Nicotiana tabacum* L. Varietal classification and selection by multivariate analysis. *Indian Journal of Genetics and Plant Breeding*, 22, 68-72.
 31. Nair, K.R. and Mukerji, H.K., 1960. Classification of natural and plantation teak (*Tectona grandis*) grown at different localities of India and Burma with respect to its physical and mechanical properties. *Sankhyā: The Indian Journal of Statistics*, pp.1-20. <https://www.jstor.org/stable/25048443>
 32. Nalla, M.K., Rana, M.K., Singh, S.J., Sinha, A.K., Reddy, P.K. and Mohapatra, P.P., 2014. Genetic divergence analysis in tomato (*Solanum lycopersicum* L.). *Electronic Journal of Plant Breeding*, 5, 517-525.
 33. Peralta, I.E., Spooner, D.M. and Knapp, S., 2008. Taxonomy of wild tomatoes and their relatives (*Solanum* section *Lycopersicon*, Solanaceae). *Systematic Botany Monographs*, 84, 1-186. <https://doi.org/10.1600/036364408784571545>

34. Perveen, R., Suleria, H.A.R., Anjum, F.M., Butt, M.S., Pasha, I. and Ahmad, S., 2015. Tomato (*Solanum lycopersicum* L.) carotenoids and lycopene chemistry; metabolism, absorption, nutrition, and allied health claims—A comprehensive review. *Critical reviews in food science and nutrition*, 55(7), pp.919-929. <https://doi.org/10.1080/10408398.2012.657809>
35. Popat, R., Patel, H. and Popat, P., 2024. Agri Analyze [Online tool]. www.agrianalyze.com
36. Ram, J. and Panwar, D.V.S., 1970. Intraspecific divergence in rice. *Indian Journal of Genetics and Plant Breeding (The)*, 30(1), pp.1-10.
37. Rao, C.R., 1952. Advanced statistical methods in biometric research. *John Wiley and Sons*.
38. Reddy, B.R., Begum, H., Sunil, N. and Reddy, T.M., 2013. Genetic divergence studies in exotic collections of tomato (*Solanum lycopersicum* L.). *International Journal of Agricultural Sciences*, 9(2), pp.588-592.
39. Roy, D., Gaur, A.K. and Pandey, I.D., 2022. Deciphering genetic diversity in 'Antenna Panel' genotypes of IRR1's Global Rice Array-IV for yield traits in Indo-Gangetic Plains. *Electronic Journal of Plant Breeding*, 13(2), pp.425-431. <https://doi.org/10.37992/2022.1302.042>
40. Saini, J., Yeasmin, S., Maurya, D. and Jatav, V., 2026. Genetic variability and trait relationships in tomato germplasm under semi-arid conditions of Haryana. *Journal of Applied Horticulture*, 28(2), pp.335-341. <https://doi.org/10.37855/jah.2026.v28i02.53>
41. Schouten, H.J., Tikunov, Y., Verkerke, W., Finkers, R., Bovy, A., Bai, Y. and Visser, R.G., 2019. Breeding has increased the diversity of cultivated tomato in the Netherlands. *Frontiers in plant science*, 10, p.1606. <https://doi.org/10.3389/fpls.2019.01606>
42. Sepat, N.K., Sepat, S.R., Sepat, S. and Kumar, A., 2013. Energy use efficiency and cost analysis of tomato under greenhouse and open field production system at Nubra valley of Jammu and Kashmir. *International Journal of Environmental Sciences*, 3(4), pp.1233-1241.
43. Singh, A. and Singh, H.N., 1976. Genetic divergence in chilli. *Indian Journal of Genetics and Plant Breeding (The)*, 36(3), pp.425-430.
44. Singh, A.K., Sharma, J.P., Kumar, S. and Chopra, S., 2008. Genetic divergence in tomato (*Lycopersicon esculentum* Mill.). *Journal of Research, SKUAST-J*, 7(1), pp.105-110.
45. Singh, J.K., Singh, J.P., Jain, S.K. and Aradhana Joshi, A.J., 2002. Studies on genetic variability and its importance in tomato (*Lycopersicon esculentum* Mill.). *Progressive Horticulture*, 34, 77-79.
46. Singh, R.K. and Chaudhary, B.D., 1977. Biometrical methods in quantitative genetic analysis. Kalyani Publishers.
47. Sudeepthi, K., Srinivas, T., Kumar, B.N.V.S.R.R., Jyothula, D.P.B. and Umar, S.N., 2020. Genetic divergence studies for anaerobic germination traits in rice (*Oryza sativa* L.). *Current Journal of Applied Science and Technology*, 39(1), pp.71-78. <https://doi.org/10.9734/cjast/2020/v39i130482>
48. Thoday, J.M., 1960. Effects of disruptive selection: III. Coupling and repulsion. *Heredity*, 14, 35-39.
49. Tieman, D., Zhu, G., Resende Jr, M.F., Lin, T., Nguyen, C., Bies, D., Rambla, J.L., Beltran, K.S.O., Taylor, M., Zhang, B. and Ikeda, H., 2017. A chemical genetic roadmap to improved tomato flavor. *Science*, 355(6323), pp.391-394.
50. Van Ginkel, M. and Ortiz, R., 2018. Cross the best with the best, and select the best: HELP in breeding selfing crops. *Crop Science*, 58(1), pp.17-30. <https://doi.org/10.2135/cropsci2017.05.0270>
51. Vennela, P.R., Singh, S.K., Singh, R., Gayatonde, V. and Singh, D.K., 2017. Genetic divergence studies in rice (*Oryza sativa* L.) for yield and yield related traits. *Vegetos-An International Journal of Plant Research*, 30(special), pp.191-195.
52. Yeasmin, S., Dukpa, P., Maurya, D., Jatav, V. and Dwary, S., 2023. Genetic variability, heritability and genetic advance for yield and yield contributing traits in tomato (*Solanum lycopersicum* L.). *International Journal of Advanced Biochemistry Research*, 7(2), 207-211. <https://doi.org/10.33545/26174693.2023.v7.i2Sc.214>