

MOLECULAR GENETICS OF LIPID METABOLISM AND ITS ROLE IN NEURODEGENERATIVE DISEASES

Dr Shrikanth Metri¹, Dr Suhasini Atharga², Dr. Sonia Rani³, Gargi Mishra⁴, Mukesh Krishnakumar⁵, Dr. C. Ramya⁶

¹Associate Professor, Dept of General Medicine, JN Medical college KAHER, Belagavi, Karnataka, India

²Assistant Professor of Cardiology, JN Medical College KAHER, Belagavi, Karnataka, India

³Prof. Mental Health Nursing, Department of Galgotias School of Nursing, Galgotias University, Greater Noida, Uttar Pradesh, India

⁴Ass. Professor, SOAH, DRIEMS University, Kataka, Odisha, India

⁵Assistant professor, Chennai Institute of Technology, Kanchipuram, Tamilnadu

⁶Senior Lecturer, Oral and Maxillofacial Pathology & Oral Microbiology, Sree Balaji Dental College and Hospital, BIHER, Chennai, Tamil Nadu, India

ABSTRACT:

The molecular regulation of lipid metabolism has emerged as an important area of investigation in understanding the biological mechanisms underlying neurodegenerative diseases. Lipids are essential components of neuronal membranes, myelin, intracellular organelles, and signaling systems, while their synthesis, transport, modification, storage, and degradation must remain tightly coordinated to preserve neuronal function. Recent research indicates that disturbances in lipid homeostasis can influence processes associated with neurodegeneration, including membrane instability, mitochondrial dysfunction, oxidative stress, neuroinflammation, impaired lysosomal activity, altered protein processing, and abnormal lipid droplet accumulation. This study examines the molecular genetic mechanisms that regulate lipid metabolism and explores their contribution to the development and progression of major neurodegenerative disorders, with particular emphasis on Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis, and related neurological conditions. The investigation considers the roles of genes and regulatory pathways involved in cholesterol transport, fatty-acid metabolism, sphingolipid synthesis, phospholipid remodeling, lipid storage, and lysosomal lipid processing. Particular attention is given to genetic variations affecting lipid-associated proteins and cellular pathways, including apolipoprotein-mediated transport, lysosomal function, lipid droplet regulation, and interactions between neurons and glial cells. The study further examines how inherited or acquired alterations in these molecular pathways may modify cellular responses to aging, metabolic stress, inflammation, and abnormal protein accumulation. Emerging evidence suggests that lipid metabolism is not an isolated metabolic process but is closely connected with mitochondrial activity, immune signaling, membrane trafficking, and neuronal survival. Understanding these interconnected mechanisms may help explain why disturbances in specific lipid pathways produce different neurological consequences across disease conditions. The study also considers the potential of molecular profiling, lipidomics, genetic analysis, and pathway-based approaches for identifying disease-associated biomarkers and therapeutic targets. By integrating molecular genetics with neurobiology and lipid metabolism, this research highlights the importance of maintaining lipid homeostasis for neural health and provides a framework for investigating personalized therapeutic strategies aimed at modifying lipid-related mechanisms involved in neurodegeneration.

KEYWORDS: Molecular Genetics, Lipid Metabolism, Neurodegenerative Diseases, Lipid Homeostasis, Neurobiology

INTRODUCTION

Lipid metabolism is a fundamental biological process that supports the structural, energetic, and signaling requirements of the nervous system. Lipid molecules constitute a major portion of neuronal and glial membranes, contribute to myelin formation, participate in intracellular signaling, and influence membrane trafficking, synaptic activity, and cellular communication. Unlike many other tissues, the brain contains a highly diverse lipid composition that must be carefully maintained despite the relatively restricted exchange of many molecules between the circulation and the central nervous system. Cholesterol, phospholipids, sphingolipids, fatty acids, and their metabolites perform distinct but interconnected functions in maintaining neuronal integrity. Their synthesis, transport, remodeling, storage, and degradation are controlled by coordinated molecular pathways involving numerous enzymes, transport proteins, receptors, transcriptional regulators, and intracellular organelles. Genetic variation in these pathways can alter the quantity, distribution, or biological activity of individual lipid species and may consequently affect neural function. This relationship has attracted increasing attention because disturbances in lipid homeostasis have been observed in several neurodegenerative conditions. Changes in lipid composition can influence membrane properties, mitochondrial activity, oxidative balance, inflammatory signaling, protein trafficking, and the functioning of lysosomal and autophagic systems. Such disturbances may become particularly important during aging, when the capacity of cells to maintain metabolic balance and repair molecular damage can decline. The study of molecular genetics therefore provides an

important perspective for understanding how inherited variants, regulatory changes, and interactions among lipid-related genes contribute to susceptibility to neurological disorders. Rather than considering lipid abnormalities simply as secondary consequences of neuronal injury, current research increasingly investigates whether alterations in lipid metabolism can participate directly in disease initiation and progression.

The genetic regulation of lipid metabolism involves interconnected pathways operating within neurons, astrocytes, microglia, oligodendrocytes, and other cells of the nervous system. Genes involved in cholesterol synthesis and transport, fatty-acid processing, sphingolipid metabolism, phospholipid remodeling, lipid storage, and lysosomal degradation can influence cellular responses to physiological and pathological stress. Cholesterol is particularly important because it contributes to membrane organization and synaptic function, while its distribution within the brain is regulated through specialized transport and turnover mechanisms. Similarly, sphingolipids and phospholipids participate in membrane architecture and signaling pathways, and alterations in their metabolism can affect neuronal survival and communication. Genetic differences in lipid-associated proteins may consequently modify cellular physiology even when their effects are not immediately apparent under normal conditions. One widely investigated example is variation in genes associated with lipid transport and neuronal homeostasis, where particular genetic variants have been linked with altered susceptibility to neurodegenerative disorders. However, genetic influence is complex and cannot generally be reduced to a single gene or pathway. Interactions between multiple genetic factors, age-related changes, environmental exposures, metabolic status, and cellular stress can influence how lipid disturbances develop and affect the nervous system. In addition, lipid metabolism is closely connected with mitochondrial function and energy production. Disrupted fatty-acid handling or abnormal lipid accumulation can interfere with mitochondrial processes, increase oxidative stress, and alter cellular energy balance. Lysosomes and autophagic pathways are similarly important because they participate in the degradation and recycling of cellular components, including lipid-containing structures. When these systems become dysfunctional, potentially harmful lipid species and damaged organelles may accumulate, creating conditions that can further disturb neuronal homeostasis. Understanding these interconnected mechanisms is therefore essential for establishing how molecular changes in lipid metabolism may contribute to long-term neuronal damage.

The relationship between lipid metabolism and neurodegeneration is particularly relevant to disorders such as Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis, and related neurological conditions. Although these diseases differ in their clinical characteristics and pathological features, they share several cellular disturbances involving protein accumulation, mitochondrial dysfunction, neuroinflammation, impaired intracellular trafficking, and progressive neuronal loss. Lipid metabolism can intersect with each of these processes. Altered membrane composition may influence the processing, movement, and aggregation of proteins, while changes in lipid signaling can affect inflammatory responses mediated by microglia and other immune-related cells in the nervous system. Lipid abnormalities may also influence the interaction between neurons and glial cells, thereby modifying the local environment in which neurodegenerative processes occur. In Alzheimer's disease, for example, cholesterol and lipid transport mechanisms have been investigated in relation to amyloid processing, neuronal membrane organization, and inflammatory responses. In Parkinson's disease, alterations involving lipid metabolism and lysosomal pathways have attracted attention because of their potential relationship with α -synuclein handling and dopaminergic neuronal vulnerability. In other neurodegenerative disorders, abnormalities in lipid storage, membrane trafficking, and energy metabolism may contribute to cellular stress and impaired neuronal maintenance. Importantly, these relationships are biologically interconnected rather than disease-specific in a simple manner. A disturbance in one lipid pathway can influence other metabolic networks through feedback mechanisms and changes in cellular signaling. The resulting effects may vary according to cell type, genetic background, disease stage, and environmental conditions. This complexity makes molecular investigation particularly important because conventional measurements of total lipid concentration may not reveal changes in specific lipid species, their cellular location, or their functional significance. Genetic approaches combined with transcriptomic, proteomic, lipidomic, and biochemical analyses can provide a more detailed understanding of these relationships and help identify molecular signatures associated with disease susceptibility and progression.

Against this background, the present study examines the **molecular genetics of lipid metabolism and its role in neurodegenerative diseases**, focusing on the biological pathways through which genetic and metabolic alterations may influence neuronal health. The study considers genes involved in lipid synthesis, transport, degradation, and signaling and evaluates their relationship with cellular mechanisms relevant to neurodegeneration. Particular attention is given to the interaction between lipid homeostasis, mitochondrial dysfunction, oxidative stress, neuroinflammation, lysosomal activity, membrane dynamics, and abnormal protein processing. The investigation also recognizes that genetic predisposition does not operate independently and that lipid-related neurological outcomes can emerge from complex interactions between inherited variants, cellular aging, metabolic conditions, and environmental influences. Advances in high-throughput sequencing, genome-wide association studies, lipidomics, single-cell analysis, and molecular profiling have created new opportunities to investigate these relationships at greater resolution. Such approaches can help identify genetic variants and lipid signatures associated with disease risk while also providing information about the biological pathways affected by those changes. A clearer understanding of these mechanisms may support the development of biomarkers for early disease detection and contribute to the identification of therapeutic targets directed

toward lipid homeostasis rather than solely toward downstream pathological effects. The significance of this field therefore extends beyond understanding lipid abnormalities as biochemical observations; it involves determining how genetic regulation of lipid pathways interacts with the fundamental processes required for neuronal survival and function. Investigating these relationships may provide a more integrated view of neurodegenerative disease biology and encourage the development of research strategies that connect molecular genetics, lipid biology, cellular neuroscience, and therapeutic discovery.

METHODOLOGY

The present study adopted an integrative molecular-genetics research approach to investigate the relationship between lipid metabolism, genetic regulation, and the development and progression of neurodegenerative diseases. The methodology was designed to combine evidence from recent peer-reviewed literature with molecular and pathway-level interpretation of genes associated with lipid synthesis, transport, remodeling, degradation, signaling, and cellular lipid storage. This approach was selected because lipid metabolism in the central nervous system is not controlled by a single biochemical pathway but by interconnected processes operating across neurons, astrocytes, microglia, oligodendrocytes, and other neural cell types. Current evidence indicates that lipid remodeling, oxidation, lipid-droplet metabolism, phospholipid processing, and intercellular lipid transfer are closely connected with neuronal and glial homeostasis and may become disrupted during neurodegeneration. The investigation therefore considered both individual lipid-related genes and the biological networks in which they participate. Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis, and selected related neurodegenerative conditions were considered as major disease contexts because abnormalities in cholesterol, sphingolipids, phospholipids, fatty acids, lysosomal metabolism, and lipid transport have been repeatedly investigated in these disorders.

RESEARCH DESIGN

A structured integrative review design was used to identify, organize, compare, and interpret evidence concerning the molecular genetics of lipid metabolism and neurodegeneration. The research process consisted of five interconnected stages: identification of relevant literature, extraction of lipid-associated genetic information, classification of metabolic pathways, mapping of genetic alterations to neurodegenerative mechanisms, and synthesis of the findings. Rather than treating genetic variants and lipid abnormalities as independent observations, the methodology examined their functional relationships with membrane integrity, mitochondrial activity, oxidative stress, lysosomal function, neuroinflammation, protein aggregation, and neuronal survival.

S. No.	Research Component	Methodological Approach
1	Research design	Integrative molecular-genetics literature analysis
2	Primary focus	Genes regulating lipid metabolism and homeostasis
3	Major lipid classes	Cholesterol, fatty acids, phospholipids, sphingolipids, glycerolipids and lipid-storage components
4	Major diseases	Alzheimer's disease, Parkinson's disease, ALS and selected related neurodegenerative disorders
5	Biological levels	Gene, protein, pathway, lipid, cell and disease phenotype
6	Analytical perspective	Molecular, biochemical, cellular and disease-level integration
7	Principal outcome	Identification of relationships between lipid-associated genetic mechanisms and neurodegeneration

Literature Identification and Database Selection

Relevant scientific publications were identified through systematic searches of major biomedical and multidisciplinary databases. PubMed/MEDLINE, Web of Science, Scopus, Google Scholar, and selected publisher databases were considered appropriate sources for locating peer-reviewed studies, review articles, genetic investigations, mechanistic studies, and recent advances in lipid biology. Greater emphasis was placed on recent literature, particularly studies published from 2020 onward, while older landmark studies were retained when they provided essential information concerning established genes, pathways, or disease mechanisms. Recent reviews have emphasized the importance of lipid metabolism and remodeling within the CNS and have identified dysregulated lipid pathways as relevant to neurodegenerative disease mechanisms.

The search strategy employed combinations of keywords and Boolean operators. Core search terms included "lipid metabolism," "lipid homeostasis," "molecular genetics," "genetic variants," "lipid-associated genes," "cholesterol metabolism," "fatty acid metabolism," "sphingolipid metabolism," "phospholipid metabolism," "lysosomal lipid metabolism," "neurodegeneration," "Alzheimer's disease," "Parkinson's disease," "amyotrophic lateral sclerosis," "lipidomics," and "neuroinflammation." Search combinations were modified according to the requirements of individual databases.

S. No.	Search Concept	Representative Search Terms
1	Molecular genetics	“molecular genetics” AND “lipid metabolism”
2	Lipid homeostasis	“lipid homeostasis” AND “brain” OR “CNS”
3	Cholesterol	“cholesterol metabolism” AND “neurodegeneration”
4	Fatty acids	“fatty acid metabolism” AND “neurodegenerative disease”
5	Sphingolipids	“sphingolipid” AND “genetic” AND “neurodegeneration”
6	Phospholipids	“phospholipid metabolism” AND “neuronal dysfunction”
7	Alzheimer’s disease	“lipid metabolism” AND “Alzheimer’s disease”
8	Parkinson’s disease	“lipid metabolism” AND “Parkinson’s disease”
9	Genetic risk	“lipid genes” AND “genetic variants” AND “neurodegeneration”
10	Cellular mechanisms	“lipid metabolism” AND “mitochondria” AND “neuroinflammation”

Selection of Relevant Studies

Studies were selected according to predefined inclusion and exclusion criteria. The selection process was designed to minimize the inclusion of publications that addressed lipid metabolism only superficially or without a meaningful relationship to molecular genetics or neurodegeneration. Peer-reviewed original research articles, systematic reviews, narrative reviews, genetic association studies, genome-wide association studies, molecular studies, lipidomic investigations, and mechanistic studies were considered. Publications were screened initially according to title and abstract, followed by evaluation of the full text where sufficient information was available.

Selection Category	Inclusion Criteria	Exclusion Criteria
Publication type	Peer-reviewed research and review articles	Editorials, opinion pieces and non-scholarly material
Research focus	Lipid metabolism, genetics and neurodegeneration	Studies unrelated to neural lipid biology
Genetic relevance	Genes, variants, expression or molecular regulation	Studies without meaningful genetic or molecular information
Disease relevance	Neurodegenerative or closely related neurological disorders	Conditions unrelated to the research objectives
Evidence	Mechanistic, clinical, genetic, biochemical or lipidomic evidence	Unsupported commentary or duplicate information
Publication period	Priority to recent literature; landmark older studies retained where necessary	Obsolete material without continuing scientific relevance
Language	Primarily English-language scientific literature	Publications without sufficient accessible scientific information

Extraction of Molecular and Genetic Information

After study selection, relevant information was systematically extracted into a structured evidence matrix. Data extraction focused on the identification of genes, gene products, lipid classes, biochemical pathways, cellular locations, disease associations, and proposed molecular mechanisms. Particular attention was given to genes involved in cholesterol transport and synthesis, fatty-acid metabolism, sphingolipid processing, phospholipid remodeling, lysosomal function, lipid-droplet regulation, and intracellular lipid trafficking.

The extracted information was organized according to the relationship between genetic alteration and lipid-related phenotype. Genetic evidence included common variants, rare variants, loss-of-function changes, disease-associated mutations, altered gene expression, and pathway-level genetic associations where reported. Recent genetic studies have highlighted the contribution of lipid-related pathways to neurodegenerative disease susceptibility, while research on rare variants has also expanded understanding of genes involved in Alzheimer’s disease biology.

S. No.	Molecular Domain	Information Extracted
1	Gene identity	Gene name and recognized biological function
2	Genetic variation	Variant, mutation or reported genetic association
3	Gene expression	Increased, decreased or altered expression
4	Protein function	Enzymatic, transport, receptor or regulatory activity
5	Lipid class	Sterol, fatty acid, sphingolipid, phospholipid or glycerolipid
6	Cellular location	Membrane, mitochondria, lysosome, endoplasmic reticulum or lipid droplet
7	Disease relationship	Reported association with neurodegenerative disease

8	Molecular consequence	Altered synthesis, transport, degradation or signaling
9	Cellular consequence	Oxidative stress, inflammation, mitochondrial or lysosomal dysfunction
10	Pathological consequence	Protein aggregation, synaptic dysfunction or neuronal injury

Classification of Lipid-Associated Genes and Pathways

The identified genes were classified according to their principal contribution to lipid biology. This classification allowed the study to compare molecular functions across different neurodegenerative diseases. Cholesterol-associated genes were considered separately from genes involved in fatty-acid, sphingolipid, phospholipid, lysosomal, and lipid-droplet metabolism. Genes with overlapping functions were assigned to multiple functional categories when necessary because lipid metabolic pathways are highly interconnected.

The study also considered the relationship between lipid metabolism and cellular organelles. The endoplasmic reticulum is central to lipid synthesis and remodeling, mitochondria participate in lipid oxidation and energy metabolism, lysosomes contribute to degradation and recycling, and lipid droplets provide dynamic storage sites for neutral lipids. Disturbances in these systems can influence one another and may produce a progressive metabolic imbalance. Recent research has particularly emphasized lipid transfer between neurons and glial cells and the role of microglial lipid dysregulation in neuroinflammation.

Disease-Specific Molecular Mapping

The extracted genetic and metabolic information was subsequently mapped against major neurodegenerative diseases. For Alzheimer's disease, the analysis focused on lipid transport, cholesterol regulation, membrane remodeling, lysosomal function, inflammatory signaling, and interactions with pathological protein processing. For Parkinson's disease, attention was given to fatty-acid and sphingolipid metabolism, lysosomal pathways, mitochondrial dysfunction, α -synuclein-associated mechanisms, and oxidative stress. For ALS and other neurodegenerative disorders, lipid homeostasis, membrane maintenance, energy metabolism, and cellular stress pathways were evaluated according to the available evidence.

Disease	Major Lipid-Related Areas Examined	Associated Cellular Mechanisms
Alzheimer's disease	Cholesterol, phospholipids, sphingolipids, lipid transport	Protein processing, inflammation, membrane dysfunction and synaptic injury
Parkinson's disease	Fatty acids, sphingolipids, cholesterol and lipid storage	α -synuclein biology, mitochondrial dysfunction and oxidative stress
ALS	Fatty acids, membrane lipids and energy-related lipid pathways	Neuronal stress, mitochondrial abnormalities and inflammation
Huntington's disease	Membrane and energy-associated lipid pathways	Cellular stress, mitochondrial dysfunction and neuronal vulnerability
Related disorders	Lysosomal and lipid-storage pathways	Autophagy, lipid accumulation and impaired cellular clearance

Evidence concerning Alzheimer's and Parkinson's disease indicates that abnormal cholesterol, sphingolipid, and glycerophospholipid metabolism can interact with protein aggregation, mitochondrial dysfunction, inflammatory signaling, and membrane abnormalities. Similarly, recent work has investigated fatty-acid dysregulation as an important component of Parkinson's disease biology.

Molecular Pathway Analysis

The study examined the biological consequences of altered lipid metabolism at several interconnected levels. First, changes in lipid synthesis or degradation were evaluated for their potential effects on membrane composition. Second, altered lipid transport and intracellular trafficking were considered in relation to organelle function. Third, the study examined whether lipid imbalance could contribute to oxidative stress, mitochondrial dysfunction, neuroinflammation, lysosomal impairment, and abnormal protein accumulation.

Particular attention was given to the possibility of feedback relationships. For example, mitochondrial dysfunction can increase oxidative stress and lipid peroxidation, while oxidized lipid products can further damage membranes and cellular organelles. Likewise, lysosomal dysfunction can interfere with lipid degradation and recycling, leading to lipid accumulation that may subsequently influence inflammatory responses. Microglial lipid dysregulation has also been associated with neuroinflammatory processes, making neuron–glia interactions an important component of the analysis.

Comparative and Integrative Analysis

A comparative analysis was conducted to identify common and disease-specific molecular patterns. Genes and pathways appearing in multiple disorders were examined as potential points of convergence, while disease-specific findings were retained to avoid oversimplifying distinct pathological mechanisms. The analysis considered whether the same lipid pathway could produce different biological consequences depending on cell type, disease stage, genetic background, or environmental context.

The evidence was synthesized across three principal levels: **genetic regulation**, **lipid metabolic alteration**, and **neurodegenerative consequence**. This framework allowed the study to move from the identification of a genetic factor toward interpretation of its potential biochemical and cellular significance.

Analytical Level	Main Question	Examples of Variables Examined
Genetic	Which genetic factors influence lipid metabolism?	Variants, mutations, gene expression and regulatory changes
Metabolic	Which lipid pathways are altered?	Cholesterol, fatty acids, sphingolipids and phospholipids
Cellular	What cellular processes are affected?	Mitochondria, lysosomes, membranes, lipid droplets
Molecular	Which pathological mechanisms are involved?	Oxidative stress, inflammation, protein aggregation
Neurological	What are the potential consequences?	Synaptic dysfunction and neuronal degeneration
Clinical	How may molecular changes relate to disease?	Risk, progression, biomarkers and therapeutic targets

Bioinformatic and Functional Interpretation

Where appropriate, the identified genes and molecular pathways were interpreted using functional annotation and pathway-based approaches. Gene-function information was organized according to biological processes, molecular functions, cellular components, and metabolic pathways. Particular emphasis was placed on pathways connecting lipid metabolism with neuronal signaling, inflammation, mitochondrial activity, lysosomal degradation, autophagy, and membrane trafficking.

The methodological framework also considered evidence generated through genome-wide association studies, transcriptomic studies, lipidomic investigations, and other high-throughput approaches. Such integration is important because genetic association alone does not necessarily establish biological causation. A reported genetic variant was therefore interpreted together with available evidence concerning gene expression, protein function, lipid alteration, cellular phenotype, and disease association. Recent research using genetic approaches, including Mendelian-randomization analyses, has further explored potential relationships between lipid traits and neurodegenerative disease risk.

Reliability and Evidence Assessment

To improve the reliability of the synthesis, greater interpretive weight was given to studies with clearly described populations or experimental systems, appropriate analytical methods, reproducible molecular measurements, established genetic associations, and biologically supported conclusions. Findings reported by independent research groups were compared where possible. When studies produced conflicting observations, the differences were considered in relation to sample characteristics, disease stage, tissue type, lipid measurement techniques, genetic background, and analytical methodology rather than forcing a uniform conclusion.

The evidence was also categorized according to its strength and biological level.

Evidence Category	Interpretation
Genetic association	Indicates a relationship between a genetic factor and disease or lipid phenotype
Functional genetic evidence	Provides evidence that a genetic alteration affects molecular function
Biochemical evidence	Demonstrates changes in lipid concentration, composition or metabolism
Cellular evidence	Demonstrates effects on neurons, glia or organelles
Mechanistic evidence	Connects lipid alteration with a defined pathological process
Translational evidence	Indicates potential relevance to biomarkers or therapeutic intervention

Methodological Integration

Finally, all extracted evidence was integrated into a conceptual framework linking **genetic regulation** → **lipid metabolic alteration** → **cellular dysfunction** → **neurodegenerative pathology**. This framework provided the basis for interpreting how molecular changes in lipid-associated genes may influence neuronal vulnerability. The methodology recognizes that lipid metabolism is a dynamic network rather than an isolated biochemical pathway and that its effects depend on interactions among neurons, glial cells, organelles, inflammatory pathways, and systemic metabolic conditions. Recent literature similarly emphasizes that lipid metabolism, remodeling, and intercellular transfer are integral to CNS homeostasis and that their disruption may contribute to neurodegenerative disease development. Through this integrative methodology, the study sought to identify recurring molecular patterns, disease-specific mechanisms, genetic contributors, and potential biological targets that can improve understanding of the relationship between lipid metabolism and neurodegeneration.

RESULTS AND DISCUSSIONS

The integrated analysis demonstrated that lipid metabolism is closely connected with the molecular and cellular mechanisms underlying neurodegenerative diseases. The evidence reviewed indicates that neuronal lipid

homeostasis depends on coordinated regulation of lipid synthesis, transport, remodeling, oxidation, storage, and degradation rather than on the activity of an individual metabolic pathway. Cholesterol, phospholipids, sphingolipids, fatty acids, and related lipid mediators were repeatedly identified as important components of neuronal membranes, myelin, synaptic structures, intracellular organelles, and cell-signaling systems. Recent research has emphasized that lipid remodeling and intercellular lipid transfer between neurons and glial cells are essential for maintaining central nervous system homeostasis, while disruption of these processes can contribute to neurodegenerative pathology. The findings of the present synthesis therefore support a network-based interpretation in which genetic alterations affecting lipid metabolism can influence several downstream processes simultaneously, including membrane instability, mitochondrial dysfunction, oxidative stress, lysosomal impairment, neuroinflammation, and abnormal protein accumulation.

Relationship Between Lipid Classes and Neurodegenerative Mechanisms

The analysis identified several major lipid classes with recurring associations across neurodegenerative disorders. Cholesterol was particularly important because of its role in membrane organization, synaptic signaling, myelin maintenance, and protein–membrane interactions. Disturbances in cholesterol distribution may alter membrane properties and influence the organization of proteins involved in neuronal signaling. Evidence also indicates that cholesterol imbalance can affect amyloid-beta and α -synuclein-related processes, providing a molecular connection between lipid homeostasis and pathological protein aggregation.

Sphingolipids and glycerophospholipids similarly demonstrated substantial relevance. Changes in these lipid classes can influence membrane curvature, vesicular trafficking, cellular signaling, and inflammatory responses. Fatty-acid metabolism emerged as another important area, particularly in Parkinson’s disease, where recent research has proposed a strong relationship between fatty-acid dyshomeostasis and α -synuclein-associated pathology. The findings suggest that changes in individual lipid species should not be interpreted independently because alterations in one lipid pathway can affect substrate availability, signaling, membrane composition, and metabolic flux in interconnected pathways.

Lipid Category	Major Biological Functions	Potential Neurodegenerative Consequences
Cholesterol	Membrane structure, synaptic function, myelin maintenance	Altered membrane organization, impaired signaling and protein aggregation
Phospholipids	Membrane architecture and cellular signaling	Membrane instability, oxidative damage and impaired trafficking
Sphingolipids	Signaling, membrane organization and myelin biology	Inflammation, lysosomal dysfunction and altered neuronal signaling
Fatty acids	Energy metabolism, membrane composition and signaling	Oxidative stress, mitochondrial dysfunction and protein–lipid interactions
Glycerolipids	Energy storage and membrane-related metabolism	Lipid accumulation and cellular metabolic stress
Oxidized lipids	Signaling and products of lipid oxidation	Membrane damage and inflammatory responses

Genetic Regulation of Lipid Homeostasis

The molecular-genetic analysis demonstrated that lipid-related genes provide an important connection between inherited susceptibility and altered neuronal lipid biology. Genes involved in lipid transport, lysosomal function, lipid sensing, membrane remodeling, and intracellular trafficking were repeatedly associated with neurodegenerative mechanisms. In Alzheimer’s disease, genetic evidence has particularly strengthened the importance of pathways involving lipid transport and microglial function. Recent research has identified rare variants affecting genes such as **ABCA7**, **SORL1**, and **TREM2** as important contributors to understanding Alzheimer disease susceptibility and pathogenesis. These findings are significant because they indicate that genetic effects can influence not only lipid concentrations but also how neural cells detect, transport, recycle, and respond to lipid molecules.

The analysis also highlighted the importance of **APOE**, which plays a central role in lipid transport within the brain. Variability in APOE-related lipid handling has been associated with differences in susceptibility to neurological disease, although genetic associations should not be interpreted as deterministic predictions of disease. The findings indicate that lipid-related genetic factors operate within broader molecular networks involving immune responses, protein clearance, membrane biology, and cellular stress. Consequently, genetic predisposition may modify the biological environment in which neurodegenerative processes develop rather than independently determining disease progression.

Comparative Disease-Level Findings

Comparison across major neurodegenerative diseases demonstrated both shared and disease-specific relationships between lipid metabolism and pathological processes. Alzheimer’s disease showed a particularly strong association with cholesterol transport, lipid-mediated membrane regulation, sphingolipid metabolism, microglial activity, and mechanisms involved in amyloid and tau biology. Parkinson’s disease showed prominent relationships involving fatty acids, sphingolipids, lysosomal metabolism, mitochondrial dysfunction, and α -synuclein. These findings correspond with recent evidence that abnormal cholesterol, sphingolipid, and

glycerophospholipid metabolism can influence Alzheimer's and Parkinson's disease mechanisms through several interconnected pathways.

Neurodegenerative Disease	Prominent Lipid-Related Processes	Major Molecular/Cellular Associations
Alzheimer's disease	Cholesterol transport, phospholipid and sphingolipid metabolism	Amyloid processing, tau pathology, microglial function and synaptic dysfunction
Parkinson's disease	Fatty-acid, sphingolipid and lysosomal lipid metabolism	α -synuclein accumulation, mitochondrial dysfunction and oxidative stress
ALS	Membrane lipid and energy-associated metabolism	Neuronal stress, mitochondrial abnormalities and inflammation
Huntington's disease	Membrane and energy-related lipid pathways	Neuronal vulnerability and altered cellular metabolism
Frontotemporal dementia	Lysosomal and lipid-handling pathways	Protein clearance, microglial dysfunction and neuronal degeneration

The comparison suggests that lipid dysregulation may represent a common biological feature across different neurodegenerative diseases while the precise lipid pathways and cellular consequences vary according to disease-specific molecular environments. This distinction is important because a therapeutic approach designed for one lipid pathway may not necessarily produce equivalent effects in another neurodegenerative disorder.

Lipid Metabolism, Mitochondrial Dysfunction and Oxidative Stress

A major finding from the integrated analysis was the close relationship between lipid imbalance and mitochondrial dysfunction. Neurons have substantial energy requirements, and mitochondrial integrity is therefore essential for maintaining synaptic activity and cellular survival. Altered fatty-acid metabolism, excessive lipid accumulation, or lipid oxidation can interfere with mitochondrial function and increase cellular oxidative stress. At the same time, impaired mitochondrial activity can increase reactive oxygen species, promoting lipid peroxidation and generating secondary lipid products capable of damaging cellular membranes. This relationship indicates the possibility of a self-reinforcing metabolic cycle. An initial disturbance in lipid metabolism may increase oxidative stress, while oxidative damage can further alter lipid composition and mitochondrial membranes. The resulting dysfunction may affect energy production, intracellular trafficking, and neuronal signaling. Such interactions help explain why lipid abnormalities may persist and become progressively more damaging during neurodegenerative disease rather than remaining isolated biochemical abnormalities.

Lysosomal Function, Lipid Storage and Neuroinflammation

The results further identified lysosomal function as an important intersection between lipid metabolism and neurodegeneration. Lysosomes participate in the degradation and recycling of cellular materials, including several lipid-containing structures. When lysosomal processes are impaired, lipid substrates and damaged cellular components may accumulate, disrupting intracellular homeostasis. Genetic evidence from neurodegenerative diseases increasingly emphasizes the importance of cellular clearance pathways, particularly in relation to disease-associated protein accumulation.

The relationship between lipid metabolism and neuroinflammation was particularly evident in microglial biology. Microglia respond to changes in their surrounding environment by altering metabolic pathways, and lipid accumulation can accompany changes in inflammatory state. Recent evidence identifies dysregulated microglial lipid metabolism as an important component of neuroinflammation and highlights progranulin, encoded by **GRN**, as a regulator of lipid-droplet formation and lysosomal lipid metabolism. These findings provide an important link between genetic regulation, lipid storage, lysosomal function, and inflammatory responses.

Interaction Between Lipids and Protein Aggregation

Another significant finding was the interaction between lipid metabolism and pathological protein accumulation. Neurodegenerative diseases are commonly characterized by abnormal accumulation or altered processing of proteins such as amyloid-beta, tau, and α -synuclein. The present synthesis indicates that membrane lipids can influence the localization, conformation, processing, and aggregation behavior of such proteins. Conversely, pathological proteins can interfere with lipid trafficking, organelle function, and membrane stability.

In Alzheimer's disease, abnormal lipid metabolism has been associated with processes involving amyloid-beta deposition and tau pathology, while Parkinson's disease demonstrates a particularly notable interaction between lipid biology and α -synuclein. This bidirectional relationship suggests that lipid imbalance may both contribute to pathological protein accumulation and be worsened by it. Consequently, targeting only protein aggregation without considering the lipid environment may overlook an important component of disease biology.

Integrated Molecular Relationship

The combined findings support a model in which genetic variation influences lipid-related enzymes, transport proteins, receptors, regulatory molecules, or lysosomal components, resulting in changes in lipid synthesis, distribution, remodeling, or degradation. These metabolic changes can subsequently affect organelles and cellular signaling systems, eventually increasing neuronal vulnerability.

Genetic/Metabolic Alteration	Immediate Effect	Downstream Cellular Effect	Possible Consequence Disease
Altered lipid transport	Abnormal lipid distribution	Membrane and signaling changes	Synaptic dysfunction
Impaired sphingolipid metabolism	Altered membrane lipid composition	Signaling and inflammatory changes	Neuronal vulnerability
Abnormal fatty-acid metabolism	Altered lipid oxidation	Mitochondrial stress	Oxidative damage
Lysosomal lipid-processing defect	Lipid accumulation	Impaired cellular clearance	Protein and organelle accumulation
Altered lipid-droplet regulation	Excess or insufficient lipid storage	Cellular metabolic stress	Neuroinflammatory responses
Cholesterol imbalance	Modified membrane organization	Altered protein–membrane interactions	Synaptic and neuronal dysfunction

Overall, the results indicate that the relationship between molecular genetics and neurodegeneration is best understood through interconnected biological pathways rather than isolated gene–disease associations. Genetic alterations can modify lipid homeostasis, but their pathological effects depend on cellular context, aging, environmental influences, metabolic state, and interactions with other genetic pathways. Recent work using genetic, lipidomic, and transcriptomic approaches supports increasingly detailed investigation of these relationships, including cell-type-specific changes in lipid metabolism.

The findings also have implications for biomarker development and therapeutic research. Lipidomic profiling combined with genetic information may help identify molecular subgroups characterized by distinct metabolic abnormalities. However, the presence of a lipid alteration does not automatically establish that it is a primary cause of neurodegeneration. Some changes may arise as compensatory responses or consequences of neuronal injury. Future investigations should therefore integrate longitudinal genetic, lipidomic, transcriptomic, and cellular data to distinguish causal mechanisms from secondary metabolic effects. The overall evidence supports lipid metabolism as an important molecular interface connecting genetic susceptibility with membrane dysfunction, mitochondrial impairment, lysosomal abnormalities, neuroinflammation, protein pathology, and progressive neuronal degeneration.

CONCLUSION

The present study establishes that the molecular regulation of lipid metabolism represents an important component of the biological processes associated with neurodegenerative diseases. Lipids are not merely structural constituents of neuronal membranes but actively participate in membrane organization, synaptic communication, energy metabolism, intracellular trafficking, myelin maintenance, and cellular signaling. The integrated analysis indicates that genetic alterations affecting lipid transport, synthesis, remodeling, storage, and degradation can disturb these functions and contribute to neuronal vulnerability. Particular relationships were identified between lipid dysregulation and mitochondrial impairment, oxidative stress, lysosomal dysfunction, neuroinflammation, abnormal protein processing, and impaired cellular clearance. The involvement of genes and pathways associated with cholesterol, phospholipids, sphingolipids, fatty acids, and lipid droplets further demonstrates that neurodegeneration is influenced by interconnected metabolic networks rather than by isolated molecular abnormalities. Recent evidence also demonstrates that specific genetic alterations can produce measurable changes in neuronal or glial lipid metabolism, strengthening the biological connection between genetic susceptibility and lipid homeostasis.

Overall, the findings support an integrated genetic–lipid–cellular framework for understanding neurodegenerative disease development and progression. The interaction between inherited or acquired molecular alterations and age-related metabolic stress may determine how effectively neural cells maintain lipid balance and respond to pathological conditions. Advances in genomics, lipidomics, transcriptomics, single-cell analysis, and multi-omics approaches provide opportunities to identify disease-associated lipid signatures, clarify cell-specific mechanisms, and investigate potential biomarkers and therapeutic targets. Emerging evidence involving genes such as **ABCA7**, **APOE**, **GRN**, and other lipid-related regulators illustrates the potential value of connecting genetic information with functional lipid changes rather than examining genetic associations independently. Nevertheless, further longitudinal and mechanistic studies are required to distinguish primary causal lipid disturbances from metabolic changes that occur as consequences of neuronal injury. A deeper understanding of these relationships may ultimately support more precise approaches to diagnosis, disease stratification, and therapeutic development while advancing knowledge of how molecular lipid regulation contributes to the complex biology of neurodegenerative diseases.

REFERENCES

1. Cooper, Oliver, Penny Hallett, and Ole Isacson. "Upstream Lipid and Metabolic Systems Are Potential Causes of Alzheimer's Disease, Parkinson's Disease and Dementias." *FEBS Journal*, vol. 291, no. 4, 2024, pp. 632–645.
2. Yang, Linda G., Zachary M. March, Roxan A. Stephenson, and Priyanka S. Narayan. "Apolipoprotein E in Lipid Metabolism and Neurodegenerative Disease." *Trends in Endocrinology & Metabolism*, vol. 34, no. 8, 2023, pp. 430–445.
3. Tong, Bin, et al. "Targeting Dysregulated Lipid Metabolism for the Treatment of Alzheimer's Disease and Parkinson's Disease: Current Advancements and Future Prospects." *Neurobiology of Disease*, vol. 196, 2024, article 106505.
4. Long, Junpeng, et al. "Targeting Lipid Metabolism in Neurodegenerative Diseases: From Experimental to Clinical." *Metabolism*, vol. 175, 2026, article 156436.
5. Kang, Yubing, et al. "Role of Lipid Droplets in Neurodegenerative Diseases: From Pathogenesis to Therapeutics." *Neuroscience & Biobehavioral Reviews*, vol. 165, 2024, article 105867.
6. Mallick, Keya, Shuchismita Paul, Sayani Banerjee, and Sugato Banerjee. "Lipid Droplets and Neurodegeneration." *Neuroscience*, vol. 548, 2024, pp. 1–15.
7. Shin, Kyung Chul, Houda Yasmine Ali Moussa, and Yongsoo Park. "Cholesterol Imbalance and Neurotransmission Defects in Neurodegeneration." *Experimental & Molecular Medicine*, vol. 56, 2024, pp. 1685–1690.
8. Kamano, Shumpei, Daisaku Ozawa, Kensuke Ikenaka, and Yoshitaka Nagai. "Role of Lipids in the Pathogenesis of Parkinson's Disease." *International Journal of Molecular Sciences*, vol. 25, no. 16, 2024, article 8935.
9. Qin, Bingqing, et al. "Lipid Metabolism in Health and Disease: Mechanistic and Therapeutic Insights for Parkinson's Disease." *Chinese Medical Journal*, vol. 138, no. 12, 2025, pp. 1411–1423.
10. Yilmaz, Ali, et al. "Lipid Profiling of Parkinson's Disease Brain Highlights Disruption in Lysophosphatidylcholines, and Triacylglycerol Metabolism." *npj Parkinson's Disease*, vol. 11, 2025, article 159.
11. Wang, Zeyu, Zixiao Yin, Guangyong Sun, Dong Zhang, and Jianguo Zhang. "Genetic Evidence for the Liver-Brain Axis: Lipid Metabolism and Neurodegenerative Disease Risk." *Lipids in Health and Disease*, vol. 24, 2025, article 41.
12. Windham, Ian A., and Sarah Cohen. "The Cell Biology of APOE in the Brain." *Trends in Cell Biology*, vol. 34, 2024, pp. 198–211.
13. Martens, Yuka A., et al. "ApoE Cascade Hypothesis in the Pathogenesis of Alzheimer's Disease and Related Dementias." *Neuron*, vol. 110, no. 8, 2022, pp. 1304–1317.
14. Liu, Chia-Chen, et al. "Peripheral ApoE4 Enhances Alzheimer's Pathology and Impairs Cognition by Compromising Cerebrovascular Function." *Nature Neuroscience*, vol. 25, no. 8, 2022, pp. 1020–1033.
15. Lou, Tianwen, Borui Tao, and Ming Chen. "Relationship of Apolipoprotein E with Alzheimer's Disease and Other Neurological Disorders: An Updated Review." *Neuroscience*, vol. 514, 2023, pp. 123–140.
16. Sun, Yu-Ying, Zhun Wang, and Han-Chang Huang. "Roles of ApoE4 on the Pathogenesis in Alzheimer's Disease and the Potential Therapeutic Approaches." *Cellular and Molecular Neurobiology*, vol. 43, no. 7, 2023, pp. 3115–3136.
17. Kurano, Makoto, et al. "Isoform-Dependent Effects of Apolipoprotein E on Sphingolipid Metabolism in Neural Cells." *Journal of Alzheimer's Disease*, vol. 85, no. 4, 2022, pp. 1529–1544.
18. Yang, Danying, et al. "Lipid Metabolism and Storage in Neuroglia: Role in Brain Development and Neurodegenerative Diseases." *Cell & Bioscience*, vol. 12, 2022, article 106.
19. Li, Hangzhen, et al. "The Potential Role of Glucose Metabolism, Lipid Metabolism, and Amino Acid Metabolism in the Treatment of Parkinson's Disease." *CNS Neuroscience & Therapeutics*, vol. 30, no. 2, 2024, article e14411.
20. Maltseva, Sofiya, Daniel Kerr, Miah Turke, Erin J. Adams, and Ka Yee C. Lee. "Parkinson's Disease-Associated Mutations in α -Synuclein Alters Its Lipid-Bound State." *Biophysical Journal*, vol. 123, no. 12, 2024, pp. 1610–1619.
21. Zhu, Hongling, et al. "Reactivation of mTOR Signaling Slows Neurodegeneration in a Lysosomal Sphingolipid Storage Disease." *Neurobiology of Disease*, vol. 204, 2025, article 106760.
22. Islam, M. Nurul, et al. "Sphingolipid Signaling and Metabolism in Neuronal and Glial Cells: Implications for Cerebrovascular and Neurodegenerative Disorders." *Aging*, 2025.
23. "The Physiological and Pathological Effects of Sphingolipid Metabolism and Signaling in the Central Nervous System." *Brain Pathology*, 2025.
24. Tsap, Mariana I., and Halyna R. Shcherbata. "The Balancing Act between Lipid Droplets and Lysosomes for Membrane Functionality in Age-Related Neurodegeneration and Inflammation." *Progress in Lipid Research*, 2025, article 101341.
25. "Lipid Metabolism and Neurodegeneration: Mechanistic Insights and Therapeutic Targets." PubMed-indexed review, 2026.

26. "The Neurolipid Atlas: A Lipidomics Resource for Neurodegenerative Diseases." *Nature Metabolism*, 2025.
27. "APOE4/4 Is Linked to Damaging Lipid Droplets in Alzheimer's Disease Microglia." *Nature*, vol. 628, 2024, pp. 154–161.
28. "TDP-43 Dysfunction Leads to Bioenergetic Failure and Lipid Metabolic Rewiring in Human Cells." *Redox Biology*, vol. 74, 2024, article 103301.
29. "Multiple Pathways of Lipid Dysregulation in Amyotrophic Lateral Sclerosis." PubMed-indexed study, 2023.
30. "Glycerophospholipids in ALS: Insights into Disease Mechanisms and Clinical Implication." PubMed-indexed review, 2025.