

STRUCTURE-PROPERTY ANALYSIS OF D- π -A NAPHTHALENE CHROMOPHORES FOR OPTOELECTRONIC APPLICATIONS

Gorak Tharangini¹, Dr. Rajamouli Boddula², Vijay Singh³, Shashanka Rajendrachari⁴, Nadeem Siddiqui⁵

¹SR University, Ananthasagar, Hasanparthy, Warangal - 506371, Telangana, India, Email: tharangini2001@gmail.com

²SR University, Ananthasagar, Hasanparthy, Warangal - 506371, Telangana, India

⁴Global Academy of Technology, Bangalore, Karnataka 560098, India

⁵Koneru Lakshmaiah Education Foundation (Deemed to be University), Vaddeswram, Green Fields, AP 522502, India,


ABSTRACT

Donor- π -acceptor chromophores are an important class of organic functional materials because their optical and electronic properties can be tuned through molecular design. In these systems, an electron-donating group and an electron-accepting group are connected through a conjugated π -bridge, enabling intramolecular charge transfer after excitation. Naphthalene is a suitable structural platform for such chromophores because of its rigid aromatic framework, chemical stability, extended conjugation and multiple substitution sites. The present article examines the relationship between molecular structure and optoelectronic behaviour in D- π -A naphthalene chromophores. The analysis focuses on donor and acceptor strength, π -conjugation, substitution pattern, molecular planarity, solvent polarity, aggregation, frontier molecular orbitals and nonlinear optical response. The study indicates that efficient optoelectronic performance depends on a balanced relationship among charge transfer, molecular rigidity, orbital overlap, fluorescence efficiency, thermal stability and solid-state packing. Extended conjugation generally shifts absorption towards longer wavelengths and narrows the optical energy gap, but excessive flexibility and aggregation may increase non-radiative losses. The article proposes an integrated experimental and computational methodology for designing and evaluating naphthalene chromophores for organic light-emitting devices, fluorescence sensing, photovoltaics, nonlinear optics and molecular electronics.

KEYWORDS: D- π -A chromophores; naphthalene; charge transfer; fluorescence; optoelectronics; nonlinear optics; density functional theory.

1. INTRODUCTION

Organic optoelectronic materials have attracted increasing scientific attention because they are lightweight, structurally adaptable and often suitable for solution-based processing. Their absorption, emission, energy levels and charge-transfer behaviour can be adjusted through chemical modification. This molecular flexibility makes them useful in organic light-emitting diodes, solar cells, sensors, optical switches, field-effect transistors and photonic devices.



Among the different molecular architectures used in organic electronics, donor- π -acceptor systems are especially significant. These molecules contain an electron-rich donor and an electron-deficient acceptor linked through a conjugated π -system. On optical excitation, electron density moves from the donor region towards the acceptor region, producing an intramolecular charge-transfer state. This process influences absorption wavelength, fluorescence colour, Stokes shift, dipole moment, oxidation and reduction potentials, excited-state lifetime and nonlinear optical response.

The π -bridge controls electronic communication between the donor and acceptor. Its length, rigidity, planarity and chemical composition determine the extent of orbital overlap and charge delocalisation. Greater conjugation often lowers the optical energy gap and shifts absorption towards longer wavelengths. However, a longer bridge may also introduce rotational freedom, molecular twisting and non-radiative relaxation.

Naphthalene is a useful framework for constructing D- π -A chromophores because its two fused benzene rings provide rigidity, conjugation and several functionalisation positions. Substitution at different positions can alter molecular symmetry, direction of electronic communication, steric interaction and solid-state packing. Therefore, the naphthalene core is not merely a structural support; it actively regulates photophysical and electronic behaviour.

The principal challenge in chromophore design is that improvement in one property may reduce another. Strong charge transfer can produce long-wavelength absorption and high molecular polarisation, but it may also weaken fluorescence. High molecular planarity improves conjugation, yet close π - π stacking can cause aggregation-related quenching. Rational molecular design must therefore balance electronic strength, geometric stability and intermolecular organisation.

2. Conceptual Basis of D- π -A Naphthalene Chromophores

2.1 Electron-donating groups

The donor unit supplies electron density to the conjugated system. Common donors include methoxy, dimethylamino, diphenylamino, triphenylamine, carbazole and phenothiazine groups. Stronger donor groups generally raise the highest occupied molecular orbital energy and strengthen intramolecular charge transfer.

Dimethylamino groups provide strong electron donation and may produce red-shifted absorption. However, they can also increase sensitivity to protonation and solvent polarity. Carbazole and triphenylamine units offer good electron-donating ability, structural rigidity and thermal stability, making them suitable for optoelectronic applications.

2.2 π -conjugated bridge

The π -spacer connects the donor and acceptor and determines the efficiency of electronic communication. Phenylene, vinylene, ethynylene and thiophene are commonly used bridges. Longer conjugation generally increases delocalisation and reduces the energy gap.

Ethynylene spacers provide a relatively linear pathway and reduce steric hindrance. Vinylene units strengthen conjugation but may undergo photoisomerisation. Thiophene spacers increase polarizability and facilitate visible-light absorption, although they may influence oxidation stability.

The effectiveness of a bridge depends not only on its length but also on its geometry. A short and planar bridge may support stronger charge transfer than a long but twisted pathway.

2.3 Electron-accepting groups


The acceptor receives electron density in the excited state. Common acceptors include cyano, nitro, aldehyde, ketone, benzothiazole, dicyanovinyl and imide groups. Strong acceptors lower the lowest unoccupied molecular orbital and promote charge separation.

Cyano and dicyanovinyl units often produce strong charge-transfer absorption and narrow energy gaps. Naphthalene-diimide derivatives are also valuable because of their electron-deficient nature, reduction stability and ability to support intermolecular organisation.

Excessively strong acceptors may, however, increase non-radiative relaxation and reduce fluorescence. The acceptor must therefore be selected according to the intended application.

2.4 Role of the naphthalene core

Naphthalene offers a rigid and conjugated aromatic surface. Its substitution position strongly influences electronic and geometric properties. Linear substitution patterns such as 2,6- or 2,7-substitution may support extended conjugation and charge transport. In contrast, 1,8-substitution may create steric crowding and through-space electronic interaction.



Positional isomers may therefore display different absorption, fluorescence, orbital distribution and molecular packing, even when they contain identical donor and acceptor groups.

3. Review of Related Literature

Table 1. Summary of Literature on D- π -A Naphthalene Chromophores

Authors (Year)	Molecule Type / System	Key Findings	Method Used	Major Observation	Reference
Bureš (2014)	Push-pull chromophores (D- π -A)	Optical and electronic properties depend on donor-acceptor strength, π -bridge length and molecular shape.	Photophysical, Electrochemical, Computational	Stronger donor and acceptor reduce HOMO-LUMO gap and enhance charge transfer.	[RSC Adv., 4, 58826–58851]
Dokládlová et al. (2014)	D- π -A chromophores with varied π -bridge	Molecular first hyperpolarizability is influenced by bridge length, planarity and polarizability.	Computational (DFT)	Longer bridge does not always give higher NLO response due to electronic decoupling.	[Org. Biomol. Chem., 12, 7412–7425]
Pannipara et al. (2017)	Naphthalene-based AIE fluorophores	Restriction of intramolecular rotation in aggregated state enhances solid-state emission.	Synthesis, Photophysical	Aggregation-induced emission leads to strong solid-state fluorescence.	[Opt. Mater., 72, 13–21]
Kumari et al. (2017)	D- π -A compounds in various solvents	Increase in solvent polarity causes red shift in absorption and fluorescence due to ICT stabilization.	UV-Vis, Fluorescence Solvatochromism	Excited state has larger dipole moment than ground state.	[RSC Adv., 7, 16520–16535]
Nath et al. (2022)	Donor-acceptor system	Directional charge transfer from donor to acceptor confirmed by spectra and DFT calculations.	Steady-state, Time-resolved, DFT	HOMO localized on donor, LUMO on acceptor; ICT increases in polar solvents.	[J. Mol. Liq., 347, 117939]
Babejová et al. (2024)	Series of D- π -A chromophores	Systematic variation in structure affects optical, electrochemical and theoretical properties.	Spectroscopy, Electrochemistry, DFT	Different donors, acceptors and bridges lead to tunable properties.	[RSC Adv., 14, 22567–22582]

Bureš (2014) critically examined push-pull chromophores and established that their optical and electronic properties are governed by the combined effects of the electron donor, π -conjugated bridge and electron acceptor. The study demonstrated that variations in molecular shape, bridge length and donor-acceptor strength can significantly modify frontier molecular orbital energies, absorption behaviour and intramolecular charge transfer. Dokládlová et al. (2014) investigated structurally modified D- π -A chromophores and reported that molecular nonlinearity depends not only on donor and acceptor strength but also on the length, planarity and polarizability of the π -linker. Their findings showed that a stronger acceptor does not always produce a higher nonlinear optical response when the donor and acceptor become electronically separated by an excessively long bridge.

Pannipara et al. (2017) studied naphthalene-based aggregation-induced emissive materials and demonstrated that restriction of intramolecular rotation in the aggregated state can suppress non-radiative energy loss. This structural rigidification can enhance solid-state fluorescence, making suitably engineered naphthalene derivatives valuable for sensors, imaging systems and light-emitting materials.

Kumari et al. (2017) observed that increasing solvent polarity caused bathochromic shifts in the absorption and fluorescence spectra of donor-acceptor compounds. The researchers attributed this behaviour to stabilisation of the polar intramolecular charge-transfer excited state. Their results also indicated that the excited state possesses a greater dipole moment than the ground state.

Nath et al. (2022) combined steady-state spectroscopy, time-resolved fluorescence and density functional theory to investigate excited-state charge transfer in a donor-acceptor molecule. A pronounced red shift in polar solvents and the calculated increase in excited-state dipole moment confirmed efficient electron-density transfer from the donor to the acceptor. The HOMO was primarily associated with the donor region, whereas the LUMO extended towards the acceptor moiety.

Babejová et al. (2024) examined a systematically designed series of nine D- π -A chromophores using optical, electrochemical and computational methods. Their study confirmed that changes in the π -linker and terminal donor-acceptor groups produced measurable differences in absorption, fluorescence, oxidation-reduction behaviour and calculated electronic structure.

Collectively, these studies confirm that the performance of D- π -A chromophores depends on the balanced interaction of donor strength, acceptor ability, bridge composition, molecular rigidity and surrounding medium. However, comparatively fewer studies have systematically examined how different substitution positions on the naphthalene core influence these properties. This limitation creates a clear need for controlled structure-property analysis using related molecular series rather than isolated chromophores.

4. Research Gap

The existing literature reveals several limitations.

First, many studies investigate only one or two compounds, making it difficult to identify the structural factor responsible for a particular optical property.

Second, absorption and fluorescence are often examined without integrating electrochemical, thermal, computational and solid-state results.

Third, positional substitution on the naphthalene core has received less systematic attention than donor and acceptor modification.

Fourth, many computational studies report high hyperpolarizability or a small HOMO-LUMO gap without experimentally confirming synthesis feasibility, stability or bulk optical behaviour.

Fifth, solution-state fluorescence is sometimes used to predict device suitability without examining aggregation, film morphology and thermal resistance.

The present study addresses these gaps through an integrated structure-property framework.

5. Aim and Objectives

Aim

To analyse the influence of molecular structure on the optical, electronic and nonlinear optical properties of D- π -A naphthalene chromophores for potential optoelectronic applications.

Objectives

- To study the effect of donor, π -spacer and acceptor units on the electronic structure of naphthalene chromophores.
- To examine the influence of conjugation length, molecular planarity and substitution position on absorption and fluorescence.
- To correlate experimental photophysical properties with calculated frontier molecular orbitals and nonlinear optical parameters.
- To propose rational molecular-design principles for application-specific naphthalene chromophores.

6. RESEARCH METHODOLOGY

The proposed investigation may combine experimental, analytical and computational methods.

6.1 Molecular design

A comparative series may be prepared using:

- methoxy, dimethylamino, carbazole and triphenylamine donors;
- phenylene, vinylene, ethynylene and thiophene spacers;
- cyano, nitro, aldehyde, benzothiazole and dicyanovinyl acceptors;
- 1-, 2-, 1,8-, 2,6- and 2,7-substituted naphthalene frameworks.

Only one major structural component should be changed at a time to ensure reliable comparison.

6.2 Synthesis and purification

The compounds may be synthesised through Suzuki–Miyaura, Heck or Sonogashira coupling, Knoevenagel condensation, nucleophilic substitution or Schiff-base formation. Reaction progress may be monitored using thin-layer chromatography. Purification may be carried out through recrystallisation, column chromatography or preparative chromatography.

6.3 Structural characterisation

The synthesised compounds should be characterised using:

- Fourier-transform infrared spectroscopy;
- proton and carbon-13 nuclear magnetic resonance spectroscopy;
- high-resolution mass spectrometry;
- elemental analysis;
- single-crystal X-ray diffraction, where possible.

These techniques can confirm functional groups, molecular structure, purity, bond geometry and molecular packing.

6.4 Photophysical analysis

UV–visible and fluorescence spectra should be recorded in solvents of different polarity. The following parameters may be determined:

- maximum absorption wavelength;
- molar extinction coefficient;
- absorption onset;
- optical energy gap;
- fluorescence maximum;
- Stokes shift;
- fluorescence quantum yield;
- excited-state lifetime.

The optical energy gap may be estimated through:

- Optical energy gap (eV) = $1240 \div$ absorption onset wavelength (nm)
- Solvatochromic analysis may be used to determine how solvent polarity affects the charge-transfer excited state.

6.5 Electrochemical and thermal analysis

Cyclic voltammetry may be used to estimate oxidation potential, reduction potential and frontier-orbital energy levels. Thermogravimetric analysis and differential scanning calorimetry may be employed to examine decomposition temperature, melting behaviour and thermal stability.

6.6 Computational analysis

Density functional theory and time-dependent density functional theory may be used to calculate:

optimised geometry;

- HOMO and LUMO energies;
- molecular orbital distribution;
- HOMO–LUMO energy gap;
- dipole moment;
- molecular electrostatic potential;
- polarizability;
- first hyperpolarizability;
- oscillator strength;
- electron–hole distribution.

Range-separated functionals should preferably be used for charge-transfer excitations because conventional functionals may underestimate long-range electronic interactions.

6.7 Nonlinear optical evaluation

Selected chromophores may be examined through theoretical hyperpolarizability calculations and experimental techniques such as the Z-scan method, second-harmonic generation or hyper-Rayleigh scattering.

7. Structure–Property Relationships

7.1 Donor strength and optical behaviour

Stronger donors generally raise the HOMO level and reduce excitation energy. This may cause red-shifted absorption and stronger charge-transfer character. However, excessive donation may increase excited-state relaxation and reduce fluorescence efficiency.

7.2 Acceptor strength and charge separation

Stronger acceptors lower the LUMO level and facilitate electron transfer. Cyano and dicyanovinyl acceptors may produce narrower optical energy gaps than aldehyde or ester groups. Strong charge separation is useful for photovoltaic systems but may weaken emission.

7.3 Conjugation length

Longer conjugation increases electron delocalisation and usually shifts absorption towards longer wavelengths. Nevertheless, the bridge must remain planar and electronically connected. Rotational flexibility can reduce effective conjugation and increase internal conversion.

7.4 Molecular planarity

Planarity supports p-orbital overlap and charge transfer. It can improve absorption intensity and reduce the energy gap. However, highly planar molecules may aggregate through π – π stacking and display fluorescence quenching in the solid state.

7.5 Substitution position

Naphthalene substitution determines the direction of electronic communication. Linear substitution may improve conjugation, while peri substitution may introduce steric crowding and molecular twisting. Positional isomers should therefore be compared experimentally and computationally.

7.6 Solvent effects

Polar solvents stabilise charge-transfer excited states and may shift emission towards longer wavelengths. Large solvent-dependent shifts indicate a substantial difference between ground- and excited-state dipole moments. Excessive stabilisation may reduce quantum yield through non-radiative relaxation.

7.7 Aggregation behaviour

Some chromophores suffer from aggregation-caused quenching, while others exhibit aggregation-induced emission. Bulky substituents may prevent close π – π stacking, while rigid structures can restrict intramolecular motion. Solid-state performance must therefore be investigated separately from solution behaviour.

7.8 Frontier molecular orbitals

An effective D– π –A chromophore generally displays a HOMO localised mainly on the donor and a LUMO concentrated towards the acceptor. A smaller HOMO–LUMO gap predicts longer-wavelength absorption, but excessive orbital separation may reduce oscillator strength and fluorescence probability.

7.9 Nonlinear optical response

Molecular asymmetry, strong intramolecular charge transfer and low excitation energy can increase first hyperpolarizability. However, high theoretical hyperpolarizability does not always produce strong bulk nonlinear behaviour because crystal symmetry and molecular orientation also influence the final response.

8. Comparative Structure–Property Framework

Structural feature	Expected benefit	Possible limitation
Strong donor	Raises HOMO and enhances charge transfer	Oxidation and fluorescence loss
Strong acceptor	Lowers LUMO and improves charge separation	Non-radiative relaxation
Extended π -bridge	Narrows optical energy gap	Molecular flexibility
Rigid planar structure	Improves orbital overlap	π – π aggregation
Bulky substituent	Reduces solid-state quenching	Molecular twisting
Polar solvent	Stabilises charge-transfer state	Reduced quantum yield
Linear substitution	Supports extended conjugation	Difficult synthesis
Peri substitution	Enables through-space interaction	Steric congestion

9. Potential Applications

9.1 Organic light-emitting devices

Naphthalene chromophores with strong fluorescence, suitable energy levels and good thermal stability may be used as emitting or charge-transport materials.

9.2 Fluorescence sensors

Their sensitivity to polarity, protonation, ions and molecular environment makes D– π –A systems useful for chemical and biological sensing.

9.3 Organic photovoltaics

Broad absorption and efficient charge separation may enable their use as molecular donors, acceptors or sensitising materials.

9.4 Nonlinear optics

Chromophores with high molecular polarisation may be explored for optical switching, frequency conversion, modulation and optical limiting.

9.5 Molecular electronics

Directional charge-transfer behaviour makes these molecules suitable for molecular rectification, charge transport and switching applications.

10. Major Findings

The analytical study produces the following findings:

- Naphthalene provides a rigid and structurally adaptable framework for donor–acceptor chromophores.
- Strong donor–acceptor interaction generally narrows the energy gap and shifts absorption towards longer wavelengths.
- Effective conjugation depends more on molecular planarity and orbital overlap than on bridge length alone.
- Strong charge transfer may improve solvatochromism and nonlinear optical response but reduce fluorescence efficiency.
- Positional substitution significantly influences molecular geometry and electronic communication.
- Aggregation and solid-state packing are critical factors in practical optoelectronic performance.
- Computational modelling is valuable for molecular screening but must be supported by experimental validation.
- No single chromophore structure is ideal for every application; molecular design must be application-specific.

11. Scientific Significance

The importance of this study lies in its integrated examination of molecular structure, excited-state behaviour and practical optoelectronic requirements. It highlights that absorption wavelength, energy gap or calculated hyperpolarizability should not be treated as isolated indicators of performance.

The proposed framework considers donor and acceptor strength, π -bridge composition, substitution pattern, molecular rigidity, solvent response and aggregation together. This approach can reduce trial-and-error synthesis and support the selection of more promising chromophore structures.

12. Limitations and Future Scope

The present work is conceptual and does not report newly synthesised molecules or original spectral data. The proposed structural trends must therefore be experimentally tested.

Future research should focus on:

- controlled synthesis of positional naphthalene isomers;
- aggregation-induced-emission naphthalene systems;
- water-compatible chromophores for biosensing;
- thin-film and device-level evaluation;
- non-centrosymmetric materials for nonlinear optics;

- machine-learning-assisted molecular screening;
- long-term photochemical and thermal stability.

13. CONCLUSION

D- π -A naphthalene chromophores are promising organic materials for optoelectronic and photonic applications. Their behaviour is controlled by the combined influence of donor strength, acceptor character, bridge composition, molecular planarity, substitution position, solvent environment and intermolecular packing.

The naphthalene core provides rigidity, conjugation and multiple functionalisation pathways. Strong donor-acceptor interaction can narrow the energy gap and enhance charge transfer, but excessive polarity or structural relaxation may reduce fluorescence. Similarly, planar structures improve orbital overlap but can promote aggregation-related quenching.

Successful chromophore design therefore requires a balanced approach. The donor and acceptor should be electronically compatible, the π -bridge should be rigid but not excessively extended, and molecular packing should be controlled. Experimental spectroscopy, electrochemistry, thermal analysis and density functional theory should be used together to establish reliable structure-property relationships.

Such an integrated strategy can support the development of efficient naphthalene-based materials for organic light-emitting devices, fluorescence sensors, photovoltaic systems, nonlinear optical devices and molecular electronics.

REFERENCES

1. Abbotto, A., Beverina, L., Bradamante, S., Facchetti, A., Klein, C., Pagani, G. A., Redi-Abshiro, M., and Wortmann, R. (2003). Cooperative effects of structure and environment in intramolecular charge-transfer chromophores. *Chemistry—A European Journal*, 9, 1991–2007.
2. Alyar, H. (2013). Nonlinear optical properties of donor-acceptor derivatives of naphthalene and azanaphthalene. *Reviews on Advanced Materials Science*, 34, 79–87.
3. Barakat, A., et al. (2015). Structural and computational investigation of a naphthalene-based nonlinear optical material. *Chemistry Central Journal*, 9.
4. Bureš, F. (2014). Fundamental aspects of property tuning in push-pull molecules. *RSC Advances*, 4, 58826–58851.
5. Bureš, F., et al. (2016). Design of heteroaromatic push-pull chromophores. *The Chemical Record*.
6. Lai, C. H., Muhammad, S., Al-Sehemi, A. G., and Chaudhry, A. R. (2019). Nonlinear optical properties of naphthalene-diimide derivatives. *Journal of Molecular Graphics and Modelling*, 87, 68–75.
7. Nath, S., et al. (2022). Spectroscopic and theoretical investigation of excited-state charge transfer in a donor-acceptor system. *Journal of Molecular Liquids*.
8. Panahi, F., et al. (2020). A donor- π -acceptor halochromic distyrylnaphthalene fluorescent compound. *RSC Advances*, 10.
9. Pannipara, M., et al. (2017). Naphthalene-based aggregation-induced-emission-active material. *Optical Materials*.
10. Parthasarathy, V., Pandey, R., Das, P. K., Castet, F., and Blanchard-Desce, M. (2018). Linear and nonlinear optical properties of merocyanine dyes. *ChemPhysChem*, 19(2), 187–197.
11. Qiu, Y. Q., Li, Z., Ma, N. N., Sun, S. L., Zhang, M. Y., and Liu, P. J. (2013). Nonlinear optical properties of aromatic diimide molecules. *Journal of Molecular Graphics and Modelling*, 41, 79–88.
12. Shkir, M., Irfan, A., AlFaify, S., Patil, P. S., and Al-Sehemi, A. G. (2019). Linear and nonlinear optical properties of donor-acceptor chromophores. *Optik*, 199, 163354.
13. Yun, Y. J., et al. (2022). Positional effects on energy and charge transfer in donor-acceptor dyads. *Photochem.*