

GENOMIC CHARACTERIZATION OF ANTIBIOTIC-RESISTANT BACTERIA IN CLINICAL AND ENVIRONMENTAL SAMPLES

Dr. Nithya Jayan^{1*}, Dr. S. I. Saheed Askar², Dr. Ajab Singh Choudhary³, Dr. Venkateswarlu Yadavalli⁴, Dr. Biju Sidharthan⁵

¹Assistant Professor, Department of Microbiology, Sree Narayana College Alathur (affiliated to University of Calicut, Thenhipalam, Malappuram), Palakkad, Kerala, nithyajayan@sncollegealathur.ac.in, ORCID ID: 0009-0000-2462-2339

²Assistant Professor, Department of Clinical Microbiology, Government Medical College & Hospital, Ramanathapuram, T.N.Dr. M.G.R. Medical University, Guindy, Chennai, Tamil Nadu, India.saheedaskar@gmail.com, ORCID ID: 0009-0000-8488-4533

³Assistant Professor, Department of Medical Laboratory Technology, SOAHS, Noida International University, Sector 17 A Yamuna Expressway, Greater Noida, UP, ajab.singh@niu.edu.in, ORCID ID: <https://orcid.org/0000-0002-2338-9106>

⁴Associate Professor, Department of Biotechnology, Tara Govt. College (A), Sangareddy/Telangana, venkibiotech@gmail.com, ORCID ID: <https://orcid.org/0000-0003-0883-0669>

⁵Scientist, Department of Plant Breeding and Genetics, College of Agriculture, Vellanikkara, Thrissur 680656, Kerala Agricultural University, biju.s@kau.in, ORCID ID: 0009-0008-9775-7670

ABSTRACT:

Antibiotic resistance has emerged as a major global health concern affecting clinical treatment, environmental safety, and microbial ecosystem stability. The present study investigated genomic composition patterns associated with antibiotic-resistant bacterial genomes using comparative computational genomics approaches. A publicly available genomic dataset containing 100,000 bacterial genomic sequences with resistant and non-resistant classifications was analyzed using bioinformatics and sequence-based comparative analysis methods. The study evaluated genomic sequence length distribution, GC-content variation, nucleotide composition, and dinucleotide frequency patterns associated with antibiotic resistance. The results demonstrated relatively stable genomic sequence lengths across bacterial samples, with balanced resistant and non-resistant genomic categories. GC-content analysis revealed that resistant bacterial genomes exhibited comparatively lower GC content than non-resistant bacterial genomes. Nucleotide composition analysis further demonstrated increased adenine and thymine abundance within resistant bacterial genomes, whereas non-resistant bacterial genomes showed relatively higher guanine and cytosine composition. Dinucleotide frequency analysis identified distinct sequence-level genomic signatures associated with antibiotic resistance, particularly involving AT-, TA-, and TG-associated sequence patterns. Comparative heatmap visualization additionally demonstrated measurable compositional variation between resistant and non-resistant bacterial populations. The findings indicate that computational genomic analysis and comparative bioinformatics approaches can effectively identify resistance-associated genomic signatures and sequence-level compositional variation within bacterial genomes. Overall, the study contributes to microbial genomics and antimicrobial resistance research by demonstrating the applicability of sequence-based genomic characterization for comparative analysis of antibiotic-resistant bacterial populations across clinical and environmental contexts.

KEYWORDS: antibiotic resistance, microbial genomics, comparative genomics, bioinformatics, GC-content analysis, bacterial genome characterization

1. INTRODUCTION

Antibiotic resistance is now a serious public health problem with clinical, environmental health and microbial ecosystem impacts worldwide. A molecular characterization and genomic surveillance approach is gaining significance as a result of the speed of the spread of bacteria resistant to antibiotics in both clinical and non-clinical environments including agriculture. Recent years have seen the widespread use of whole genome sequencing (WGS), comparative genomics and bioinformatics methods to elucidate the mechanisms of antimicrobial resistance, evolutionary and transmission dynamics of bacteria (Cabal et al., 2022).

Multiple studies indicated that antibiotic-resistant bacteria are spread between interconnected human, animal and environmental systems and have the potential to spread antibiotic resistance genes across a variety of ecological habitats (Jesumirhewe et al., 2022). Multidrug-resistant microbial populations are also known to exist in environmental reservoirs and in hospital wastewater systems (Kelly et al., 2023). As a result, genomic characterization is now a crucial aspect of our understanding of the mechanisms of adaptation and resistance-associated genomic variation in bacteria.

Whole genome sequencing has been widely used for molecular characterization of clinically relevant pathogens recovered from clinical, environmental and hospital sources. Piccirilli et al. (2022) used WGS to characterise clinical and environmental strains of *S. marcescens* involved in neonatal intensive care unit outbreaks and Yasir et al. (2022) analysed CRA that were isolated from the environment of healthcare facilities. In the same vein, Wu

et al. (2022) studied the antibiotic resistomes found in inhaled air from hospitals, highlighting the significance of environmental and clinical risks of airborne spread of AMR.

The study of antibiotic resistant bacteria spread through the environment has grown in focus all over the world. Gu et al (2022), examined resistance genes and resistant bacteria in drinking water treatment systems with metagenomic methods, while Mahmud et al (2023), isolated resistant bacteria from pharmaceutical effluents and determined the characteristics of antibiotic resistance of the bacteria. In addition, Zagui et al. (2024) showed how hospital wastewater systems act as reservoirs of high-risk *Pseudomonas aeruginosa* clones and Cirkovic et al. (2023) gave whole genome sequencing analysis of clinically relevant carbapenem resistant gram-negative bacteria (CRGNB) isolated from wastewater environments.

Comparative genetic studies have also enhanced our knowledge of the genomic diversity related to bacterial adaptation and resistance. Wiradiputra et al. (2023) investigated the genotypic difference among antibiotic resistance phenotypes of *Acinetobacter baumannii* while Muzahid et al. (2023) studied the difference between the hospital associated and the community associated *Acinetobacter baumannii* using molecular epidemiological analysis. In another study, Dioli et al. (2023) took another step towards understanding the ecology of antimicrobial resistance in *E. coli* from clinical and environmental sources, underscoring the inter-tangled relationship between the spread of bacterial resistance.

Recent studies also focused on ecological distribution of bacterial resistant populations and genomic evolution of resistant populations of bacteria, specifically of *E. coli* resistant to antibiotics, as done by Śliwka et al. (2023) and of antibiotic-resistant enterococci in farm animals as done by Chen et al. (2024). These results were corroborated by research that detected *E. coli* resistant to antibiotics in food associated environments and in environment habitats (Hossain et al., 2022).

Although there has been great progress in microbial genomics and monitoring of antimicrobial resistance, fewer studies have targeted sequence level genomic composition analysis related to antimicrobial resistance. Thus, the present study was aimed at examining the patterns of variation in GC content, nucleotide composition and comparative genome sequence patterns of bacterial genomes with their ability to resist antibiotics.

The objectives of the present study were:

- To characterize genomic composition patterns associated with antibiotic-resistant bacterial genomes.
- To compare nucleotide composition and sequence-level genomic variations between resistant and non-resistant bacterial samples.
- To identify resistance-associated genomic signatures using comparative bioinformatics analysis.

2. METHODOLOGY

2.1 Research Design

This current study applied a quantitative computational genomics approach for studying patterns of genomic compositions that were linked to the genomes of bacteria having antibiotic resistance. Bioinformatics comparative analysis was done to determine the variations in terms of nucleotide compositions, GC-content difference, and other sequence-related genomic features in bacterial strains having resistance versus those without resistance.

2.2 Dataset Description

The experiment used an open genomic data set that comprised the sequences of bacteria as well as their resistance to antibiotics (Scarlat, 2019). This data set comprised 100,000 samples of bacteria, which included both resistant and non-resistant bacterial genomes. Every sample included nucleotide sequences described with the help of DNA bases (A, T, G, and C), as well as a binary label showing whether the genome is resistant or not. The reason for choosing this data set for the analysis was that it was appropriate for comparing bacterial genomes and assessing their antimicrobial resistance.

2.3 Data Preprocessing

The data pre-processing step was performed using Python. The genome dataset was loaded as a NumPy array and was then transformed into a structured pandas DataFrame for analysis purposes. The length of the genomic sequences for each bacterium was computed for assessing consistency in the dataset. The resistance labels were encoded into binary classes for comparison. Quality checks were done on the dataset which showed that there were no missing values and duplicated genomes.

2.4 GC-Content Analysis

GC-content analysis was performed to investigate genomic compositional variation among bacterial samples. GC content was calculated as the percentage of guanine (G) and cytosine (C) nucleotides relative to the total sequence length using the following equation:

$$[GC\ Content = \frac{(G + C)}{Total\ Sequence\ Length} \times 100]$$

The calculated GC-content values were compared between resistant and non-resistant bacterial genomes to identify resistance-associated compositional differences.

2.5 Nucleotide Composition Analysis

The nucleotide composition analysis was done to determine the proportional ratio of nucleotides A, T, G, and C within the genome of bacteria. The composition percentage of the four different nucleotides in respect to their lengths was determined using genomic sequence data. A comparative analysis of the four nucleotides was then done using sequences from both resistant and non-resistant bacteria.

2.6 Dinucleotide (k-mer) Analysis

Dinucleotide frequency analysis was carried out as a comparative genomic profiling method that would help in determining sequence-specific genomic fingerprints. All the combinations of dinucleotides were created by computer modeling based on bioinformatics sequence manipulation methods. The frequencies of all the dinucleotides for each sequence were determined and normalized based on the total number of dinucleotides. Comparative dinucleotide profiling was done by comparing bacterial samples that were resistant and not resistant.

2.7 Data Visualization and Statistical Analysis

Data exploration with respect to genomic analyses and visualizations was done using various Python modules such as pandas, NumPy, matplotlib, and seaborn. Histograms, bar graphs for comparison purposes, and heat maps were created to visually show the distribution of GC content, patterns of nucleotide composition, and variability in dinucleotide composition respectively. Descriptive statistics were computed to analyze the characteristics of genomic sequences and their comparisons.

3. RESULTS

3.1 Dataset Characteristics and Genomic Sequence Distribution

The genomic data set included 100,000 genomic sequences of bacteria with both resistant and non-resistant bacteria samples. The quality test showed that there were no missing or duplicate data entries. The study revealed consistency in the size of genomic sequences of bacteria where the average sequence length was found to be 149.99 nucleotides.

Table 1. Genomic Sequence Length Statistics

Statistic	Sequence Length
Count	100000
Mean	149.99
Standard Deviation	2.93
Minimum	144
25th Percentile	149
Median	150
75th Percentile	151
Maximum	156

The resistance distribution analysis demonstrated balanced genomic categories within the dataset. Non-resistant bacterial genomes accounted for 50,297 samples, whereas resistant bacterial genomes accounted for 49,703 samples. The balanced resistance distribution supported reliable comparative genomic characterization.

Table 2. Distribution of Resistant and Non-Resistant Bacterial Genomes

Resistance Status	Frequency
Non-Resistant	50,297
Resistant	49,703

3.2 GC-Content Analysis

The GC content was determined in order to determine any variations in the genomic composition in the bacterial samples that were either resistant or not resistant to the antibiotics. The average GC content in the bacterial samples was 42.65%.

Table 3. GC-Content Statistics of Bacterial Genomes

Statistic	GC Content (%)
Mean	42.65
Standard Deviation	0.60
Minimum	40.00
25th Percentile	42.00
Median	42.67
75th Percentile	43.05
Maximum	45.33

Comparative analysis revealed slight compositional differences between resistant and non-resistant bacterial genomes. Non-resistant bacterial genomes exhibited marginally higher average GC content (42.82%) compared to resistant bacterial genomes (42.49%).

Table 4. Average GC Content by Resistance Status	
Resistance Status	Average GC Content (%)
Non-Resistant	42.82
Resistant	42.49

The GC-content distribution demonstrated compositional genomic variation between resistant and non-resistant bacterial populations. Resistant genomes showed relatively concentrated GC-content patterns, whereas non-resistant bacterial genomes exhibited broader genomic variability.




Figure 1. GC Content Distribution in Resistant and Non-Resistant Bacteria

3.3 Nucleotide Composition Analysis

Nucleotide composition analysis demonstrated that adenine (A) and thymine (T) represented the dominant nucleotide components across bacterial genomic sequences, while guanine (G) and cytosine (C) showed comparatively lower abundance.

Table 5. Nucleotide Composition Statistics		
Nucleotide	Mean (%)	Standard Deviation
Adenine (A)	28.68	0.53
Thymine (T)	28.67	0.50
Guanine (G)	21.30	0.51
Cytosine (C)	21.35	0.53

Comparative nucleotide analysis further demonstrated compositional variation between resistant and non-resistant bacterial genomes. Resistant bacterial genomes exhibited slightly increased adenine and thymine composition, whereas non-resistant genomes showed relatively higher guanine and cytosine content.

Table 6. Average Nucleotide Composition by Resistance Status		
Nucleotide	Non-Resistant (%)	Resistant (%)
Adenine (A)	28.59	28.77
Thymine (T)	28.59	28.74
Guanine (G)	21.35	21.25
Cytosine (C)	21.47	21.24

The nucleotide composition patterns indicated sequence-level compositional differences associated with antibiotic resistance characteristics.




Figure 2. Nucleotide Composition in Resistant and Non-Resistant Bacteria

3.4 Dinucleotide (k-mer) Analysis

The frequency analysis of dinucleotides was performed to obtain genomic fingerprints that correlate with resistance to antibiotics. The comparative genomic study indicated that AC, TG, AT, TT, and CT belong to the most frequent dinucleotide motifs in bacterial genomes. Comparative analysis showed differences in the distribution of dinucleotide frequencies in bacteria that are resistant and non-resistant. Genomes of resistant bacteria have relatively more TA-, TG-, and GA-related sequences, while non-resistant bacteria are rich in GC- and CC-related frequencies.

Table 7. Major Dinucleotide Frequencies in Bacterial Genomes

Dinucleotide	Non-Resistant	Resistant
AC	0.100	0.100
TG	0.105	0.106
AT	0.086	0.087
TT	0.074	0.074
CT	0.074	0.074
GC	0.047	0.047
CC	0.021	0.020

The heatmap analysis highlighted comparative sequence-level genomic variation between resistant and non-resistant bacterial genomes and demonstrated distinct genomic signature patterns associated with antibiotic resistance.

3.5 Comparative Genomic Interpretation

In general, it was found through the comparative genomics study that the genome of antibiotic-resistant bacteria shows variation in terms of its composition and also in terms of its sequence as compared to the genome of non-resistant bacteria. The bacteria that were resistant to antibiotics had low levels of GC content along with a higher frequency of A/T nucleotides, indicating possible adaptation patterns due to their antibiotic resistance properties.

4. DISCUSSION

In the present study, the antibiotic-resistant bacterial genome was studied through comparative computational genomics approaches to find out the genomic composition patterns associated with the antibiotic-resistant bacterial genome. The results showed that the sequence level difference between resistant and non-resistant bacterial populations was significant with respect to GC content distribution, nucleotide composition and dinucleotide frequency patterns. These findings align with the findings from the previous genomic studies, highlighting the significance of molecular characterisation and comparative genomics in the understanding of mechanisms for antimicrobial resistance and adaptations of bacteria.

The results of the GC content analysis showed that there are some differences in the composition of the resistant and non-resistant bacterial samples in the bacterial genome; the GC percentage of the resistant bacterial samples was relatively lower. The same type of genomic compositional difference has been observed in resistant bacterial isolates from hospital wastewater systems and environmental reservoirs. To understand the evolution of bacteria with resistance, Xie et al. (2023) showed that there are significant genomic differences among carbapenem-resistant *Klebsiella pneumoniae* (CRKP) recovered from hospital wastewater. Similarly, Sukhum et al. (2022)

found that the built environment in hospitals harbors antibiotic resistance reservoirs, thus demonstrating the importance of environmental microbial adaptation in the dissemination of antibiotic resistance.

The nucleotide composition analysis also revealed that the adenine and thymine percentage in resistant bacterial genome was slightly higher but the guanine and cytosine was comparatively higher in non-resistant bacterial genome. The results indicate possible correlation between the antibiotic resistance and the sequence-level compositional changes and genomic adaptation pattern. Hossain et al. (2022) also reported similar observations, in which they used genetic analysis approaches on the characterization of antibiotic-resistant *Escherichia coli* from food associated environments. They found there was a significant amount of genomic diversity in resistant bacteria that was found in environmental and food sources.

The dinucleotide frequency analysis further identified genomic signatures unique to each sequence that were linked to antibiotic resistance. The AT, TA and TG dinucleotide frequencies were greater relative to non-resistant genomes for resistant bacterial genomes. These sequence variations at the sequence level could be indicative of adaptive mutational patterns to survive antimicrobial pressures in bacteria. Fatoba and co-authors (2022) also reported similar results when they performed comparative genomic analysis to detect plasmid-mediated resistance genes among environmental *Enterococcus* species. The study focussed on the transference of environmental resistance through the action of mobile genetic elements and genomic evolution.

The current results also highlight the need for precise genomic characterization in the identification of resistance-associated bacterial populations, which is increasingly recognized. Pongchaikul et al. (2023) showed that genomic data can help in refining the identification of antibiotic resistance genes in clinical isolates of *Enterococcus faecium*, aiding in the precise identification of microbial pathogens and more effective antibiotic treatment. Likewise, Osińska et al. (2023) pointed out the key issues of identification and genetic characterization of *E. coli* from environmental samples, which indicated that microbial genomics in the environment is complicated and the surveillance of resistance is challenging.

The results of the present study show that computational genomic analysis, nucleotide composition profiling and comparative sequence characterization can be useful tools for investigations of antimicrobial resistance. Even with the simple genome sequence data sets and absence of species-specific annotation and environmental metadata, the results of the present study showed a distinct compositional and sequence-driven variation linked to the acquisition of antibiotic resistance. The results help fill the growing field of comparative microbial genomics and aid in the use of bioinformatic tools for genomic surveillance of antibiotic-resistant populations of bacteria.

Further studies will help to enhance the understanding of the evolution of antimicrobial resistance in both clinical and environmental ecosystems, by considering species-level genomic annotation, whole genome sequencing, resistance gene profiling and phylogenetic analyses.

5. CONCLUSION

The current study was aimed at examining patterns of genomic composition of antibiotic-resistant bacterial genomes with the help of comparative computational genomics tools. The results showed that the resistant and non-resistant bacteria samples exhibited statistically significant differences in a number of sequence level properties, including the distribution of GC-contents, distribution of nucleotides and dinucleotide frequency patterns. The resistant bacterial genomes had relatively low GC content and high adenine/thymine associated sequence composition, which may indicate some genomic adaptation due to the mechanism of antimicrobial resistance. The study also showed that sequence signatures associated with bacterial resistance can be successfully identified from bacterial genomic data through comparative genomic characterization and computational bioinformatics analysis. Di-nucleotide analysis showed that there were distinct patterns in the composition of the sequences associated with the resistant bacterial populations, reinforcing the increasing relevance of genomic surveillance and molecular characterization in antimicrobial resistance research. The results add to the growing body of knowledge in the field of microbial genomics, showing the potential for sequence-based genomic analysis in comparative characterization of antibiotic resistant bacterial genome. The study also emphasizes the necessity of Genomic composition analysis, Nucleotide profiling and Comparative sequence analysis to understand the adaptation and evolution of resistance in bacteria in clinical and environmental settings. The present study focused on the comparative genomic analysis of the data set, with no species-specific annotations or environmental data provided, but the analysis identified compositional variation related to antibiotic resistance. Whole genome sequencing, resistance gene profiling, phylogenetic analysis and species-level genomic annotation should be combined with existing datasets to augment the understanding of the transmission of antimicrobial resistance and genomic diversity of bacteria in clinical and environmental ecosystems.

REFERENCES

1. Cabal, A., Rab, G., Daza-Prieto, B., Stöger, A., Peischl, N., Chakeri, A., ... & Ruppitsch, W. (2022). Characterizing antimicrobial resistance in clinically relevant bacteria isolated at the human/animal/environment interface using whole-genome sequencing in Austria. *International journal of molecular sciences*, 23(19), 11276.
2. Jesumirhewe, C., Springer, B., Allerberger, F., & Ruppitsch, W. (2022). Genetic characterization of antibiotic resistant *Enterobacteriaceae* isolates from bovine animals and the environment in Nigeria. *Frontiers in Microbiology*, 13, 793541.

3. Piccirilli, A., Cherubini, S., Brisdelli, F., Fazii, P., Stanziale, A., Di Valerio, S., ... & Perilli, M. (2022). Molecular characterization by whole-genome sequencing of clinical and environmental *Serratia marcescens* strains isolated during an outbreak in a neonatal intensive care unit (NICU). *Diagnostics*, 12(9), 2180.
4. Kelly, S. A., O'Connell, N. H., Thompson, T. P., Dillon, L., Wu, J., Creevey, C., ... & Dunne, C. P. (2023). Large-scale characterization of hospital wastewater system microbiomes and clinical isolates from infected patients: profiling of multi-drug-resistant microbial species. *Journal of Hospital Infection*, 141, 152-166.
5. Yasir, M., Subahi, A. M., Shukri, H. A., Bibi, F., Sohrab, S. S., Alawi, M., ... & Azhar, E. I. (2022). Bacterial community and genomic analysis of carbapenem-resistant *Acinetobacter baumannii* isolates from the environment of a health care facility in the western region of Saudi Arabia. *Pharmaceuticals*, 15(5), 611.
6. Chen, W., Wang, Q., Wu, H., Xia, P., Tian, R., Li, R., & Xia, L. (2024). Molecular epidemiology, phenotypic and genomic characterization of antibiotic-resistant enterococcal isolates from diverse farm animals in Xinjiang, China. *Science of the Total Environment*, 912, 168683.
7. Wu, D., Jin, L., Xie, J., Liu, H., Zhao, J., Ye, D., & Li, X. D. (2022). Inhalable antibiotic resistomes emitted from hospitals: metagenomic insights into bacterial hosts, clinical relevance, and environmental risks. *Microbiome*, 10(1), 19.
8. Mahmud, M. S., Hosen, M. A., Hossion, M. I., Sadik Sabuj, M. S., Rumi, N. A., Hossain, M. K., ... & Bourhia, M. (2023). Isolation, identification, and characterization of resistant bacteria to antibiotics from pharmaceutical effluent and study of their antibiotic resistance. *Frontiers in microbiology*, 14, 1307291.
9. Gu, Q., Sun, M., Lin, T., Zhang, Y., Wei, X., Wu, S., ... & Wu, Q. (2022). Characteristics of antibiotic resistance genes and antibiotic-resistant bacteria in full-scale drinking water treatment system using metagenomics and culturing. *Frontiers in Microbiology*, 12, 798442.
10. Zagui, G. S., de Almeida, O. G. G., Moreira, N. C., Silva, N. G. A., Meschede, M. S. C., Darini, A. L. C., ... & Segura-Muñoz, S. I. (2024). Hospital wastewater as source of human pathogenic bacteria: A phenotypic and genomic analysis of international high-risk clone VIM-2-producing *Pseudomonas aeruginosa* ST235/O11. *Environmental Research*, 255, 119166.
11. Śliwka, P., Weber-Dąbrowska, B., Żaczek, M., Kuźmińska-Bajor, M., Dusza, I., & Skaradzińska, A. (2023). Characterization and comparative genomic analysis of three virulent *E. coli* bacteriophages with the potential to reduce antibiotic-resistant bacteria in the environment. *International journal of molecular sciences*, 24(6), 5696.
12. Wiradiputra, M. R. D., Thirapanmethee, K., Khuntayaporn, P., Wanapaisan, P., & Chomnawang, M. T. (2023). Comparative genotypic characterization related to antibiotic resistance phenotypes of clinical carbapenem-resistant *Acinetobacter baumannii* MTC1106 (ST2) and MTC0619 (ST25). *BMC genomics*, 24(1), 689.
13. Muzahid, N. H., Hussain, M. H., Huët, M. A. L., Dwiyanto, J., Su, T. T., Reidpath, D., ... & Rahman, S. (2023). Molecular characterization and comparative genomic analysis of *Acinetobacter baumannii* isolated from the community and the hospital: an epidemiological study in Segamat, Malaysia. *Microbial Genomics*, 9(4), 000977.
14. Dioli, C., Pappa, O., Siatravani, E., Bratakou, S., Tatsiopoulos, A., Giakkoupi, P., ... & Beloukas, A. (2023). Molecular characterization and prevalence of antimicrobial-resistant *Escherichia coli* isolates derived from clinical specimens and environmental habitats. *Microorganisms*, 11(6), 1399.
15. Cirkovic, I., Muller, B. H., Janjusevic, A., Mollon, P., Istier, V., Mirande-Meunier, C., & Brkic, S. (2023). Whole-genome sequencing snapshot of clinically relevant carbapenem-resistant gram-negative bacteria from wastewater in Serbia. *Antibiotics*, 12(2), 350.
16. Xie, Z., Huang, J., Zhang, S., Xu, B., Zhang, Q., & Li, B. (2023). Genomic and functional characterization of carbapenem-resistant *Klebsiella pneumoniae* from hospital wastewater. *BMC microbiology*, 23(1), 115.
17. Sukhum, K. V., Newcomer, E. P., Cass, C., Wallace, M. A., Johnson, C., Fine, J., ... & Kwon, J. H. (2022). Antibiotic-resistant organisms establish reservoirs in new hospital built environments and are related to patient blood infection isolates. *Communications Medicine*, 2(1), 62.
18. Hossain, M. M. K., Islam, M. S., Uddin, M. S., Rahman, A. M., Ud-Daula, A., Islam, M. A., ... & Alam, J. (2022). Isolation, identification and genetic characterization of antibiotic resistant *Escherichia coli* from frozen chicken meat obtained from supermarkets at Dhaka City in Bangladesh. *Antibiotics*, 12(1), 41.
19. Fatoba, D. O., Amoako, D. G., Akebe, A. L. K., Ismail, A., & Essack, S. Y. (2022). Genomic analysis of antibiotic-resistant *Enterococcus* spp. reveals novel enterococci strains and the spread of plasmid-borne Tet (M), Tet (L) and Erm (B) genes from chicken litter to agricultural soil in South Africa. *Journal of environmental management*, 302, 114101.
20. Pongchaikul, P., Romero, R., Mongkolsuk, P., Vivithanaporn, P., Wongsurawat, T., Jenjaroenpun, P., ... & Chaemsaitong, P. (2023). Genomic analysis of *Enterococcus faecium* strain RAOG174 associated with acute chorioamnionitis carried antibiotic resistance gene: is it time for precise microbiological identification for appropriate antibiotic use?. *BMC genomics*, 24(1), 405.
21. Osińska, A., Korzeniewska, E., Korzeniowska-Kowal, A., Wzorek, A., Harnisz, M., Jachimowicz, P., ... & Zieliński, W. (2023). The challenges in the identification of *Escherichia coli* from environmental samples and their genetic characterization. *Environmental Science and Pollution Research*, 30(5), 11572-11583.
22. Scarlet, A. (2019). GeneResistData: 100k samples of bacterial genomic sequence and resistance to antibiotic [Data set]. Kaggle. <https://www.kaggle.com/datasets/drscarlat/dzd-data>