

EPIGENETIC CONTROL OF TRANSCRIPTIONAL PLASTICITY UNDER MULTI-STRESS ENVIRONMENTAL CONDITIONS

Dr. Muninathan N¹, Dr. Karpagavalli², Dr. Durga Devi G³

¹Scientist, Central Research Laboratory, Meenakshi Medical College Hospital & Research Institute, Meenakshi Academy of Higher Education and Research, Enathur, Kanchipuram, Tamil Nadu – 631552, India, Email: muninathan@maher.ac.in

²Professor cum HoD, Pharmaceutics, Meenakshi College of Pharmacy, Meenakshi Academy of Higher, education and Research, Email: karpagavalli@maher.ac.in

³Professor & HOD, Anatomy, ORCID: <https://orcid.org/0000-0003-1852-8711>

ABSTRACT

Transcriptional plasticity is a key biological mechanism through which organisms can dynamically regulate gene expression to changing environmental conditions. This adaptive ability is especially vital during times of stress that is; during rapid and synchronized responses by molecules, survival and resilience are acquired. Although there is an increasing interest in epigenetic regulation, the mechanisms of transcriptional plasticity in the combined or multi-stress conditions are poorly understood. Specifically, it remains unclear how particular stress signals are combined at the epigenetic level to tune gene expression programs. Our hypothesis in this work is that multi-stress exposure causes the coordinated epigenetic remodeling in increasing transcriptional plasticity by modulating chromatin accessibility and gene-regulatory network dynamics. To examine this, a multi-omics (combining transcriptomic (RNA-seq), epigenomic (DNA methylation profiling), and chromatin state (ChIP-seq of histone modifications) analyses were used under regulated single and combined stresses. The analysis of differential gene expression showed that when stress is exposed in multi-stress, the stress-responsive transcriptional activity of cells increased significantly as compared to when the same cells were subjected to a single stress treatment. At the same time, the epigenetic profiling revealed a significant drop in promoter methylation and enrichment of activating histone marks (e.g., H3K27ac, H3K4me3) of core regulatory genes, with reference to increased chromatin accessibility. Integrated multi-omics analysis also revealed closely orchestrated regulatory networks of stress-responsive transcription factors and signaling pathways such as MAPK and reactive oxygen species (ROS)-mediated signaling. These results are an indication that the effect of multi-stress condition is a synergism of the effect of individual stresses and not additive. All in all, this paper offers proof that under complicated environmental circumstances, epigenetic processes are of prime importance in concerting transcriptional plasticity. Results emphasize how the chromatin-level regulation may be paramount to adaptive responses and provide a new understanding of the molecular foundations of stress resilience, which can be subsequently utilized in agriculture, environmental biology, and biomedical research.

KEYWORDS: Epigenetics, transcriptional plasticity, multi-stress response, DNA methylation, histone modification, chromatin remodelling, RNA-seq, gene regulation.

1. INTRODUCTION

Transcriptional plasticity is the capacity of organisms to dynamically tune gene expression to changing environmental conditions, allowing a quick adaptation to, and survival in a stressful situation. It is an inherent aspect of biological systems that enable plasticity to affect developmental processes, stress tolerance, and evolutionary fitness (Duncan et al., 2022). Complex transcriptional responses, reprogramming cellular functions, can be elicited by environmental stressors like the temperature change, oxidative stress, and pathogen exposure. These adaptive output levels are not only controlled by the genetic sequences but are rapidly being acknowledged to be controlled by epigenetic processes that offer a flexible and reversible overlay of regulation to the expression of genes (Feil & Fraga, 2012). Epigenetic regulation is very important in the regulation of transcriptional activity in the absence of modifying the underlying DNA sequence. The most significant ones are DNA methylation, histone alterations, and chromatin remodeling, which affect accessibility of chromatin and transcription factor binding (Atlasi & Stunnenberg, 2017). DNA methylation is generally a repressive marker, but the context of histone modifications can activate or suppress transcription, which includes acetylation and methylation. The processes of these epigenetic mechanisms are very dynamic and sensitive to environmental signals that enable cells to respond quickly to stress changing their gene expression patterns (Szyf et al., 2016). Moreover, those genetic alterations caused by stress have been demonstrated to control the genes related to resilience, adaptation to metabolic circumstances, and cell recovery processes (Matosin et al., 2017; Miller et al., 2025).

Although there has been a lot of progress in known epigenetic regulation, most studies have been done on response to individual stress conditions, e.g. heat, drought or oxidative stress. Although these studies have yielded useful

information, they do not reflect the real world situation where organisms are often subjected to a combination of several stressors. There is evidence that the pressures of combined conditions are not just additive but lead to distinct and in many cases synergistic transcriptional responses (Loughland et al., 2021). This shows that there are built in regulatory processes that orchestrate gene expression involving a combination of signaling pathways. Multi-stress environmental situations add even more compliance of regulation, necessitating the assembly of different signaling networks and epigenetic alterations. The conditions have the ability to change the chromatin structure, break the regulatory feedback loops, as well as activate different transcription factor networks, resulting in non-linear patterns of gene expression. An example is that epigenetic reprogramming under stress has been associated with both immediate adaptive changes and enduring memory of stress especially in plant and cell systems (Tan, 2026). Additionally, the new literature indicates that epigenetic signaling might serve as key integrators of the environmental cues, and that transcriptional responses to a range of stress pathways are coordinated by epigenetic signaling.

Nonetheless, the critical research gap concerning the need to learn how a combination of epigenetic alterations can control the potential of transcriptional plasticity in multi-stress situations still persists. The existing models do not seem to extensively incorporate epigenomic and transcriptomic data and, therefore, do not allow us to comprehensively describe the way a particular adaptive response is triggered at the molecular level. Specifically, the interplay among DNA methylation, histone modification and chromatin accessibility in determining gene regulatory networks during combined stress is still under-researched. Our observations suggest we can test the hypothesis that multi-stress exposure triggers coordinated epigenetic remodeling that promotes transcriptional plasticity by dynamically changing chromatin accessibility and gene regulatory network interactions. This study aims to examine the epigenetic mechanisms of transcriptional plasticity in the context of multi-stress conditions in the environment using an integrated multi-omics strategy. This study will use transcriptomic analysis and epigenomic analysis to reveal important regulatory pathways and molecular signatures that are related to adaptive stress response.

This work offers a new integrative model of the relationships between the mechanisms of epigenetic regulation in coordinating transcriptional plasticity to the complex environment. In contrast to traditional single-stress experiments, it has identified the nature of multi-stress reactions as synergistic and non-linear and provided novel understanding into the regulatory landscape of chromatin and gene network responses. The results help in the future of stress biology and epigenomics and have potential applications in creating stress-resistant crops, enhancing better strategies to adapt to environmental changes, and inform biomedical research on stress-related diseases.

2. Molecular Basis of Epigenetic Regulation

Epigenetic regulation is an essential dimension of gene expression regulation that facilitates organisms to react dynamically in response to environmental cues even without causing a change in the underlying DNA sequence. These processes play a key role in transcriptional plasticity, especially during stressful responses when there is a need in the fast and reversible changes in gene expression. Epigenetic mechanisms work via several interrelated pathways, comprising of DNA methylation, histone modifications, chromatin remodeling, and non-coding RNA-based regulation. The combination of these processes affects the chromatin structure and accessibility which in turn regulated the transcriptional activity in relation to environmental signals.

2.1 DNA Methylation

One of the most widely researched epigenetic modifications is DNA methylation, which is a key to controlling gene expression. This is done by the methylation of the 5-carbon site of cytosine bases, most commonly found in a CpG dinucleotide. Transcriptional repression, in general, is linked with CpG methylation, especially at promoter regions, since it prevents transcription factor binding and recruitment of repressive protein complexes. As a result, hypermethylation commonly results in silencing of genes, and hypomethylation with transcriptional activation. DNA methylation patterns are very plastic under stress conditions. Methylation profiles can change on a genome-wide scale due to environmental stressors, including changes in temperature, oxidative stress and nutrient deprivation. These active changes in methylation allow the selective activation or repression of stress-responsive genes, which allow adaptive responses. Notably, the changes in stress-induced methylation may be short and long-term, with the contribution to both short-term adaptation and epigenetic memory. This flexibility highlights the importance of DNA methylation as an important modulator of transcriptional plasticity in a multi-stress setting.

2.2 Histone Modifications

Another crucial part of the epigenetic regulation process is histone modifications, which can alter the structure of chromatin and modulate the expression of genes. Histone proteins, which surround DNA to create nucleosomes, are the subject to a broad spectrum of post-translational alterations, such as acetylation, methylation, phosphorylation and ubiquitination. These changes modify the relationship between histones and DNA and, thus, control chromatin availability. Certain histone marks are linked to either transcriptional activation or repression. The activation marks, e.g., histone H3 lysine 4 trimethylation (H3K4me3) and histone H3 lysine 27 acetylation (H3K27ac), are generally localized to active genes and promoter regions, and activate an open chromatin structure that facilitates binding of transcription factors. Conversely, repressive marks like histone H3 lysine 27 trimethylation (H3K27me3) are linked with compaction in the chromatin and transcription silencing. In multi-stress situations, histone modification patterns are very dynamic, indicating the necessity of fast transcriptional reprogramming. The aligning deposition and erasing of activating and repressive marks enable cells to modulate

gene expression to intricate environmental signals. This dynamic histone environment has a significant impact on the plasticity and reversibility of transcriptional plasticity.

2.3 Chromatin Remodeling

Chromatin remodeling is a central mechanism that governs the physical structure of chromatin, and thus, determines access to genetic information. In contrast to the chemical modifications, like methylation or acetylation, chromatin remodeling entails the ATP-dependent protein complex, repositioning, ejection or restructuring of the nucleosomes along the DNA. The SWI/SNF (Switch/Sucrose Non-Fermentable) family of chromatin remodeling complexes is one of them, and these are essential in assisting with transcriptional activation. These complexes exploit the hydrolysis of ATP to change the position of the nucleosome exposing promoter and enhancer regions to transcription factors and RNA polymerase. Chromatin remodeling is a highly flexible regulatory mechanism as nucleosome positioning promotes or suppresses gene expression based on context. Chromatin remodeling is specifically valuable in multi-stress settings as this allows the rapid response of transcription. The reconfiguration of chromatin accessibility can be caused by stress signaling the recruitment of remodeling complexes to particular genomic loci. This mechanism guarantees that genes that respond to stress are quickly activated and non-essential genes are suppressed, which help to efficiently allocate resources and adapt cells.

2.4 Non-Coding RNA Regulation

Another level of gene regulation, introduced by non-coding RNAs (ncRNAs), is an interaction between them and genes in different ways. These non-protein coding RNA molecules have a very important role in the regulation of transcriptional and post-transcriptional processes. MicroRNAs (miRNAs) and long non-coding RNAs (lncRNAs) are some of them that are quite instrumental in transcriptional plasticity regulation. miRNAs usually work through binding to target messenger RNAs (mRNAs) with complementary sequences, which results in mRNA degradation or translational repression. This system enables the precise control of gene expression, especially in response to stresses when there is need to modulate the level of protein rapidly. The miRNA expression profiles in response to stress have been reported to be able to regulate responses and survival-related pathways. lncRNAs, however, act at various levels to control gene expression, such as chromatin modification, transcriptional regulation, and RNA processing. They are able to bind directly to chromatin modifying complexes and can direct them to definite regions of the genome to either activate or silence the expression of genes. lncRNAs are important in establishing the epigenetic environment and orchestrating multifaceted transcriptional reactions, through these interactions. ncRNAs that are present in the multi-stress environments serve as very important regulators of integration of signals across multiple pathways, therefore, responding to coordinated gene expressions. They are also necessary to retain transcriptional plasticity in complex environmental conditions due to their capacity to tune both epigenetic states and transcriptional networks.

3. Multi-Stress Environmental Interaction Model

In the natural environments, organisms are not often exposed to one form of stress, rather, they are subjected to complexes of both abiotic and biotic stresses which may be concomitant or sequential. Such multi-stress reactions have great burdens on cellular mechanisms which demand a dynamic and well-organized regulation. In contrast, the multi-stress environments in contrast to single-stress situations result in extremely complicated and in many cases, non-linear changes in gene expression. This biological complexity comes with the combination of various signaling pathways and epigenetic changes, which together maintain transcriptional plasticity and adaptive capacity.

3.1 Types of Stress

It is possible to broadly divide environmental stressors into abiotic and biotic, all of which have a profound effect on gene expression and epigenetic status. Physical and chemical factors like heat, drought, salinity and oxidative stress are abiotic stresses. These stresses cause disturbances in cellular homeostasis including protein stability, membrane integrity, and metabolism. As an example, heat stress has the ability to denature proteins and trigger heat shock response pathways, whereas oxidative stress results in the buildup of reactive oxygen species (ROS) that damage DNA, lipids and proteins. Biogenic stresses, on the other hand, are caused by interactions with living organisms, e.g., pathogens, pests and microbial toxins. The responses to these stresses stimulate immune and defense-related pathways, which frequently entail intricate signal transduction cascades and transcriptional remodelling. Notably, abiotic and biotic stresses often are interconnected and as a result, result in overlapping and occasionally even conflicting cellular responses. The concomitant presence of multifactorial stressors will require a unified regulatory system with the capability to offset competing physiological needs.

3.2 Stress Crosstalk Mechanisms

The intricate crosstalk between signaling pathways mediates the effect of multi-stress conditions on the cellular response. Common signaling molecules, especially reactive oxygen species (ROS), are of a central concern both as damaging components and signaling intermediates. ROS serve as second messengers to stimulate downstream signaling pathways, such as mitogen-activated protein kinase (MAPK) cascades, to promote the expression of gene responses to stress. Moreover, hormonal signaling pathways, including those based on abscisic acid (ABA), salicylic acid (SA) and jasmonic acid (JA) can be viewed as prominent integrators of stress signals, which orchestrate the responses to different types of stress. Transcription factors are key nodes in this crosstalk network and combine signals of a variety of pathways and adjust the gene expression in response. Several stress signals can be combined to activate stress-responsive transcription factors, which include heat shock factors (HSFs),

dehydration-responsive element-binding proteins (DREBs) and NAC domain proteins. Their combinatorial interactions allow the modulation of gene networks that are set-specific to specific combinations of stressors and not to individual stressors. This underscores the role of the dynamics of transcription factors in the context-dependent transcriptional plasticity.

3.3 Systems-Level Gene Regulation

Gene regulation in multi-stress conditions at the systems level is best comprehended as a network based process, and not a linear pathway. Gene regulatory networks are networks of interconnected nodes (genes, transcription factors and epigenetic modifiers), linked by regulatory interactions. These networks have emergent properties, such as robustness, adaptability, the capacity to respond to a variety of responses using a limited input. Feedback loops are important in the stability of systems and the ability to respond adaptively. Stress signals can be intensified by positive feedback loops which keep the defense mechanisms quickly activated, whereas negative feedback loops can prevent overactivation and homeostasis is restored after mitigating the stress. Furthermore, epigenetic changes are also involved in stabilization of such regulatory states so that cells can have a memory of the previous stresses and react better to the future challenges. These regulatory networks are extremely dynamic under multi-stress conditions, changing both the state of gene expression and epigenetics. Combination of various signaling inputs gives way to non-linear responses that one will not be able to predict by researching each stress factor independently. Thus, systems-level view of epigenetic regulation of transcriptional plasticity and how adaptive responses to complex environmental conditions can be orchestrated by epigenetics is necessary.

4. MATERIALS AND METHODS

4.1 Experimental Design

The aim of the study was to examine transcriptional and epigenetic responses to single and combined stress conditions using a controlled experimental design. The model system was *Arabidopsis thaliana* (Col-0 ecotype) because the genome is well characterized and can be subjected to multi-omics analyses. Plants were grown under controlled growing conditions of 22°C and 16-hour light and 8-hour dark photoperiod during 21 days before stress treatment. The experimental conditions were divided into three main groups: control group that remained under normal conditions, single stress group where subjects exposed to either heat or oxidative stress, and a multi-stress group where subjects were exposed to both stresses. There were three independent biological replications ($n = 3$) of each given condition and each replication contained pooled leaf tissue of ten individual plants in total to minimize biological variation. Overall, twenty-seven samples were subjected to the analysis under all the conditions of the experiment.

4.2 Stress Conditions

The application of stress treatments was done under controlled conditions in a laboratory to enhance reproducibility and consistency. The stress of heat was induced by subjecting the plants to a temperature of 42°C over the period of 3 hours, which is known to induce vigorous transcriptional responses in plants. Foliar treatment was done with 10 mM hydrogen peroxide (H_2O_2) and exposure was conducted after three hours under normal light conditions. In the multi-stress experiment, plants were subjected to both heat stress (42°C) and oxidative stress (10 mM H_2O_2) concurrently, and at the same time. As soon as the exposure to the stressor was over, the leaf tissues were harvested and frozen in liquid nitrogen and frozen at -80°C to maintain the RNA and the epigenetic conditions intact until future analysis.

4.3 Data Acquisition

The TRIzol reagent was used to extract total RNA and the Agilent Bioanalyzer was used to check the quality of RNA with a minimum RNA Integrity Number (RIN) of 7.5 being used to ensure that all samples had a minimum of 7.5. Illumina TruSeq RNA Library Preparation Kit was used to prepare RNA sequencing libraries and sequenced on the Illumina NovaSeq 6000 platform with 150-base-paired reads per sample of about 30 to 40 million paired-ends. In the case of epigenetic profiling, chromatin immunoprecipitation sequencing (ChIP-seq) is conducted in connection with histone mark antibodies related to transcriptional activation, such as H3K27ac and H3K4me3. About 20-25 million high-quality reads were obtained in each ChIP-seq sample. One particular DNA methylation experiment was performed by whole-genome bisulfite sequencing, with an average coverage of 15-20 times per CpG site which provided sufficient precision in quantifying the levels of methylation throughout the genome.

4.4 Bioinformatics Analysis

Bioinformatics pipelines were used to process the sequencing data. HISAT2 was used to align RNA-seq reads to the *Arabidopsis thaliana* reference genome (TAIR10) and featureCounts was used to quantify gene expression. DESEQ2 was used to analyse differential gene expression with a threshold of an absolute log₂ fold change of greater equal to 1.5 and adjusted p-value of less than 0.05. The ChIP-seq data were aligned with Bowtie2 and MACS2 was used to call peaks that had a q-value lower than 0.01 to mark the significant enrichment regions. Bismark was used to handle the DNA methylation data, which were then analyzed as the percentage of methylated cytosines at CpG sites. Differentiated methylated regions were found using a cut-off threshold of 25% and significant difference of $p < 0.01$. Differentiated expressed genes were analyzed through functional enrichment analysis based on Gene Ontology and KEGG pathways databases, and statistical significant was set at the false discovery rate set at 0.05.

4.5 Statistical Analysis

R software (4.2.0) was used to conduct all the statistical analyses. One-way analysis of variance (ANOVA) was used to test the differences within experimental groups, and it was supplemented by a post hoc test which compares a number of different variables (Tukey, 2009). Student t-test was used in situations where necessary to carry out pair wise comparisons. The BenjaminiHochberg technique was used to control multiple hypothesis testing and the false discovery rate was calculated and a FDR value below 0.05 was taken as a statistically significant value. The Pearson correlation analysis was used to evaluate the correlation between the extent of DNA methylation and the expression of genes with $p = 0.01$ considered as significant. It is in the form of all of the experimental data that are in the form of mean values with the values of standard deviation obtained through three independent biological replications.

5. RESULTS

5.1 Differential Gene Expression Analysis

Multi-stress exposure was transcriptomically profiled to result in a greater transcriptional response than single stressors, heat or oxidative stress. As is demonstrated in Figure 1, the integrated heat and oxidative stress initiated signal perception pathways, and MAPK signaling pathways, ROS response pathways and hormonal pathways that in turn affected genomic and transcriptomic regulation. Multi-stress conditions were able to identify 1245 differentially expressed genes (DEGs) with $FDR < 0.05$, of which there were 732 upregulated and 513 down regulated genes. This implies that the multi-stress exposure created a widespread transcriptional reprogramming response and not a specific stress response.




Fig 1. Integrated Multi-Stress Epigenetic and Transcriptional Regulation Framework.

The key upregulated genes were HSP70, DREB2A, NAC1, HSF1, WRKY33, ZAT12, BZIP60 and ERF1. HSP70 was the most highly expressed with a log₂ fold change +3.21 indicating a robust heat-protective response. DREB2A was enhanced by +2.85, which implies that it activated the drought and stress response transcriptional regulation. The increase in NAC1 was +2.44, which indicated that it is a central stress-responsive transcription regulator. On the contrary, MYC2 and ABI1 were suppressed and their log₂ fold changes were -1.75 and -1.52 respectively. This implies the inhibition of the chosen hormonal and jasmonate-dependent regulatory pathways under joint exposure to stress. The key DEGs under multi-stress conditions are summarized in Table 1.

Table 1. Differentially Expressed Genes under Multi-Stress Conditions

Gene ID	Gene Name	Functional Role	log ₂ FC	p-value	Regulation
AT5G02490	HSP70	Heat shock protein; stress protection	+3.21	<0.001	Upregulated
AT5G05410	DREB2A	Drought-responsive transcription factor	+2.85	<0.005	Upregulated
AT1G56010	NAC1	Stress-responsive transcription regulator	+2.44	<0.005	Upregulated
AT1G32640	HSF1	Heat shock transcription factor	+2.18	<0.01	Upregulated
AT2G38470	WRKY33	Defense-related transcription factor	+1.96	<0.01	Upregulated
AT4G27410	ZAT12	Oxidative stress response protein	+1.89	<0.01	Upregulated
AT1G32660	BZIP60	Stress signaling regulator	+1.78	<0.01	Upregulated
AT2G36270	ERF1	Ethylene response factor	+1.64	<0.02	Upregulated
AT3G15540	ABI1	ABA signaling phosphatase	-1.52	<0.01	Downregulated
AT1G17380	MYC2	Jasmonate signaling regulator	-1.75	<0.01	Downregulated

5.2 Epigenetic Modification Profiles

The transcriptional activation in multi-stress conditions was highly correlated with promoter hypomethylation and activation of promoters with histone marks as shown by epigenetic profiling. As Figure 2 demonstrates, genes that are stress-responsive and upregulated exhibited more activation-related histone modifications, in particular, H3K4me3 and H3K27ac ones. The 1.8-fold and 2.1-fold increases were observed in these marks, respectively, when subjected to multi-stress. This pattern shows that, together with stress, an open and transcriptionally active chromatin configuration was favored. Analysis of DNA methylation revealed that the promoter regions of activated genes had average 28% reduction in their methylation levels, whereas downregulated genes displayed about 22% augmented in their methylation levels, particularly the body of the gene or regulatory elements. This methylation pattern implies that hypomethylation of promoters played a role in the activation of gene transcription and hypermethylation played a role in transcriptional repression. Downregulated regions also saw an increase in repressive histone modification. Mark H3K27me3 was up-regulated significantly (1.5-fold) and was enriched around genes including MYC2 and ABI1, which suggested the condensation of the chromatin and the decreased transcription. Figure 2 heatmap makes it evident that stress-response upregulated genes are distinctly separate of regulatory downregulated genes, demonstrating that multi-stress exposure yielded a coordinated-scale of epigenetics instead of arbitrary variability of gene expression. The relationship between the transcriptional outcomes and epigenetic markers is indicated in Table 2.

Table 2. Correlation between Epigenetic Modifications and Transcriptional Outcomes

Feature Category	Specific Marker	Multi-Stress Trend	Pearson r	Transcription Outcome
DNA methylation	Promoter methylation	Hypomethylation, avg. -28%	-0.68	Activation, e.g., HSP70
DNA methylation	Gene body methylation	Hypermethylation, avg. +22%	+0.45	Repression, e.g., MYC2
Histone mark	H3K4me3	1.8-fold increase	+0.65	High expression
Histone mark	H3K27ac	2.1-fold increase	+0.72	Active enhancers
Histone mark	H3K27me3	1.5-fold increase	-0.54	Gene silencing
Chromatin state	ATAC-seq peak	+35% openness	+0.79	TF accessibility




Fig 2. Epigenetic and Chromatin Remodelling Profiles Under Multi-Stress Conditions.

5.3 Chromatin Accessibility Changes

The analysis of chromatin accessibility showed that there was an increased regulation region openness of about 35% in comparison to the 18-22% openness on single-stress conditions. This means that synergistic stress led to a greater chromatin remodeling effect. Figure 1 demonstrates the change in chromatin accessibility panel between control state that is characterized by limited openness and multi-stress state characterized by greater binding capacity between transcription factors. The accessibility of genes around promoter and enhancer sites increased in genes like HSP70, DREB2A, and NAC1. This favors the explanation that chromatin remodeling allowed transcription factors to more effectively regulate stress-responsive loci. The enrichment of H3K27ac and H3K4me3 was also in line with the enhanced transcriptional plasticity mediated by chromatin accessibility and histone activation marks.

5.4 Integrated Multi-Omics Analysis

Combined transcriptomic, DNA methylation, histone modification, and chromatin accessibility data analysis revealed that there was a strong correlation between epigenetic remodeling and transcriptional output. Fig. 3 indicates that there is a negative correlation between promoter DNA methylation and expression of genes, so the Pearson $r = -0.68$, $p = 0.01$. This implies that the genes with less promoter methylation tended to be more expressed during multi-stress induced conditions. The most significant positive correlation was found between the

chromatin openness and transcription factor accessibility with $r = +0.79$. Histone activation mark, H3K27ac, was significantly positively correlated with the expression of genes, but H3K4me3 associates $r = +0.72$. These values demonstrate that transcriptional activation had active histone marks and open chromatin as significant contributors. On the other hand, H3K27me3 was negatively correlated with expression ($r = -0.54$), which agrees with its repressive nature. Figure 3 is an integrated module analysis that is divided into several genes under the activation and repression modules. Hypomethylation of promoters, enrichment of the promoters with the H3K27ac/H3K4me3, and elevated ATAC openness were the main features of the activation module which includes HSP70, DREB2A, NAC1, and HSF1. MYC2 and ABI1, which are linked to hypermethylation, H3K27me3 enrichment, and closed chromatin were included in the repression module.




Fig 3. Multi-Omics Correlation between Epigenetic Modifications and Gene Expression.

5.5 Key Regulatory Genes and Pathways

The pathway enrichment analysis revealed a number of key regulatory pathways of multi-stress adaptation. The most enriched pathway as presented in Table 3 is MAPK signaling whose enrichment score stands at 4.21 and $p = 0.001$. This pathway was linked to perception of stress and signal transduction especially by the regulatory genes DREB2A and HSF1. The ROS response pathway with an enrichment score of 3.87 and $p < 0.005$, included NAC1, HSF1 and ZAT12. This validates the fact that multi-stress response significantly involved oxidative stress defense mechanism. Hormonal signaling was also enriched significantly with a score of 3.45 and $p < 0.01$ including MYC2, NAC1 and ERF1 and this indicates regulation of trade-offs in growth-defense. Figure 4 shows the multi-stress integrated regulatory network. NAC1, DREB2A, Hsf1, WRKY33 are the important hub regulators identified in the network. These genes linked various stress-response systems, such as heat stress, redox regulation, combined adaptation and defense response. This figure also indicates that the multi-stress conditions enhanced the hub connectivity over single-stress conditions, which is an indicator of a synergistic regulatory model.




Fig 4. Integrated Multi-Stress Gene Regulatory Network and Pathway Interactions.

Table 3. Pathway Enrichment and Key Regulatory Hubs

Biological Pathway	Enrichment Score	p-value	Key Regulatory Genes	Functional Significance
MAPK signaling	4.21	<0.001	DREB2A, HSF1	Signal transduction and stress perception

ROS response	3.87	<0.005	NAC1, HSF1, ZAT12	Reactive oxygen species detoxification
Hormonal signaling	3.45	<0.01	MYC2, NAC1, ERF1	Growth-defense trade-off regulation
Protein folding	3.12	<0.01	HSP70, HSF1	Proteostasis under heat stress
Transcription reprogramming	2.98	<0.02	WRKY33, BZIP60	Large-scale genomic shifts

6. DISCUSSION

The current research offers a thorough examination of transcriptional plasticity in multi-stress environmental circumstances and sheds light on the critical role of epigenetic management in coordinating the adaptive responses. These findings show that exposure to combined stress is associated with a much more extensive and coordinated transcriptional response when compared to single-stress conditions. The discovery of 1,245 differentially expressed genes, and significant epigenetic remodelling indicate that transcriptional plasticity is not only a passive process but an active one under regulation by chromatin-scale changes and signal integration. One of the major discoveries of the research is that multi-stress reactions are not to be explained as the mere accumulation of the effects of the individual stress. Rather, the transcriptional pattern observed indicates the existence of two types of regulation that are synergistic and non-linear. The expression levels of the genes like HSP70, DREB2A and NAC1 were found to be much higher than that when the stress was exposed individually meaning that there was amplification of stress-response pathways. This effect is probably mediated by common signaling elements, including ROS and MAPK signatures, and correlated action of transcription factors. The increased chromatin accessibility (35% up-folding) and increase of activating histone marks also contribute to the fact that there is a regulatory regime that combines various environmental signals into a single transcriptional response. The other significant value brought to light by the present study is the contribution of the epigenetic memory in the development of adaptive responses. The noted trends of promoter hypomethylation and chronic enrichment of histone marks of activation indicates that previous stress can pre-tune the genome to respond more effectively to future stresses. This type of epigenetic memory has the potential to aid an adaptive benefit, namely faster and more resilient transcriptional activation in response to repeated or chronic stressors. It can be especially applicable to changing environments, where organisms need to continually adapt to changing stress combinations. The results of the research are not novel to the research conducted by other scientists in proving the significance of DNA methylation and histone modifications in stress responses, however, it adds to the current body of knowledge by highlighting the complexity of multi-stress interactions. Previous reports have mostly concentrated on individual stressors and their independent action on the expression of genes. Conversely, the present findings indicate that multi-stress states and conditions engage specific regulatory networks, which entail synergized epigenetic alterations and interactions among transcription factors. The fact that the relationship between chromatin accessibility and gene expression ($r = +0.79$) is strong further demonstrates the role of chromatin remodeling as a key tool of transcriptional control in complicated environmental conditions. In biological terms, the multi-stress signal integration, which results in the coordinated transcriptional response, is of critical importance in cellular homeostasis and survival. Key regulatory hubs such as DREB2A, NAC1 and HSF1 have been identified, highlighting the role of transcription factor networks in the responses. They are hubs that bring two signaling pathways together and promote the communication between environmental stimuli and gene expression machinery. Engagement of additional pathways like MAPK signaling, ROS detoxification and hormone regulation also reflect the interconnected nature of stress-response systems. Epigenetically-mediated transcriptional plasticity is an evolutionary force of its own, as it offers an effective means of fast adaptation without necessitating genetic changes. The reversible quality of epigenetic changes enables organisms to be dynamic to changes in the environment with maintenance of genomic stability. Such adaptive responses can also be evolutionary fitness over time by increasing resilience to environmental stressors. The capacity to display context-dependent patterns of gene expression in multi-stress situations can thus be a major determinant of survival and diversification of species during dynamic ecosystems. In sum, this paper sheds some new light regarding the molecular mechanism of transcriptional plasticity in that multi-stress conditions initiate orchestrated epigenetic and transcriptional responses which are intrinsically dissimilar to those produced by single-stresses. Such results help recognize the necessity to use integrative and multi-omics methods to comprehend properly the complexity of stress biology and provide new opportunities in environmental adaptation, agriculture, and biomedical research.

7. Applications

The results of this paper have far-reaching implications in various scientific and practical fields, especially the field of agriculture, biotechnology, and biomedical research. This study, through its show that multi-stress transcriptional plasticity is mediated by integrated epigenetic processes, forms the basis of developing strategies that can be used to increase stress resilience and adaptive capacity in complex environments. Among the most crucial applications is climate-resistant crop engineering. The climate change is subjecting agricultural systems to concurrent stressors like heat, drought, and oxidative stress. Key regulatory genes and epigenetic changes that are found to play a role in responses to multi-stresses such as the hypomethylation of promoters and epigenetic

activation-related histone marks provide useful crop improvement targets. In the case of HSP70 and DREB2A, a stress-responsive gene, the expression of either of these genes can be increased by editing these epigenetic processes using genetic engineering or epigenome editing genomic-editing systems like CRISPR/dCas9-based systems. This will result in the creation of crops that are more resistant to a variety of environmental stresses thus guaranteeing food security in changing climatic conditions.

The research also has significant implications to stress-adaptive biotechnology especially in the development of artificial biological systems and industrial microorganisms. Knowing the mechanisms of epigenetic control of gene expression under combined stress can allow the construction of engineered systems with the ability to stable performance under varying environmental conditions. As an example, microbial strains employed in the production of biofuels or in bioremediation can be engineered to resist several stressors by including regulatory circuits that are based on natural epigenetic control systems. These systems can change the expression of genes dynamically in response to environmental cues, enhancing performance and resilience in industry. Within the framework of biomedical modeling of stress-response, the findings are resourceful in terms of understanding the role of epigenetic regulation in cellular responses to the complex stress conditions. Epigenetic processes have also been implicated in dysregulation in many human diseases such as neurodegenerative disorders and metabolic syndromes as well as stress-related psychiatric diseases. The correlation between DNA methylation and histone changes and gene expression in multi-stress conditions can be used to develop predictive algorithms about disease progression and therapeutic response. Furthermore, it is possible that by attacking epigenetic regulators, new intervention opportunities can be presented in stress-related disorders by altering the patterns of gene expression related to the resilience and adaptation.

Lastly, the study is relevant to the evolution of environmental adaptation mechanisms especially in the ecological and systems biology. The discovery of network-based regulatory modules and feedback mechanisms furnish a paradigm through which organisms adaptation to complex and changing environments can be understood. Such understandings may be used in ecological modeling, conservation biology, and environmental monitoring systems where it is essential to forecast how organisms react to various stress factors. Moreover, more precise predictions can be made in terms of species survival, ecosystem stability, and biodiversity in stressful conditions by incorporating epigenetic data in environmental models. All in all, the scope of application of this study is beyond a basic research with practical recommendations that can be used to tackle real life problems related to environmental stress. This study offers a flexible avenue to enhance innovation in the agricultural, biotechnological, medical, and environmental science fields by connecting epigenetics regulation with transcriptional plasticity.

8. Limitations

Although this study offers valuable information regarding the epigenetic control of transcriptional plasticity during multi-stress condition, the study has a number of limitations that must be taken into account when interpreting the results. The main issue is that multi-stress modeling is intrinsically complex. Exposure to stress factors in nature can be in different intensities, durations and patterns but the experimental design uses constant and concomitant stress exposure. Though this method will provide reproducibility, it might not be as dynamic and heterogeneous as real-world environmental conditions, which can restrict the extrapolation of the results. The other major constraint is associated with integration of data across multi-omics platforms. Although transcriptomic, DNA methylation, histone modification, and chromatin accessibility datasets were combined in one analysis, it is conceptually difficult to combine the various types of data. Disagreements in data resolution, normalization methods, and analysis pipeline may bias the results, potentially influencing the analysis of correlations among epigenetic marks and gene expression. Despite finding strong associations, direct causal links between epigenetic alterations and transcriptional outcomes are only possible with additional experimental support. Temporal resolution is also a limitation to the study. The measurement was done at one or short term period after stress exposure and this limits the time dynamic changes of gene expression and epigenetic states to be measured over time. Transcriptional plasticity and epigenetic remodeling are quite dynamic and lack of time-course data constrains how these responses change, stabilize, or reverse during chronic or repeated exposure to stress.

Also, the experimental limitations of sample size, choice of model system, and controlled laboratory conditions can affect the results. A single well-characterized model organism might not be representative of the diversity of responses that are seen across species. Moreover, although biological replicates were used, an increase in sample size and validating the results independently in other systems would make the conclusions more robust. On the whole, these shortcomings demonstrate that future studies should include more complicated experimental designs, better methods of data integration, and longitudinal analyses to identify a more profound picture of epigenetic regulation in the multi-stress environmental conditions.

9. Future Directions

The results of this research lead to a number of prospective research directions that will help to deepen the knowledge of the epigenetic regulation of transcriptional plasticity in the conditions of multi-stresses. A significant new direction is the use of single-cell epigenomics that can reveal cell-to-cell heterogeneity in stress responses that cannot be found in bulk studies. Although the current research offers population-wide data, cells that make up a tissue can have different epigenetic patterns and transcriptional functions. The combination of

single-cell RNA sequencing with single-cell chromatin accessibility and methylation profiling would allow mapping regulatory mechanisms induced by stress in high-resolution and demonstrate heterogeneity in adaptive responses.

The development of artificial intelligence (AI)-based regulatory network modeling is another important direction. Multi-stress responses are more complex, non-linear interactions between signaling pathways, transcription factors, and epigenetic modifications, which can only be interpreted with advanced computational methods to influence the accuracy. Multi-omics datasets can be integrated with machine learning and deep learning models to make predictions of regulatory relationships, key hub genes, and simulate system wide responses to changes in combinations of stresses. These models would improve greatly the capability to anticipate transcriptional results and the guide intended interventions in agricultural and biomedical investigations. Long-term epigenetic memory and its contribution to adaptive resilience should also be researchable in the future. Though this paper offers indications of epigenetic remodeling with acute stress exposure, the maintenance and inheritance of these changes have been little studied. Studies that involve longitudinal data with time course analysis would be required to establish whether the stress-induced epigenetic alterations are sustained over time over the developmental period or be passed over generations. A deeper comprehension of the processes involved in epigenetic memory may help in explaining how organisms adapt to repeated environmental stressors and how organisms are able to retain increased levels of stress resistance during later growth.

Moreover, the creation of stress-response monitoring systems in real time is also a big step towards research and applied science. The combination of biosensors and omics technologies could allow ensuring continuous gene expression and epigenetic changes upon exposure to environmental stimuli. These systems would enable dynamic monitoring of transcriptional plasticity and enable quick identification of stress-induced changes in the molecules. The method has potential uses in precision agriculture, environmental monitoring, and personalized medicine, where real-time understanding of stress changes can be used to implement timely responses. In general, these future directions underline the importance of incorporating innovative experimental and computational methods to comprehensively clarify the mechanisms of transcriptional plasticity when faced with multi-stress factors. Future studies can create a more comprehensive and predictive picture of the importance of epigenetic regulation on adaptive responses within complex environments by integrating single-cell technologies, AI-enabled models, longitudinal research, and real-time tracking.

10. CONCLUSION

To summarize, the research shows that multi-stress environmental conditions have a highly coordinated epigenetic regulatory response that regulates transcriptional plasticity. Transcriptomic and epigenomic findings are integrated to show that dynamic fluctuations in methylations of DNA and histone levels as well as chromatin accessibility carefully regulate the expression of genes in complicated stress conditions. Multi-stress exposure can cause synergistic and non-linear regulatory actions in contrast to single-stress exposure, leading to increased stress-responsive gene expression and inhibition of non-essential pathways. The results emphasize the crucial role of epigenetic changes as the key regulators of adaptive gene expression and the value of multi-omics in the context of the discovery of these sophisticated processes. Altogether, the piece of work offers an important insight on the molecular nature of stress adaptation, and has critical implications in the creation of climate-resilient crops, stress-adaptive biotechnological systems, and enhanced knowledge of stress-related mechanisms in biomedical studies.

REFERENCES

1. Amirmahani, F., Kumar, S., & Muthukrishnan, S. D. (2025). Epigenetic mechanisms of plasticity and resistance in glioblastoma: therapeutic targets and implications. *Frontiers in Epigenetics and Epigenomics*, 3, 1519449.
2. Atlasi, Y., & Stunnenberg, H. G. (2017). The interplay of epigenetic marks during stem cell differentiation and development. *Nature Reviews Genetics*, 18(11), 643-658.
3. Duncan, E. J., Cunningham, C. B., & Dearden, P. K. (2022). Phenotypic plasticity: what has DNA methylation got to do with it?. *Insects*, 13(2), 110.
4. Feil, R., & Fraga, M. F. (2012). Epigenetics and the environment: emerging patterns and implications. *Nature reviews genetics*, 13(2), 97-109.
5. Hunter, R. G., Gray, J. D., & McEwen, B. S. (2018). The neuroscience of resilience. *Journal of the Society for Social Work and Research*, 9(2), 305-339.
6. Loughland, I., Little, A., & Seebacher, F. (2021). DNA methyltransferase 3a mediates developmental thermal plasticity. *Bmc Biology*, 19(1), 11.
7. Matosin, N., Cruceanu, C., & Binder, E. B. (2017). Preclinical and clinical evidence of DNA methylation changes in response to trauma and chronic stress. *Chronic stress*, 1, 2470547017710764.
8. Miller, M., McAfee, C., Brown, R. D., Sears, S., Krautkramer, C., Gogia, R., & Champagne, F. A. (2025). Impact of acute stress exposure on genome-wide DNA methylation. *Scientific Reports*, 15(1), 23931.
9. Sherov, U., Saliyeva, N., Anvarova, M., Jurayev, Z., Salomova, F., Iskandarov, S., & Adilova, S. (2026). Exploring Epigenetic Responses to Occupational Stress in Tourism Workers and Implications for Mental Health. *Genetics and Molecular Research*, 25(1), 1-11.
10. Szyf, M., Tang, Y. Y., Hill, K. G., & Musci, R. (2016). The dynamic epigenome and its implications for behavioral interventions: a role for epigenetics to inform disorder prevention and health promotion. *Translational behavioral medicine*, 6(1), 55-62.

11. Tan, U. (2026). Plant Stress Memory: Epigenetic and Mitotic Mechanisms Underlying Stress Adaptation in Plants. Polat, T. & Baytin Alaci, R. *Multidisciplinary Approaches in Agricultural and Aquatic Sciences*, 35-69.
12. Zhang, T. Y., & Meaney, M. J. (2010). Epigenetics and the environmental regulation of the genome and its function. *Annual review of psychology*, 61(1), 439-466.