

ROLE OF BIOLOGICAL HARDENING IN ENHANCING THE SURVIVAL OF *IN VITRO* RAISED POTATO PLANTLETS

Lokesh T H^{1*}, Vidya A², Shalini M^{3*}, Suneetha C⁴ and Ashika S⁵

^{1*} Ph.D. Scholar, Department of Horticulture, College of Agriculture, University of Agricultural Sciences, Gandhi Krishi Vigyana Kendra, Bengaluru 560 065, Karnataka, India

² Professor, Department of Horticulture, College of Agriculture, University of Agricultural Sciences, Gandhi Krishi Vigyana Kendra, Bengaluru 560 065, Karnataka, India

^{3*} Associate Professor, Department of Horticulture, College of Agriculture, University of Agricultural Sciences, Gandhi Krishi Vigyana Kendra, Bengaluru 560 065, Karnataka, India

⁴ Associate Professor, Department of Horticulture, College of Agriculture, University of Agricultural Sciences, Gandhi Krishi Vigyana Kendra, Bengaluru 560 065, Karnataka, India

⁵ Ph.D. Scholar, Department of Horticulture, College of Agriculture, University of Agricultural Sciences, Gandhi Krishi Vigyana Kendra, Bengaluru 560 065, Karnataka, India

*Correspondence Author: Lokesh T H; Email: lokeshth1999@gmail.com

*Correspondence Author: Shalini M; Email: shalinigowda1978@gmail.com

ABSTRACT

The experiment entitled “Role of biological hardening in enhancing the survival of *in vitro* raised potato plantlets” was conducted at the Plant Tissue Culture Laboratory, Department of Horticulture, University of Agricultural Sciences, GKVK, Bengaluru, Karnataka, during 2025 to 2026. Biological hardening is an important step in potato micropropagation, as plantlets are highly susceptible to moisture stress and environmental changes during transfer from *in vitro* to *ex vitro* conditions. The study aimed to evaluate the effectiveness of beneficial microbial bioagents, applied individually and in combination with glycerol, in improving the survival and growth of micropropagated potato plantlets of cv. Kufri Himalini. The bioagents used were *Pseudomonas fluorescens*, *Bacillus subtilis* and *Trichoderma viride*, along with 0.5 per cent glycerol as an anti transpirant. Plant survival was recorded during the first, second and third weeks, while plant height, internodal length and root length were assessed initially and after three weeks. Among the treatments, T₁₀ (*P. fluorescens* + *T. viride* + 0.5 % glycerol) performed best, recording 83.33 per cent survival, 19.03 cm plant height, 2.05 cm internodal length and 10.83 cm root length after three weeks. The second-best treatment, T₅ (*P. fluorescens* + *T. viride*), recorded 80.00 per cent survival, 18.17 cm plant height, 2.02 cm internodal length and 10.07 cm root length. The third-best treatment, T₆ (*T. viride* + 0.5% glycerol), recorded 76.67 per cent survival, 17.13 cm plant height, 2.01 cm internodal length and 9.53 cm root length after three weeks. In contrast, the control recorded the lowest survival (56.67 %) and plant height (14.47 cm). The findings indicate that the combined application of *P. fluorescens*, *T. viride* and glycerol is a promising approach for improving the acclimatization, survival and establishment of micropropagated potato plantlets.

KEYWORDS: Biological hardening, potato, micropropagation, *Pseudomonas fluorescens*, *Trichoderma viride*, glycerol, acclimatization.

INTRODUCTION

Potato (*Solanum tuberosum* L.), a natural autotetraploid (2n=4x=48) belonging to the family Solanaceae, is the fourth most important food crop in the world after wheat, rice and maize. It originated in tropical South America and exhibits considerable genetic diversity. Potato is propagated both sexually through true potato seeds and asexually through tubers, with seed tubers being the most widely used planting material. However, conventional tuber-based propagation has several limitations, including low multiplication rates and a high susceptibility to diseases (Mirza *et al.*, 2001).

Micropropagation offers an effective biotechnological approach for producing disease-free, genetically uniform potato plantlets through clonal propagation, thereby facilitating the large-scale production of healthy planting material and microtubers. Despite its advantages, micropropagation presents challenges during the transition from *in vitro* to *ex vitro* conditions, as plantlets are highly sensitive to abiotic stresses, such as temperature, light intensity and humidity fluctuations, as well as biotic stresses caused by soil microorganisms. These stresses can result in considerable plantlet losses during acclimatization (Nguyen *et al.*, 2016). Therefore, acclimatization is a critical stage that requires the gradual adaptation of *in vitro* plantlets to greenhouse and field conditions. Owing to their poorly developed protective mechanisms and high dependence on humidity, *in vitro* plantlets are susceptible to rapid water loss and wilting. The application of anti-transpirants, such as glycerol, abscisic acid (ABA), paraffin and grease, may help reduce transpiration and improve plantlet survival. Furthermore, pre-hardening strategies, including adjustment of sucrose concentration, application of desiccants and improvement of ventilation, can enhance plantlet establishment under *ex vitro* conditions (El-Din *et al.*, 2020).

Biological hardening is a promising strategy for reducing acclimatization stress by introducing beneficial microorganisms into the growing substrate. Since *in vitro*-raised plantlets are generally maintained under aseptic conditions and are physiologically delicate, the introduction of beneficial rhizospheric or endophytic microorganisms, such as *Pseudomonas* and *Bacillus*, can promote root development, plant growth and tolerance to environmental stresses (Mirza *et al.*, 2001;

Lugtenburg and Kamilova, 2009). These beneficial bacteria possess several plant growth-promoting traits, including 1-aminocyclopropane-1-carboxylate (ACC) deaminase activity, nitrogen fixation, phosphate solubilization and the production of indole-3-acetic acid (IAA), which plays an important role in regulating plant growth and development (Jimtha *et al.*, 2014; Zhao *et al.*, 2011). In addition, certain bacterial strains exhibit antagonistic activity against plant pathogens, thereby contributing to improved plant health and resilience (Müller *et al.*, 2018). Fungal bioagents, particularly *Trichoderma* spp., also serve as effective biocontrol agents and biofertilizers by enhancing root colonization, nutrient uptake, plant growth and productivity (Vinale *et al.*, 2008).

In the present study, different combinations of biological hardening agents, with and without the addition of 0.5% glycerol, were evaluated to determine their effects on the survival of *in vitro*-raised potato plantlets during acclimatization. The study aimed to identify the most effective combination of biological hardening agents and glycerol for improving plantlet survival and successful establishment under *ex vitro* conditions.

MATERIAL AND METHODS

Experimental Location

The present investigation entitled “Role of biological hardening in enhancing the survival of *in vitro* raised potato plantlets” was conducted at the Plant Tissue Culture Laboratory, Department of Horticulture, University of Agricultural Sciences, GKVK (Gandhi Krishi Vigyan Kendra), Bengaluru-560 065, Karnataka, India, during 2025 to 2026.

Plant Material and Explant Source

Sprouts and nodal segments of the medium-maturing, late blight-resistant potato cultivar ‘Kufri Himalini’ were used as explants. Sprouts were excised from healthy, disease-free tubers, while single-node segments were collected from pot-grown plants after removing the leaves. The explants were initially washed thoroughly with tap water to remove dirt and debris. A few drops of Tween-20 were added, and the explants were washed under running tap water for 15 to 20 minutes before surface sterilization.

Micropropagation of Plantlets

The explants were surface-sterilized using 2 per cent sodium hypochlorite for 5 minutes and subsequently rinsed 4 to 5 times with sterile distilled water to remove residual sterilant and prevent tissue damage. Shoot regeneration was carried out on Murashige and Skoog (MS) medium supplemented with 30 g sucrose, 6 g agar and different plant-derived extracts like coconut water, potato homogenate, banana homogenate and tomato homogenate at 10, 15 and 20 per cent concentration. The sterilized sprouts and nodal segment explants were inoculated onto the regeneration medium, maintaining the correct polarity of the explants. The cultures were incubated at $25 \pm 2^\circ\text{C}$ under a photoperiod of 16 hours of light and 8 hours of darkness, with a light intensity of 3,000 to 3,500 lux. Regenerated shoots measuring 5 to 6 cm in length were transferred to rooting medium containing half-strength MS medium with different plant-derived extracts at 10, 15 and 20 per cent. Before biological hardening, plantlets with well-developed adventitious roots were pre-hardened by removing from the culture vessels and maintaining them in the culture room for one week. This procedure was intended to promote leaf adaptation to lower humidity and water stress through stomatal regulation. The rooted plantlets were subsequently used for the biological hardening experiments (Fig. 1).

Preparation of Bioagents

To enhance the survivability of *in vitro*-raised potato plantlets during the hardening phase, beneficial microbial bioagents, namely *Pseudomonas fluorescens*, *Bacillus subtilis* and *Trichoderma viride*, were used individually and in different combinations. Pure cultures of these bioagents were procured from the Department of Microbiology, University of Agricultural Sciences (UAS), GKVK, Bengaluru. For the preparation of bacterial bioagents, a loopful of inoculum from each pure culture was inoculated separately into 100 mL of nutrient broth in culture flasks. The inoculated flasks were incubated at $28 \pm 2^\circ\text{C}$ for 3 to 5 days. Similarly, the fungal bioagent *T. viride* was cultured by inoculating a loopful of fungal inoculum into 100 mL of potato dextrose broth (PDB), followed by incubation at $28 \pm 2^\circ\text{C}$ for 3 to 5 days. Before application in the biological hardening treatments, the culture broths of the individual bioagents and their combinations were filtered through filter paper. The resulting filtrates were used for root treatment of *in vitro*-raised potato plantlets during *ex vitro* acclimatization. The different bioagents used in the study and glycerol at a concentration of 0.5 per cent was used as an anti transpirant to reduce transpiration losses and improve the survival of *in vitro*-raised plantlets during the hardening process presented in Table 1 and Fig. 2.

Ex Vitro Acclimatization

Well-rooted *in vitro*-raised potato plantlets were carefully removed from the culture vessels using forceps to avoid mechanical damage. The agar medium adhering to the roots was gently washed off under running tap water to remove residual medium and minimize the risk of contamination. The roots were then dipped in the respective bioagent culture broths for 2 to 3 minutes, according to the treatment. Anti transpirant treatment was provided using 0.5 per cent glycerol, in which the shoot portions, including the stems and leaves, were dipped for five minutes. The treatments were applied according to the experimental treatments, with or without prior bioagent application. Immediately after treatment, the plantlets were transferred to protrays containing sterile cocopeat and maintained in a growth chamber for approximately 15 days for primary hardening. Humidity was maintained by regularly spraying water inside the covering bags to minimize moisture loss and reduce transplanting stress. After primary hardening, the established plantlets were transplanted into pots containing a potting mixture of farmyard manure (FYM), sand and soil in a 1:1:1 ratio for further establishment. Observations on plant survival were recorded at the end of the first, second and third weeks of acclimatization. Plant height, internodal length and root length were measured initially, at the time of bioagent root dipping, and again after three weeks of acclimatization to assess plant growth and establishment under *ex vitro* conditions.

Experimental Design and Statistical Analysis

The *in vitro* microtuberization experiment was conducted in a completely randomized design (CRD) with five replicates and were used to evaluate all of the data. The significance of treatment effects was assessed using the F-test at the 1 per cent level of significance. Statistical analysis was carried out using OPSTAT online software (Sekhar *et al.*, 2019).

RESULTS

Effect of different biological hardening treatments on survival of *in vitro* raised plantlets

The effect of different bio-agents, applied individually and in combination with glycerol (0.5 %), on the survival of *in vitro* raised potato plantlets under partially controlled conditions was evaluated after the 1st, 2nd and 3rd weeks of transplanting, as presented in Table 2. The bio-agents were applied as a root dip for 3 minutes, while glycerol (0.5 %) was applied as a shoot and leaf dip for 5 minutes. The observations on plantlet survival were recorded at each interval to assess the effectiveness of the different biological hardening treatments. The results revealed that T₁₀ (*Pseudomonas fluorescens* + *Trichoderma viride* + glycerol 0.5 %) recorded the highest plant survival of 93.33, 86.67 and 83.33 per cent after the 1st, 2nd and 3rd week of transplanting, respectively. This was followed by T₅ (*Pseudomonas fluorescens* + *Trichoderma viride*), which recorded 90.00, 83.33 and 80.00 per cent survival, respectively. The third-best treatment was T₆ (*Trichoderma viride* + glycerol 0.5 %), with 86.67, 80.00 and 76.67 per cent survival at the respective intervals. In contrast, the lowest survival was recorded in T₁₁ (control), with 63.33, 60.00 and 56.67 per cent, respectively. The superior survival under T₁₀ may be attributed to the synergistic effect of *Pseudomonas* and *Trichoderma*, along with glycerol-mediated reduction of transplanting stress and improved acclimatization.

Effect of different biological hardening treatments on plant height of *in vitro* raised plantlets

Significant differences were observed among the treatments with respect to plant height, both initially and after 3 weeks of transplanting (Table 3). Initially, the highest plant height was recorded in T₈ (*Bacillus subtilis* + glycerol 0.5 %) at 10.80 cm, followed by T₅ (*Pseudomonas fluorescens* + *Trichoderma viride*) and T₇ (*Pseudomonas fluorescens* + glycerol 0.5 %), both recording 10.70 cm. After 3 weeks, T₁₀ (*Pseudomonas fluorescens* + *Trichoderma viride* + glycerol 0.5 %) recorded the highest plant height (19.03 cm), followed by T₅ (18.17 cm) and T₆ (*Trichoderma viride* + glycerol 0.5 %) (17.13 cm) (Fig. 3). The lowest plant height after 3 weeks was recorded in T₁₁ (control) at 14.47 cm. The superior performance of T₁₀ may be attributed to the synergistic effects of *Pseudomonas* and *Trichoderma*, along with glycerol, which may enhance nutrient availability, root development and plantlet acclimatization.

Effect of different biological hardening treatments on internodal length of *in vitro* raised plantlets

Significant differences were also observed among the treatments for internodal length, both initially and after 3 weeks of transplanting (Table 4). Initially, the maximum internodal length was recorded in T₁ (*Trichoderma viride*) at 1.67 cm, followed by T₂ (*Pseudomonas fluorescens*) and T₈ (*Bacillus subtilis* + glycerol 0.5 %), both recording 1.63 cm. After 3 weeks, T₁₀ (*Pseudomonas fluorescens* + *Trichoderma viride* + glycerol 0.5 %) recorded the highest internodal length (2.05 cm), followed by T₅ (*Pseudomonas fluorescens* + *Trichoderma viride*) (2.02 cm) and T₆ (*Trichoderma viride* + glycerol 0.5 %) (2.01 cm). The lowest internodal length was recorded in T₁₁ (control) at 1.83 cm. The increased internodal length under T₁₀ may be associated with the combined growth-promoting effects of *Pseudomonas*, *Trichoderma* and glycerol, which supported better plantlet growth during acclimatization.

Effect of different biological hardening treatments on root length of *in vitro* raised plantlets

Root length was significantly influenced by the different bio-agents, applied alone or in combination with glycerol (0.5 %) (Table 5). Initially, the highest root length was recorded in T₁₀ (*Pseudomonas fluorescens* + *Trichoderma viride* + glycerol 0.5 %) at 7.27 cm, followed by T₅ (*Pseudomonas fluorescens* + *Trichoderma viride*) at 7.17 cm and T₆ (*Trichoderma viride* + glycerol 0.5 %) at 7.10 cm. After 3 weeks of transplanting, T₁₀ maintained the highest root length (10.83 cm), followed by T₅ (10.07 cm) and T₆ (9.53 cm). The lowest root length was recorded in T₁₁ (control), with 6.27 cm initially and 7.67 cm after 3 weeks. The enhanced root development under T₁₀ may be attributed to the synergistic action of *Pseudomonas* and *Trichoderma*, along with glycerol, which may promote root growth, nutrient acquisition and better acclimatization of plantlets.

DISCUSSION

The experiment was conducted to evaluate the effect of different biological hardening treatments on the acclimatization and establishment of micropropagated potato plantlets. Acclimatization is a critical stage in micropropagation because plantlets are sensitive to environmental stresses during transfer from *in vitro* to *ex vitro* conditions (Hazarika *et al.*, 2006). In the present study, plant survival gradually decreased from the first to the third week across all treatments. T₁₀ (*Pseudomonas fluorescens* + *Trichoderma viride* + 0.5 % glycerol) recorded the highest survival rates of 93.33, 86.67 and 83.33 per cent during the first, second and third weeks, respectively, whereas the control recorded the lowest survival rates of 63.33, 60.00 and 56.67 per cent. The improved survival under T₁₀ may be attributed to the complementary effects of the bioagents and the anti transpirant action of glycerol in reducing transplanting stress and excessive water loss (Gawad, 2015). Similar beneficial effects of bioagents and glycerol have been reported by Trivedi and Pandey (2007), Cheng *et al.* (2008), Bhushan and Gupta (2010), Santos *et al.* (2010), Anith *et al.* (2011) and Gupta *et al.* (2017). Plant height increased in all treatments during the three-week acclimatization period. The maximum plant height was recorded in T₁₀ (19.03 cm), followed by T₅ (*P. fluorescens* + *T. viride*) at 18.17 cm and T₆ (*T. viride* + 0.5 % glycerol) at 17.13 cm, while the control recorded the minimum height of 14.47 cm. In T₁₀, plant height increased from 9.90 to 19.03 cm. The superior growth may be associated with improved nutrient availability, root development and plant-growth-promoting activities of the bioagents. The growth-promoting effects of *Trichoderma* species have been associated with

the production or modulation of auxins, gibberellins and other physiological compounds (Montealegre *et al.*, 2010; Druzhinina *et al.*, 2011; Brotman *et al.*, 2013). Similar beneficial effects of microbial treatments on micropropagated potato and banana have been reported by Lavanya *et al.* (2009), Lincy and Sasikumar (2010), Sparta and Emilda (2020) and Naik *et al.* (2025).

Internodal length also increased during acclimatization, with the maximum value recorded in T₁₀ (2.05 cm), followed by T₅ (2.02 cm) and T₆ (2.01 cm). The control recorded the minimum internodal length of 1.83 cm. The increase in internodal length in T₁₀ from 1.64 to 2.05 cm suggests improved shoot elongation under the combined treatment. This response may be related to the plant-growth-promoting effects of *P. fluorescens* and *T. viride*, including improved nutrient uptake and the possible modulation of growth-regulating compounds such as auxins and gibberellins (Harman *et al.*, 2004; Druzhinina *et al.*, 2011). These findings indicate that the combined application of bioagents and glycerol may support vegetative growth during the hardening process.

Root development was also enhanced by the biological hardening treatments. T₁₀ recorded the maximum root length of 10.83 cm, followed by T₅ (10.07 cm) and T₆ (9.53 cm), whereas the control recorded the minimum root length of 7.67 cm. In T₁₀, root length increased from 7.27 to 10.83 cm. The improved root growth may be attributed to the complementary effects of *P. fluorescens* and *T. viride* in promoting root development and nutrient uptake, while glycerol may help reduce moisture stress during acclimatization. Similar beneficial effects of microbial inoculation on plant growth and root development have been reported in *Azadirachta indica* (Lavanya *et al.*, 2009), ginger (Lincy and Sasikumar, 2010), banana (Sparta and Emilda, 2020) and potato (Naik *et al.*, 2025). Overall, T₁₀ was the most effective treatment, recording the highest survival, plant height, internodal length and root length after three weeks, demonstrating its potential for improving the acclimatization and establishment of micropropagated potato plantlets.

CONCLUSION

Overall, T₁₀ (*P. fluorescens* + *T. viride* + 0.5 % glycerol) was the best-performing treatment across all four parameters after three weeks of acclimatization, recording the highest plant survival (83.33 %), plant height (19.03 cm), internodal length (2.05 cm) and root length (10.83 cm). It was followed by T₅ (*P. fluorescens* + *T. viride*), while T₆ (*T. viride* + 0.5 % glycerol) also showed comparatively favourable results. The control consistently recorded lower values for survival and growth parameters. The superior performance of T₁₀ suggests that the combined application of *P. fluorescens*, *T. viride* and glycerol may have contributed to improved acclimatization, root establishment and vegetative growth through complementary microbial and anti transpirant effects. Thus, the integration of beneficial bioagents with glycerol appears to be a promising approach for improving the survival and establishment of micropropagated potato plantlets during *ex vitro* hardening.

REFERENCES

1. Anith KN, Faseela KM, Archana PA and Prathapan KD (2011). Compatibility of *Piriformospora indica* and *Trichoderma harzianum* dual inoculants in black pepper (*Piper nigrum* L.). *Symbiosis*, 55(1): 11-17.
2. Bhushan A and Gupta RK (2010). Adventitious shoot regeneration in different explants of six genotypes of tomato. *Indian Journal of Horticulture*, 67(4): 224-227.
3. Brotman Y, Landau U, Cuadros-Inostroza Á, Takayuki T, Fernie AR, Chet I, Viterbo A and Willmitzer L (2013). *Trichoderma*-plant root colonization: Escaping early plant defense responses and activation of the antioxidant machinery for saline stress tolerance. *PLoS Pathogens*, 9(3): 1003221.
4. Cheng Y, Dengsha B, Sun L, Feldmann F and Gu F (2008). Utilization of arbuscular mycorrhizal fungi during production of micropropagated potato (*Solanum tuberosum*). *Mycorrhiza Works*, pp. 47-59.
5. Druzhinina IS, Seidl-Seiboth V, Herrera-Estrella A, Horwitz BA, Kenerley CM, Monte E, Mukherjee PK, Zeilinger S, Grigoriev IV and Kubicek CP (2011). *Trichoderma*: The genomics of opportunistic success. *Nature Reviews Microbiology*, 9(10): 749-759.
6. El-Din WMS and Mokhtar NAYO (2020). Effect of irrigation scheduling and some anti transpirants on water relations and productivity of *Mentha viridis* L. *American-Eurasian Journal of Agricultural and Environmental Sciences*, 20: 367-390.
7. Gawad NMA (2015). Effect of some anti transpirants to reduce amount of irrigation water added to banana cv. 'Grand Nain' in sandy soil. *Egyptian Journal of Horticulture*, 42(1): 69-86.
8. Gupta RK, Verma VS, Bhushan A and Raina V (2017). Biological hardening of micropropagated tomato plantlets: A case study with *Piriformospora indica*. In: Verma V and Gange AC (eds.). *Mycorrhiza: Nutrient Uptake, Biocontrol, Ecorestoration*. Springer International Publishing, Cham, pp. 301-311.
9. Harman GE, Howell CR and Viterbo A (2004). *Trichoderma* species-opportunistic, avirulent plant symbionts. *Nature Reviews Microbiology*, 2: 43-56.
10. Hazarika BN, Teixeira da Silva JA and Talukdar A (2006). Effective acclimatization of *in vitro*-cultured plants: Methods, physiology and genetics. In: *Floriculture, Ornamental and Plant Biotechnology*, 2: 427-438.
11. Jimtha JC, Smitha PV, Anisha C, Deepthi T, Meekha G, Radhakrishnan EK, Gayatri GP and Remakanthan A (2014). Isolation of endophytic bacteria from embryogenic suspension culture of banana and assessment of their plant growth-promoting properties. *Plant Cell, Tissue and Organ Culture*, 118: 57-66.
12. Lavanya M, Venkateshwarlu B and Devi BP (2009). Acclimatization of neem micro shoots adaptable to semi-sterile conditions. *Indian Journal of Biotechnology*, 8: 218-222.

13. Lincy A and Sasikumar B (2010). Enhanced adventitious shoot regeneration from aerial stem explants of ginger using TDZ and its histological studies. *Turkish Journal of Botany*, 34(1): 21-29.
14. Lugtenberg B and Kamilova F (2009). Plant-growth-promoting rhizobacteria. *Annual Review of Microbiology*, 63: 541-556.
15. Mirza MS, Ahmad W, Latif F, Haurat J, Bally R, Normand P and Mallik KA (2001). Isolation, partial characterization, and the effect of plant growth-promoting bacteria (PGPB) on micro-propagated sugarcane *in vitro*. *Plant and Soil*, 237: 47-54.
16. Montealegre J, Valderrama L, Sánchez S, Herrera R, Besoain X and Pérez LM (2010). Biological control of *Rhizoctonia solani* in tomatoes with *Trichoderma harzianum* mutants. *Electronic Journal of Biotechnology*, 13: 1-2.
17. Müller T, Ruppel S, Behrendt U, Lentzsch P and Müller MEH (2018). Antagonistic potential of fluorescent pseudomonads colonizing wheat heads against mycotoxin-producing *Alternaria* and *Fusaria*. *Frontiers in Microbiology*, 9: 2124.
18. Naik R, Bhushan A, Singh B, Kumar S, Sharma S, Kumar R and Nishant (2025). Enhancing acclimatization of *in vitro*-raised potato seedlings via biological hardening. *Indian Journal of Horticulture*, pp. 291-296.
19. Nguyen QT, Xiao Y and Kozai T (2016). Photoautotrophic micropropagation. In: Kozai T, Niu G and Takagaki M (eds.). *Academic Press*, Burlington, pp. 271-283.
20. Santos RD, Girardi CG, Pescador R and Stürmer SL (2010). Effects of arbuscular mycorrhizal fungi and phosphorus fertilization on post-*in vitro* growth of micropropagated *Zingiber officinale* Roscoe. *Revista Brasileira de Ciência do Solo*, 34: 765-771.
21. Sekhar V, Umakrishna K, Rao VS and Satyani C (2019). Completely randomised design (CRD) analysis-by manual and MS-Excel. *International Journal of Current Microbiology and Applied Sciences*, 8(3): 954-959.
22. Sparta A and Emilda D (2020). Growth evaluation of banana cv. Barangan as the effect of *Trichoderma* sp. and covering types during acclimatization process. *Caraka Tani: Journal of Sustainable Agriculture*, 35(2): 268-277.
23. Trivedi P and Pandey A (2007). Biological hardening of micropropagated *Picrorhiza kurroa* Royle ex Benth., an endangered species of medical importance. *World Journal of Microbiology and Biotechnology*, 23(6): 877-878.
24. Vinale F, Sivasithamparam K, Ghisalberti EL, Marra R, Woo SL and Lorito M (2008). *Trichoderma* plant pathogen interactions. *Soil Biology and Biochemistry*, 40(1): 1-10.
25. Zhao Q, Shen W, Xiao T, Xu D and Xu Y (2011). Inoculation of soil by *Bacillus subtilis* Y-IVI improves plant growth and colonization of the rhizosphere and interior tissues of muskmelon (*Cucumis melo* L.). *Biology and Fertility of Soils*, 47(5): 507-514.

Fig. 1. The rooted plantlets were subsequently used for the biological hardening experiments



Fig. 2. Biological hardening of *in vitro* raised potato plantlets using bio-agents: (a) *Pseudomonas fluorescens*, (b) *Bacillus subtilis*, (c) *Trichoderma viride*, (d) *Pseudomonas fluorescens* + *Trichoderma viride* + Glycerol (0.5 %), (e) Glycerol (0.5 %) - root dip, and (f) Glycerol (0.5 %) - shoot dip

Fig. 3. Effect of bio-agents on plant height after 3 weeks of hardening: (a) Control, (b) *Trichoderma viride* + Glycerol (0.5 %), (c) *Pseudomonas fluorescens* + *Trichoderma viride* + Glycerol (0.5 %), and (d) *Pseudomonas*



fluorescens + *Trichoderma viride*

Table 1. Effect of biological hardening treatments on *in vitro* raised potato plantlets

Treatments	Combinations
T ₁	<i>Trichoderma viride</i>
T ₂	<i>Pseudomonas fluorescens</i>
T ₃	<i>Bacillus subtilis</i>
T ₄	<i>Pseudomonas fluorescens</i> + <i>Bacillus subtilis</i>
T ₅	<i>Pseudomonas fluorescens</i> + <i>Trichoderma viride</i>
T ₆	<i>Trichoderma viride</i> + Glycerol (0.5 %)
T ₇	<i>Pseudomonas fluorescens</i> + Glycerol (0.5 %)
T ₈	<i>Bacillus subtilis</i> + Glycerol (0.5 %)
T ₉	<i>Pseudomonas fluorescens</i> + <i>Bacillus subtilis</i> + Glycerol (0.5 %)
T ₁₀	<i>Pseudomonas fluorescens</i> + <i>Trichoderma viride</i> + Glycerol (0.5 %)
T ₁₁	Control (seedling dip in distilled water)

Table 2. Effect of bio-agents on the survival percentage of *in vitro* raised potato plantlets

Treatments		Internodal length (cm)	
		Initial length	After 3 weeks
T ₁	<i>Trichoderma viride</i>	1.67	1.96
T ₂	<i>Pseudomonas fluorescens</i>	1.63	1.90
T ₃	<i>Bacillus subtilis</i>	1.60	1.85
T ₄	<i>Pseudomonas fluorescens</i> + <i>Bacillus subtilis</i>	1.47	1.93
T ₅	<i>Pseudomonas fluorescens</i> + <i>Trichoderma viride</i>	1.60	2.02
T ₆	<i>Trichoderma viride</i> + Glycerol (0.5%)	1.57	2.01
T ₇	<i>Pseudomonas fluorescens</i> + Glycerol (0.5%)	1.57	1.91
T ₈	<i>Bacillus subtilis</i> + Glycerol (0.5%)	1.63	1.87
T ₉	<i>Pseudomonas fluorescens</i> + <i>Bacillus subtilis</i> + Glycerol (0.5%)	1.57	1.94
T ₁₀	<i>Pseudomonas fluorescens</i> + <i>Trichoderma viride</i> + Glycerol (0.5%)	1.64	2.05
T ₁₁	Control (seedling dip in distilled water)	1.53	1.83
F-Test		NS	**
SEm ±		0.08	0.03
CD at 1 %		0.34	0.11

Table 3. Effect of bio-agents on the plant height of *in vitro* raised potato plantlets

Treatments		Plant survival rate (%)		
		After 1 st week	After 2 nd week	After 3 rd week
T ₁	<i>Trichoderma viride</i>	83.33	76.67	70.00
T ₂	<i>Pseudomonas fluorescens</i>	73.33	70.00	66.67
T ₃	<i>Bacillus subtilis</i>	66.67	63.33	60.00
T ₄	<i>Pseudomonas fluorescens</i> + <i>Bacillus subtilis</i>	80.00	73.33	70.00
T ₅	<i>Pseudomonas fluorescens</i> + <i>Trichoderma viride</i>	90.00	83.33	80.00
T ₆	<i>Trichoderma viride</i> + Glycerol (0.5 %)	86.67	80.00	76.67
T ₇	<i>Pseudomonas fluorescens</i> + Glycerol (0.5 %)	76.67	73.33	70.00
T ₈	<i>Bacillus subtilis</i> + Glycerol (0.5 %)	73.33	70.00	63.33
T ₉	<i>Pseudomonas fluorescens</i> + <i>Bacillus subtilis</i> + Glycerol (0.5 %)	80.00	73.33	66.67
T ₁₀	<i>Pseudomonas fluorescens</i> + <i>Trichoderma viride</i> + Glycerol (0.5 %)	93.33	86.67	83.33
T ₁₁	Control (seedling dip in distilled water)	63.33	60.00	56.67
F-Test		**	**	**
SEm ±		0.28	0.27	0.25
CD at 1 %		1.13	1.06	0.98

Table 4. Effect of bio-gents on the internodal length of *in vitro* raised potato plantlets

Treatments		Plant height (cm)	
		Initial length	After 3 weeks
T ₁	<i>Trichoderma viride</i>	9.70	16.97
T ₂	<i>Pseudomonas fluorescens</i>	10.40	15.67
T ₃	<i>Bacillus subtilis</i>	10.30	15.07
T ₄	<i>Pseudomonas fluorescens</i> + <i>Bacillus subtilis</i>	10.50	16.17
T ₅	<i>Pseudomonas fluorescens</i> + <i>Trichoderma viride</i>	10.70	18.17
T ₆	<i>Trichoderma viride</i> + Glycerol (0.5%)	10.00	17.13
T ₇	<i>Pseudomonas fluorescens</i> + Glycerol (0.5%)	10.70	16.00
T ₈	<i>Bacillus subtilis</i> + Glycerol (0.5%)	10.80	15.17
T ₉	<i>Pseudomonas fluorescens</i> + <i>Bacillus subtilis</i> + Glycerol (0.5%)	10.30	16.80
T ₁₀	<i>Pseudomonas fluorescens</i> + <i>Trichoderma viride</i> + Glycerol (0.5%)	9.90	19.03
T ₁₁	Control (seedling dip in distilled water)	10.10	14.47
F-Test		**	**
SEm ±		0.13	0.74
CD at 1 %		0.53	2.96

Table 5. Effect of bio-gents on the root length of *in vitro* raised potato plantlets

Treatments		Root length (cm)	
		Initial length	After 3 weeks
T ₁	<i>Trichoderma viride</i>	6.83	9.33
T ₂	<i>Pseudomonas fluorescens</i>	6.53	8.53
T ₃	<i>Bacillus subtilis</i>	6.33	8.03
T ₄	<i>Pseudomonas fluorescens</i> + <i>Bacillus subtilis</i>	6.83	9.07
T ₅	<i>Pseudomonas fluorescens</i> + <i>Trichoderma viride</i>	7.17	10.07
T ₆	<i>Trichoderma viride</i> + Glycerol (0.5%)	7.10	9.53
T ₇	<i>Pseudomonas fluorescens</i> + Glycerol (0.5%)	6.80	8.67
T ₈	<i>Bacillus subtilis</i> + Glycerol (0.5%)	6.40	8.50
T ₉	<i>Pseudomonas fluorescens</i> + <i>Bacillus subtilis</i> + Glycerol (0.5%)	6.80	9.23
T ₁₀	<i>Pseudomonas fluorescens</i> + <i>Trichoderma viride</i> + Glycerol (0.5%)	7.27	10.83
T ₁₁	Control (seedling dip in distilled water)	6.27	7.67
F-Test		**	**
SEm ±		0.11	0.43
CD at 1 %		0.43	1.71