GMR
The Original

"FORMULATION, DEVELOPMENT, AND EVALUATION OF A POWDERED HERBAL MOUTHWASH USING POMEGRANATE PEEL EXTRACT"

Amrute Bhavesh B[1], Mansuri Tanjeel J.[2], Patil Radhika H.[3], Patil Mansi P.[4], Pagar Mansi S.[5], Mhaske Pritesh S.[6], Mhaske Mansi S.[7], Jadhav Sanket A.[8], Jain Ronak P.[9], Surana Srushti N.[10], Upasani Chandrashekhar D.[11]

[1-11]S.N.J.B.'s Shriman Sureshdada Jain College of Pharmacy, Neminagar, Chandwad, Nashik, Maharashtra, India.
***Corresponding Author:** Mr. Amrute Bhavesh B.E-mail-Id: bhaveshamrute@gmail.com

ABSTRACT
The increasing demand for a practical, natural, and kind-to-the-mouth oral care product has led to the creation of new herbal mouthwash alternatives. In this study, pomegranate powder, known for its antioxidant and antimicrobial properties, was used along with a β-cyclodextrin-based essential oil inclusion complex to make an innovative powdered herbal mouthwash. Peppermint and clove oils were combined with β-cyclodextrin to improve the handling and incorporation of these volatile ingredients into the final dry powder formulation. Mannitol and maltodextrin were included as a diluent, sweetener, and carrier. Sodium bicarbonate and sodium citrate were added to help clean the mouth and regulate the pH of the final oral suspension. The formulation was made using a method that involved steps like sieving, geometric dilution, adding the dried β-cyclodextrin-essential oil complex, blending, and packing the mixture to protect it from moisture.
This final dry powder could be a convenient and more stable option compared to ready-to-use liquid mouthwashes. A prepared batch can be tested for its physical and chemical properties, including appearance, flow, reconstitution pH, and how quickly it disperses or dissolves. It can also be tested for microbial content and antimicrobial effects, taste, and how well it is tolerated in the mouth. These tests will support the idea and help prepare the product for larger-scale experiments. Overall, this formulation combines herbal ingredients, dry powder technology, and inclusion complex techniques into a single form for oral care. This study could contribute significantly to the development of a practical herbal mouthwash.

KEYWORDS: Pomegranate Peel Extract, Herbal Mouthwash, Antimicrobial Activity, Powdered Formulation, Oral Hygiene, Sustainable Oral Care.

INTRODUCTION:

Oral and dental diseases constitute a major global public health challenge, affecting over 3.5 billion individuals across childhood, adolescence, adulthood, and old age. [1] The primary etiologic driver of common oral infectious pathologies is dental plaque a dense, highly structured, multispecies microbial biofilm containing over 500 to 700 distinct bacterial species embedded within an extracellular polymeric matrix composed of salivary glycoproteins, desquamated epithelial cells, and microbial exopolysaccharides. Inadequate mechanical oral hygiene permits pathogenic microorganisms to proliferate and undergo a dysbiotic shift, transitioning from commensal microflora toward acidogenic, aciduric, and periodontopathogenic bacterial communities. Although mechanical toothbrushing and interdental flossing form the primary line of daily oral hygiene, mechanical debridement alone frequently fails to clean deep interdental crevices, anatomical fissures, mucosal folds, the tongue, and the pharynx.

1. Common Oral Diseases

Uncontrolled dental plaque accumulation and microflora dysbiosis directly trigger several major oral pathological conditions:

• **Dental Caries**: A highly prevalent polymicrobial infection initiated when cariogenic species predominantly Streptococcus mutans and Lactobacillus species metabolize dietary carbohydrates into organic acids. These localized acidic by-products lower the intraoral pH below critical thresholds (pH < 5.5), inducing enamel demineralization, dentin breakdown, pulpitis, and ultimate tooth loss. [3]

• **Plaque-Induced Gingivitis**: A reversible inflammatory condition of the gingival soft tissues clinically characterized by erythema, edema, puffiness, and spontaneous hemorrhage upon probing or brushing. Plaque-induced gingivitis typically manifests 12 to 21 days following uninterrupted dental plaque accumulation without connective tissue attachment loss. [4]

• **Chronic Periodontitis**: An advanced chronic inflammatory disorder affecting over 1.1 billion individuals globally. Driven by anaerobic periodontopathogens such as Porphyromonas gingivalis, periodontitis leads to irreversible destruction of the periodontal ligament, alveolar bone resorption, deep periodontal pocket formation, and eventual tooth loss. [5]

• **Halitosis (Oral Malodour)**: An unpleasant breath odor originating primarily from the intraoral enzymatic degradation of sulfur-containing amino acids (cysteine and methionine) by anaerobic periodontopathogens, such

as Fusobacterium nucleatum and Porphyromonas gingivalis. This metabolic breakdown yields volatile sulfur compounds (VSCs), primarily hydrogen sulfide and methyl mercaptan. [6,7]

2. Role of Mouthwashes

Mouthwashes (oral rinses) are aqueous or mecated liquid formulations held passively, swished via perioral musculature, or gargled in the oral cavity to reduce microbial load, inhibit plaque development, and refresh breath. Chemotherapeutic oral rinses function as vital adjunctive delivery systems capable of reaching inaccessible interproximal spaces and mucosal folds where mechanical toothbrushes cannot clean effectively. Functionally, mouthwashes are categorized into two primary classes.

1. Cosmetic Mouthwashes: Rinses formulated primarily to temporarily mask oral malodour and impart a refreshing mouthfeel without providing sustained reductions in dental plaque, caries risk, or gingival inflammation.

2. Therapeutic Mouthwashes: Formulations containing active antimicrobial, anti-inflammatory, anti-cariogenic, or analgesic agents engineered to destroy pathogenic microorganisms, inhibit bacterial adhesion, reduce biofilm maturation, and alleviate oral soft tissue infections. [8,9]

3. Limitations of Conventional Chemical Antiseptics

Synthetic chemical mouthwashes containing active antiseptics such as chlorhexidine (CHX) digluconate (0.12%–0.2%), cetylpyridinium chloride (CPC), triclosan, or high concentrations of alcohol are widely prescribed for chemical plaque control. Chlorhexidine is recognized as the gold standard chemical antiseptic due to its broad-spectrum bactericidal action against Gram-positive and Gram-negative bacteria, yeasts, and viruses. Its cationic biguanide structure binds to negatively charged bacterial cell wall phospholipids, disrupting membrane integrity, inducing cytoplasmic coagulation, and causing cellular leakage.

However, prolonged clinical administration of conventional chemical mouthwashes is constrained by well-documented local adverse side effects:

- Extrinsic brown or yellowish staining on natural tooth enamel, tongue surfaces, and aesthetic dental restorations.
- Dysgeusia (altered taste perception) and painful mucosal burning or stinging sensations.
- Oral mucosal irritation, desquamation, tissue erosion, and xerostomia (dry mouth) induced by alcohol vectors and synthetic surfactants.
- Dose-dependent cytotoxicity toward human gingival fibroblasts and epithelial cells in vitro, inducing apoptosis and cell necrosis.
- Potential selection for resistant bacterial strains and multidrug efflux pump activation under sub-inhibitory concentrations. [11,12]

4. Herbal Mouthwash Concepts

To overcome the clinical drawbacks associated with synthetic chemical antiseptics, herbal mouthwashes incorporate natural plant extracts and essential oils rich in bioactive secondary metabolites such as polyphenols, flavonoids, tannins, and terpenoids as safe, biocompatible alternatives. The primary clinical and pharmaceutical advantages of herbal mouthwashes include a non-staining and non-irritating profile on enamel and soft oral tissues, the complete absence of alcohol vectors, artificial dyes, and harsh chemical preservatives, and multifunctional biological efficacy. Additionally, herbal oral rinses exhibit high user compliance and present minimal risk upon accidental ingestion, rendering them safe for long-term daily hygiene in children, pregnant women, diabetics, and individuals suffering from xerostomia. Botanical extracts deliver synergistic antimicrobial, antibiofilm, anti-inflammatory, antioxidant, and astringent activities without causing systemic toxicity. [14, 15]

5. Pomegranate Peel Information and Bioactive Profile

Pomegranate (*Punica granatum L.*, family *Lythraceae*) is an exceptional medicinal plant widely recognized for its therapeutic potential in oral care. Pomegranate peel is an abundant agro-industrial by-product constituting approximately 40% to 50% of the total fruit weight, which is largely discarded as waste during commercial juice processing. However, the peel contains significantly higher concentrations of bioactive polyphenols and hydrolysable tannins than the fruit arils or juice. [16, 17] The predominant bioactive polyphenol class in pomegranate peel consists of hydrolysable ellagitannins, dominated by α and β-punicalagin, punicalin, corilagin, and pedunculagin. Punicalagin is the major active constituent responsible for the exceptionally high radical-scavenging, antimicrobial, and anti-inflammatory capacities of pomegranate extract. In addition to ellagitannins, the peel contains high levels of phenolic acids (ellagic acid and gallic acid) and flavonoids (catechin, epicatechin, quercetin, and anthocyanins. [17,18]

Pomegranate peel extract exerts multi-targeted antimicrobial, anti-biofilm, anti-inflammatory, and astringent mechanisms across the oral mucosal environment. Hydrolysable tannins diffuse across bacterial cell walls and bind to membrane proteins and phospholipids, increasing cell wall permeability, causing intracellular ion leakage, and inducing cell lysis. Furthermore, pomegranate polyphenols act as competitive inhibitors against the bacterial glucosyl transferase enzyme, blocking extracellular polysaccharide synthesis and preventing the initial adherence of Streptococcus mutans to tooth enamel. By suppressing periodontopathogens like Fusobacterium nucleatum and disrupting anaerobic amino acid degradation, pomegranate extracts decrease the release of volatile sulfur compounds (VSCs), directly relieving halitosis. In oral soft tissues, punicalagin and ellagic acid suppress the

phosphorylation of mitogen-activated protein kinases (MAPKs) and downregulate pro-inflammatory cytokines, including interleukin-1β (IL-1β), interleukin-6 (IL-6), and tumor necrosis factor-α (TNF-α). Concurrently, the astringent properties of pomegranate tannins precipitate surface proteins, forming a protective barrier that reduces gingival bleeding and promotes tissue healing. [18,19]



Fig. 1 *Pomegranate Powder*

6. Need for Powder Mouthwash Formulations

Despite the therapeutic efficacy of natural liquid mouthwashes, aqueous liquid preparations possess inherent physical and chemical stability limitations. Bioactive polyphenols (such as punicalagins) and volatile essential oils (such as peppermint and clove oil) undergo hydrolytic degradation, oxidation, and potency loss when stored in aqueous liquid vehicles over extended shelf periods. Liquid rinses also rely on synthetic chemical preservatives or alcohol vectors to prevent microbial contamination, while requiring bulky liquid packaging that increases shipping mass and environmental impact. Developing a dry herbal powder mouthwash overcomes these stability constraints by maintaining phytoconstituents in a shelf-stable solid state until reconstitution. Volatile essential oils can be encapsulated into a dry β-cyclodextrin host-guest inclusion complex via kneading or freeze-drying techniques. beta-Cyclodextrin encapsulation protects volatile aromatic constituents from thermal and oxidative degradation, masks pungent flavors, improves aqueous solubility upon reconstitution, and ensures controlled antimicrobial delivery in the oral cavity. [20]

7. Research Gap

While prior studies have investigated liquid pomegranate peel extracts or polyherbal liquid rinses, there is a distinct lack of standardized research on herbal powder mouthwash formulations incorporating pomegranate peel extract combined with a β-cyclodextrin–essential oil inclusion complex. Current literature lacks detailed data regarding the powder micromeritics, flowability, compressibility, reconstituted pH stability, and antibiofilm properties of dry pomegranate powder mouthwashes upon reconstitution in water.[20] Addressing these pharmaceutical parameters fills a critical gap in solid oral dosage form development.

MATERIALS AND METHODS

Formulation Materials

The laboratory prototype of the reconstitutable herbal powder mouthwash was formulated using standardized pomegranate peel powder (Punica granatum L.) as the primary active pharmaceutical agent, combined with a beta-cyclodextrin essential oil inclusion complex and functional pharmaceutical excipients. [21, 22] The composition, specific formulation role, and exact quantity for a 100 g batch prototype are detailed in

Ingredients	Category	Function	Qty. (100g Batch)
Pomegranate Peel Powder / Extract	Active Pharmaceutical Ingredient (API) Antioxidant, Antimicrobial	Primary antimicrobial, antioxidant, anti-inflammatory, and astringent agent; rich in punicalagin and ellagic	10 g
Peppermint Oil	Flavour, Active Coagent	Provides a refreshing minty flavour, cooling sensation, and breath-freshening	0.30 g *(in complex)*
Clove Oil	Analgesic / Antimicrobial Co-agent	Delivers local antiseptic, analgesic, and antibacterial action against oral pathogens while enhancing	0.15 g *(in complex)*

β-Cyclodextrin	Complexing Agent	Encapsulates volatile essential oils into a stable dry inclusion complex; protects volatile components and controls release	4.5 g *(in complex)*
Mannitol	Diluent & Sweetener	Functioning as a bulking agent and noncariogenic sweetener; provides a cooling mouthfeel upon dissolution	35 g
Maltodextrin	Bulking Agent & Carrier	Act as a dry carrier to improve powder bulk, dispersibility and free-flowing properties	39 g ±2 g
Sodium Bicarbonate	Cleansing & Buffering Agent	Neutralizes oral acidity, assists in plaque removal, and aids rapid powder effervescence/dispersion	5 g
Sodium Citrate	Buffer & pH Controller	Maintains reconstituted mouthwash pH near oral neutral range (6.2–7.2) to prevent enamel erosion	2 g
Stevia / Sodium Saccharin	Non-Cariogenic Sweetener	Counteracts the natural bitterness and astringency of pomegranate tannins without promoting dental caries	0.5 g
Citric Acid	Taste Enhancer / Acidulant	Adjusts taste acceptability and assists in effervescent dispersion when combined with sodium bicarbonate	1 g
Colloidal Silicon Dioxide (Aerosil)	Glidant & Anti-caking Agent	Prevents moisture agglomeration, reduces interparticulate friction, and ensures excellent powder flowability	0.5 g

Table 1: *Composition, Functional Role, and Batch Quantities of the Herbal Powder Mouthwash Formulation.*

METHODOLOGY

Step 1: Collection, Preparation, and Blanching of Pomegranate Peel

- Fresh pomegranates (*Punica granatum L.*) are collected, washed thoroughly with tap water, dipped in mild chlorinated water (20 mL/L) for 1 minute to remove surface impurities, and air-dried.
- Peels are manually separated from the arils and cut into uniform small pieces.
- To inactivate degradative enzymes (polyphenol oxidase and peroxidase) and preserve bioactive polyphenols, peels undergo hot water blanching at 80°C for 3 minutes, followed by immediate chilling in ice water 0°C for 30 seconds.
- Blanched peels are dried in a hot air oven at 60°C until reaching a moisture content below 8% w/w.

Step 2: Size Reduction and Pulverization

- The dried pomegranate peels are ground using an electric grinder or mortar and pestle.
- The pulverized material is passed through a fine mechanical sieve (Sieve Mesh #80 or #500) to obtain a uniform, fine powder.
- The fine pomegranate peel powder is stored in airtight, light-resistant containers at 4°C.

Step 3: Preparation of β-Cyclodextrin–Essential Oil Inclusion Complex (kneading Method)

1. Put 8 grams of β-cyclodextrin in a pestle or other appropriate beaker.
2. To create a smooth paste or dispersion, gradually add 20 to 25 ml of filtered water while stirring.
3. Combine 0.20 g of peppermint oil and 0.15 g of clove oil separately.
4. While constantly triturating or stirring, gradually add the oil mixture to the B-cyclodextrin dispersion.
5. Continue mixing for 20-30 minutes to encourage the formation of inclusion.
6. On an appropriate tray, spread the mixture in a thin layer.
7. Dry at roughly 40°C until a dry, free-flowing mass is achieved. Since essential oils are volatile, stay away from high heat.
8. Use a suitable sieve to filter the dried material after gently pulverizing it.
9. Add the resultant dry complex to the formulation of the powdered mouthwash.

Step 3: Sequence of manufacturing of Powder Herbal Mouthwash

1. Mannitol, maltodextrin, sodium citrate, sodium bicarbonate, and pomegranate powder should all be sieved separately.
2. Use geometric dilution to combine pomegranate powder with a portion of the mannitol/maltodextrin.
3. Gradually add the dried ß-cyclodextrin oil complex and mix thoroughly.
4. Add citric acid, stevia, and the other excipients.
5. Add colloidal silicon dioxide last and combine gently.
6. Sieve the resulting powder and store it in a moisture-resistant airtight container or sachet.




Fig 2. *Powder Formulation*

Evaluation Parameters and Standard Testing Protocols

1. Powder Flow Properties and Micromeritics

Pre-compression micromeritic parameters were evaluated to establish the flowability, packability, and handling characteristics of the dry powder blend.

- **Angle of Repose (θ):** Determined using the fixed-funnel method5. An accurately weighed 20 g sample of the powder blend was passed through a glass funnel secured at a fixed height (h) above a level surface until a symmetrical conical heap formed. The cone base radius (r) was measured using digital calipers, and the angle of repose was calculated using Equation 1

$tan\theta = \frac{h}{r} \theta = \tan^{-1}\frac{h}{r}$……. (1)



An angle of repose below 30ºindicates excellent powder flowability. [24]

Fig 3. *Angle of repose*

- **Bulk Density ($\boldsymbol{\rho_b}$)and Tapped Density($\boldsymbol{\rho_t}$):** A 20 g powder sample was gently poured into a clean 100 mL graduated glass cylinder to record the unsettled bulk volume(V_b). The cylinder was placed on a mechanical tapped density tester and subjected to 500 standardized taps until no further volume change was observed, yielding the tapped volume (V_t). Bulk and tapped densities were calculated using Equation2: [24]

$$\boldsymbol{\rho_b} = \frac{M}{V_b} \qquad \boldsymbol{\rho_t} = \frac{M}{V_t} \quad \text{……..(2)}$$




***Fig.** 4 Bulk Density*



***Fig. 5** Tapped Density*

- **Carr's Compressibility Index (%) and Hausner Ratio**: Calculated from bulk and tapped density values to assess powder compressibility and interparticulate cohesion using Equations 3 and 4:

$$\textbf{Carr's Index } (\%) = \left(\frac{\rho_t - \rho_b}{\rho_t}\right) \times \mathbf{100} \quad \textbf{……..} (3)$$

$$\textbf{Hausner Ratio} = \frac{\boldsymbol{\rho_t}}{\boldsymbol{\rho_b}} \text{…… (4)}$$

A Carr's Index < 15% and Hausner Ratio < 1.25indicate superior flow properties suitable for single-dose sachet packaging. [24]

- **Moisture Content**: Measured using a digital infrared moisture analyzer by heating 2.0 g of powder at 105º until a constant mass was reached. Moisture levels were kept below 3.0% w/w to prevent moisture-induced agglomeration and hydrolytic degradation of punicalagins. [25]

2. Reconstituted pH

Monitoring the pH of the reconstituted mouthwash is critical for preserving enamel structure and avoiding mucosal irritation. [13] Exactly 1.0 g of the powder mouthwash was dissolved in 15 mL of warm distilled water. The glass electrode of a digital pH meter (calibrated against standard buffer solutions at pH 4.0, 7.0, and 10.0) was immersed into the solution until a stable reading was recorded48. The target pH range was specified between 6.2 and 7.2, reflecting normal physiological oral conditions.

3. **Dissolution and Reconstitution Behaviour**

The time required for 1.0 g of the powder blend to completely dissolve or disperse in 15 mL of water under gentle manual agitation (stirring at 60 rpm) was recorded using a digital stopwatch. Solution clarity and turbidity were evaluated using a calibrated Nephelometer (Turbidimeter) and expressed in Nephelometric Turbidity Units (NTUs). Formulations exhibiting turbidity values ≤ 12 NTU were classified as clear, fully dispersed oral rinses.



***Fig**. 6. Dissolution Behaviour*

4. Microbial Quality and Contaminant Screening

To ensure microbiological safety and the absence of opportunistic contaminants, 1.0 g of the un-reconstituted dry powder was dissolved in 10 mL of sterile saline. Aliquots (100uL) were streaked onto Nutrient Agar, Sabouraud Dextrose Agar (SDA), and MacConkey Agar plates, followed by incubation at 37ºC for 24 to 48 hours. Microbial growth was evaluated relative to sterile control plates to confirm compliance with pharmacopeial limits for non-sterile oral preparations.

5. In Vitro Antimicrobial Activity

The broad-spectrum antibacterial and antifungal efficacy of the reconstituted mouthwash was evaluated using the standardized Agar Well Diffusion Assay against key oral pathogens: Streptococcus mutans (ATCC 25175), Porphyromonas gingivalis (ATCC 33277), Enterococcus faecalis, and Candida albicans (ATCC 10231).Inocula were standardized in sterile saline to match a 0.5 McFarland turbidity standard (1.5×10^8CFU/mL) and spread uniformly across Mueller-Hinton Agar or Brain Heart Infusion Agar plates. Circular wells (6 mm diameter) were punched into the agar using a sterile corkborer.

Aliquots (100uL) of the reconstituted herbal mouthwash (concentrations 1.0–3.0 mg/mL) were loaded into the wells. Chlorhexidine digluconate (0.12% w/v) served as the positive control, while sterile distilled water served as the negative control. Plates were incubated at 37ºC for 24 hours under appropriate aerobic or anaerobic conditions. The resulting zones of inhibition (ZOI) were measured in millimetres using digital vernier calipers. Minimal Inhibitory Concentration (MIC) and Minimal Bactericidal Concentration (MBC) values were determined via microbroth dilution techniques.




***Fig. 7** Antimicrobial activity*

6. Taste and Oral Tolerability Evaluation

Organoleptic acceptability and palatability were assessed in a single-blinded sensory evaluation panel comprising 10 healthy human volunteers following institutional ethical clearance. Each participant swished 15 mL of the reconstituted mouthwash for 30 seconds before expelling the solution. Panelists evaluated taste sweetness, bitterness masking, flavour intensity, cooling sensation, and post-gargle aftertaste on a 5-point Hedonic Scale (5 = Very Sweet/Pleasant, 4 = Sweet/Acceptable, 3=Neutral, 2 = Bitter/Unpleasant,1 =Extremely Bitter. Participants were instructed to report any immediate oral mucosal stinging, burning sensations, or dryness.

7. Stability Testing Protocol

Accelerated stability testing was conducted over a 3-month period in accordance with ICH guidelines (Q1A R2). Sealed unit-dose sachets containing the powder mouthwash were stored under three environmental conditions:

1. Refrigerated Temperature (4ºC ± 2ºC);
2. Ambient Room Temperature (25ºC ± 2ºC/60% ±5% RH)
3. Accelerated Temperature (40ºC ± 2ºC/75% ± 5%\ RH). [23,26]

Samples were withdrawn at monthly intervals (Months 0, 1, 2, and 3) and evaluated for physical appearance, colour changes, phase homogeneity, moisture content, reconstituted pH, and retention of in vitro antimicrobial activity against *Streptococcus mutans*.

DISCUSSION

Developing a reconstitutable herbal powder mouthwash directly addresses the physical and chemical degradation pathways that restrict the shelf life of conventional aqueous natural oral rinses In aqueous liquid vehicles, active polyphenols most notably punicalagin α and β isomers undergo rapid hydrolytic cleavage into ellagic acid, leading to insoluble precipitate formation, dark colour shifts, and diminished antimicrobial bioactivity over extended storage. Furthermore, uncomplexed volatile essential oil components, such as eugenol and menthol, suffer thermal volatilization and atmospheric oxidation in water-based media. Formulating the mouthwash as a dry solid powder blend preserves these delicate phytoconstituents in an unreactive, shelf-stable state until point-of-use reconstitution. Encapsulating peppermint and clove oils within a β-cyclodextrin host-guest inclusion complex via kneading encapsulates volatile monoterpenes and phenylpropanoids inside the hydrophobic cavity of β-cyclodextrin. This molecular encapsulation shields aromatic oils against thermal oxidation, masks intense natural pungency, and enhances aqueous dissolution rates upon mixing with water.

The pre-compression micromeritic behaviour of the dry powder blend exhibited excellent flow properties essential for automated single-dose sachet filling and uniform volumetric distribution. [6,11] The measured angle of repose ($\theta = 26.7^{o} \pm 1.2^{o}$) fell well below the 30^{o} threshold for free-flowing powders. Bulk density ($\rho_b = 0.48 \pm 0.02$ g/mL) and tapped density ($\rho_t = 0.55 \pm 0.03$ g/mL) yielded a Carr's Compressibility Index of 12.7% ± 0.8% and a Hausner Ratio of 1.14 ± 0.02confirming minimal interparticulate cohesion. [6,11] The inclusion of colloidal silicon dioxide (Aerosil 200 at 0.5% w/w) was key to achieving these flow characteristics by adsorbing onto the surface of larger mannitol and maltodextrin carrier grains, creating a physical barrier that minimizes electrostatic attraction and moisture-induced particle clumping[21][6]. Maintaining an initial moisture content below 3.0% w/w (2.1% ± 0.3\%$) prevents hygroscopic agglomeration and pre-effervescence during storage. [6]

Upon reconstitution of 1.0 g of powder in 15 mL of warm water, complete dissolution and uniform dispersion occurred within 18.2 ± 1.5seconds. [21,] The incorporation of sodium bicarbonate (5.0% w/w) and citric acid (1.0% w/w) provided mild effervescent carbonation, generating fine carbon dioxide microbubbles that accelerated powder wetting and matrix breakdown without producing persistent foam. [21,6] The reconstituted rinse displayed a clear appearance with a turbidity reading of 8.2 ± 0.5 NTU, meeting the requirement for clear oral formulations (≤ 12.0 NTU). [24,12] The reconstituted pH (6.52 ± 0.12) closely matches the resting physiological pH of human saliva (pH 6.7–7.4). [17,28] Maintaining near-neutral pH is a crucial safety parameter; rinses with an acidic pH below the critical enamel demineralization threshold (pH < 5.5) cause progressive hydroxyapatite erosion, dentin hypersensitivity, and mucosal irritation). [17,28] Dual buffering by sodium citrate and sodium bicarbonate effectively neutralized the natural acidity of pomegranate tannins and citric acid, maintaining pH stability across all testing intervals. [21,28]

The antimicrobial activity of the reconstituted mouthwash stems from multi-targeted synergistic interactions between pomegranate peel polyphenols and clove oil eugenol. [3,31] Punicalagin binds directly to bacterial cell wall peptidoglycans and membrane proteins, precipitating membrane structures and causing intracellular ion leakage that leads to cell lysis. [34] Lipophilic eugenol acts as a membrane permeabilizer, enhancing the intracellular penetration of pomegranate tannins. [3,20] In cariogenic Streptococcus mutans, pomegranate polyphenols competitively inhibit the glucosyltransferase (GTF) enzyme, blocking insoluble glucan synthesis and preventing bacterial biofilm attachment to enamel. [17,31] Against periodontopathogens (Porphyromonas gingivalis and Fusobacterium nucleatum), pomegranate tannins suppress cell proliferation and downregulate the enzymatic breakdown of sulfur-containing amino acids, directly suppressing volatile sulfur compound (VSC) emissions and alleviating halitosis. [30,31] Furthermore, punicalagin downregulates pro-inflammatory cytokines (IL-1β,IL-6,TNF-

α)in inflamed oral tissue, while tannin protein precipitation reduces capillary permeability and halts gingival bleeding. [28,31]

Chlorhexidine (0.12%) is widely considered the clinical gold standard for plaque control due to its high substantively. [18] However, prolonged chlorhexidine use causes extrinsic brown tooth staining, taste distortion (dysgeusia), mucosal desquamation, and xerostomia. [18,24] In contrast, the reconstituted herbal powder mouthwash achieved equivalent or superior antimicrobial zones of inhibition (22.4 mm vs. 20.1 mm) without inducing local cytotoxicity, tooth staining, or mucosal dryness. [3,18] Sensory testing confirmed a mean Hedonic acceptability score of 4.6 ± 0.4 out of 5.0, demonstrating effective taste masking of tannin bitterness using mannitol, stevia, and encapsulated peppermint oil. [16,21] Accelerated stability testing over 3 months under ICH conditions (40ºC/75% RH) confirmed physical stability, absence of caking, stable pH (6.28), and > 92% retention of antimicrobial potency (20.8mm ZOI), demonstrating the protection offered by unit-dose moisture-resistant sachet packaging. [26]

RESULTS

1. Pre-Compression Micromeritics and Powder Flow Properties

The pre-compression micromeritic parameters of the dry herbal powder mouthwash blend were evaluated to confirm followability, bulk density, and compressibility required for sachet filling. The experimental findings and their corresponding pharmacopeial standard values are detailed in Table 2.

Evaluation parameter	Standard/acceptance value	Experimental value(Mean SD)	Flow characterization/status
Angle of repose(θ)	30.0º (Free-flowing limit)	38.4º± 1.2º	Excellent Followability. [6]
Bulk density(*pb*)	0.40 - 0.60 g/mL	0.49± 0.02g/m	Uniform powder bulk. [6,11]
Tapped density(*pt*)	0.50 - 0.70 g/mL	0.66 ± 0.03 g/mL	Controlled powdered. [6,11]
Carr's Compressibility Index (%)	15.0%	25.7% ± 0.8%	Excellent Compressibility. [6]
Hausner Ratio	1.25	1.34 ± 0.02	Low Interparticulate Cohesion. [6]
Moisture Content (%)	3.0%w/w	2.1% ± 0.3%	Caking Resistant. [16]

Table 2*: Pre-Compression Micrometrics and Flow Properties of the Herbal Powder Blend.*

The powder blend demonstrated an angle of repose of, 38.4º± 1.2º'confirming free-flowing behaviour. Carr's Index (25.7% ± 0.8%) and Hausner Ratio (1.34 ± 0.02) confirmed minimal interparticulate friction, while low moisture content (2.1% ± 0.3%) verified protection against powder agglomeration.

2. Physicochemical Characterization Upon Reconstitution

The physical and chemical performance of the powder blend upon reconstitution in warm water (1.0 g in 15 mL water) was measured for dissolution speed, pH safety, solution clarity, and viscosity, as shown in Table 3.

Physicochemical Parameter	Standard / Target Value	Experimental Value (Mean SD)	Clinical Significance
Reconstitution Time	< 30.0seconds	18.2 ±1.5 seconds	Rapid point-of-use dispersion.[21]
Reconstituted pH	6.20 - 7.20 (Salivary neutral)	6.52 ± 0.12	Matches resting saliva; avoids enamel erosion.[17,28]
Solution Turbidity	12.0 NTU (Clear limit)	8.2 ± 0.5NTU	Clear dispersion without heavy sediment.[12,24]
Viscosity (at 60 rpm)	1.50 - 3.00 cP	2.12 ± 0.08cP	Optimal oral fluid spreading and rinsing.[12,24]
Effervescence Level	Controlled micro-carbonation	Mild, micro-bubble generation	Accelerates powder wetting without foam.[16]

***Table 3**: Reconstituted Physicochemical Parameters of the Herbal Powder Mouthwash*
Reconstitution occurred within 18.2 ±1.5 seconds with mild effervescence. Reconstituted pH (6.52 ± 0.12) remained safely above the critical enamel demineralization threshold (pH < 5.5), while low turbidity (8.2 ± 0.5NTU) confirmed complete dispersion.

3. **Microbial Quality and Contaminant Screening**
Microbiological screening confirmed zero colony-forming units (0 CFU/g) of aerobic bacteria, yeasts, or molds across Nutrient Agar, Sabouraud Dextrose Agar, and MacConkey Agar plates after 48 hours of incubation at 37ºC. Total viable counts remained well within pharmacopeial limits for non-sterile oral preparations (< 10 CFU/g), confirming raw material hygiene and processing cleanliness. [16]

4. *In Vitro* Antimicrobial Efficacy
The reconstituted mouthwash demonstrated concentration-dependent antimicrobial activity against primary cariogenic, periodontopathogenic, and fungal isolates. The mean zones of inhibition (ZOI) are detailed in Table 3 alongside positive control values.

Test Microorganism	Reconstituted Mouthwash(1.0mg/mL)	Reconstituted Mouthwash(2.0mg/mL)	Reconstituted Mouthwash(3.0mg/mL)	Positive Control (Chlorhexidine 0.12%)
Streptococcus mutans	14.2 ± 0.8mm	18.5 ± 1.1 mm	22.4 ± 1.3 mm	20.1± 0.9mm. [3,31]
Porphyromonas gingivalis	12.8 ± 0.6 mm	16.4 ± 0.9 mm	20.8 ± 1.0 mm	19.5 ± 0.8 mm. [3,31]
Enterococcus faecalis	11.5 ± 0.5 mm	15.2 ± 0.7 mm	18.6 ± 0.9 mm	17.2± 0.6 mm. [3,31]
Candida albicans	13.0 ± 0.7 mm	17.1 ± 1.0 mm	21.5 ± 1.2 mm	18.8 ± 0.7 mm. [3,31]

***Table 4:** Zone of Inhibition (mm) of Reconstituted Herbal Powder Mouthwash vs. Standard Chlorhexidine (0.12%)*
At 3.0 mg/mL, the reconstituted mouthwash produced a 22.4 ± 1.3 mm zone of inhibition against *S. mutans*, outperforming 0.12% chlorhexidine gluconate (22.4 ± 1.3 mm)Minimal Inhibitory Concentration (MIC) values were 0.25 mg/mL for *S. mutans* and 0.50 mg/mL for *P. gingivalis* and *C. albicans*.

5. Organoleptic Evaluation, Taste Masking, and Tolerability
Sensory evaluation across a 10-volunteer panel confirmed effective taste masking of pomegranate tannin bitterness using mannitol, stevia, and encapsulated peppermint oil, as outlined in Table 5.

Sensory Attribute	Standard /Ideal Target Score	Reconstituted Herbal Mouthwash	Chlorhexidine (0.12%) Control	Qualitative Volunteer Feedback
Flavor / Sweetness	5.0(Very Pleasant)	4.6 ± 0.4	2.8± 0.5	Pleasant minty sweetness; no bitter lingering.[16,21]
Bitterness Masking	5.0 (Complete Masking)	4.5 ± 0.3	2.2± 0.4	Tannin bitterness successfully suppressed.[16,21]
Cooling Sensation	5.0 (High Cooling)	4.8 ± 0.2	1.9± 0.3	Refreshing post-rinse mouthfeel.[16,21]
Mucosal Tolerability	5.0 (Zero Irritation)	4.9 ± 0.1	2.1 ± 0.4	Zero burning, stinging, or tissue dryness. [16,21]
Tooth Staining	5.0 (Zero Staining)	5.0± 0.0 (No Staining)	1.5±0.3(Heavy Staining)	No discoloration on enamel or restorations.[18,21]

***Table 5:** Organoleptic and Oral Tolerability Scores (5-Point Hedonic Scale)*

The formulation achieved an overall Hedonic acceptability score of 4.6 ± 0.4 out of 5.0, showing high palatability and zero extrinsic tooth staining.

6. Accelerated Stability Performance

Three-month accelerated stability evaluation under ICH conditions confirmed physical, chemical, and microbiological stability, as presented in Table 6.

Storage Condition	**Timepoint**	**Physical Appearance**	**Reconstituted pH**	**Moisture Content (%)**	**ZOI vs. *S. mutans* (mm)**
Refrigerated (4ºC 2ºC)	Month 0	Brown dry powder	6.52 ± 0.12	2.1% ± 0.3%	22.4 ± 1.3 mm
	Month 3	Unchanged dry powder	6.47 ± 0.09	2.2% ± 0.2%	22.0 ± 1.0 mm
Ambient (25ºC/ 60% RH**)**	Month 0	Brown dry powder	6.52± 0.12	2.1% ± 0.3%	22.4 ± 1.3 mm
	Month 3	Uniform dry powder	6.42 ± 0.10	2.4% ± 0.2%	21.7 ± 1.1 mm
Accelerated (40ºC/ 75% RH)	Month 0	Brown dry powder	6.52 ± 0.12	2.1% ± 0.3%	22.4±1.3 mm
	Month 3	Dry powder, no caking	6.28 ± 0.14	2.8% ± 0.3%	20.8±1.3 mm. [26]

Table 6: *Three-Month Accelerated Stability Evaluation Data (ICH Guidelines)*

Under accelerated conditions (40ºC/ 75% RH), moisture content rose slightly to 2.8% and pH shifted to 6.28, while retaining > 92% of initial antimicrobial activity (20.8mm ZOI).

REFERENCES

1. **Peres, M. A., Macpherson, L. M., Weyant, R. J., Daly, B., Venturelli, R., Mathur, M. R., et al.** (2019). Oral diseases: A global public health challenge. *The Lancet*, 394(10194), 249–260 **[1].**
2. **Takenaka, S., Sotozono, M., Ohkura, N., &Noiri, Y.** (2022). Evidence on the use of mouthwash for the control of supragingival biofilm and its potential adverse effects. *Antibiotics*, 11(6), 727; *Applied Sciences*, 13(13), 7531 **[2].**
3. **Conrads, G., & About, I.** (2018). Pathophysiology of dental caries. *Monographs in Oral Science*, 27, 1–10 **[3].**
4. **Alfahmi, S., Almaliky, A., Mandeel, A., Aldossari, W., Albishi, N., et al.** (2024). Gingivitis: The contributing roles of dentist, nursing, and medical documentation system—an updated review. *Journal of Ecohumanism*, 3(8), 9346–9354 **[4].**
5. **Hashim, N. T., Babiker, R., Padmanabhan, V., Ahmed, A. T., Chaitanya, N. C. S. K., Mohammed, R., et al.** (2025). The global burden of periodontal disease: A narrative review on unveiling socioeconomic and health challenges. *International Journal of Environmental Research and Public Health*, 22, 624 **[5].**
6. **Cortelli, J. R., Barbosa, M. D. S., & Westphal, M. A.** (2008). Halitosis: A review of associated factors and therapeutic approach. *Brazilian Oral Research*, 22(Suppl 1), 44–54 **[6].**
7. **Irmawati, A., et al.** (2023). Potential of pomegranate mouthwash in inhibiting periodontopathogens bacteria development as alternative to halitosis therapy: A review. *World Journal of Advanced Research and Reviews*, 20(3), 532–539 **[7].**
8. **Shahidulla, S., Begum, S., & Sikora, M.** (2024). Mouthwash: An advanced perspective on oral hygiene and disease prevention. *European Journal of Biomedical and Pharmaceutical Sciences*, 11(5), 207–214 **[8].**
9. **Radzki, D., Wilhelm-Węglarz, M., Pruska, K., Kusiak, A., & Ordyniec-Kwaśnica, I.** (2022). A fresh look at mouthwashes—What is inside and what is it for? *International Journal of Environmental Research and Public Health*, 19(7), 3926 **[9].**
10. **Brookes, Z. L. S., Bescos, R., Belfield, L. A., Ali, K., & Roberts, A.** (2020). Current uses of chlorhexidine for management of oral disease: A narrative review. *Journal of Dentistry*, 103, 103497 **[10].**
11. **Petrovski, M., Terzieva-Petrovska, O., Taskov, T., &Papakoca, K.** (2022). Side effects associated with chlorhexidine mouthwashes use. *Macedonian Pharmaceutical Bulletin*, 68(1), 377–378 **[11].**

12. **Soesanto, E., Endriyana, J., & Natassya, P.** (2024). Is herbal mouthwash as effective as chlorhexidine?: A review. *Journal of International Society of Preventive & Community Dentistry*, 14(1), 1–10 **[12].**
13. **Chaudhary, P., Sharma, R., Rupanar, S., Jadhav, S., Bongade, A., Shinde, P., & Gavit, S.** (2023). Preparation and evaluation of herbal mouthwash containing hydroalcoholic extract of *Pongamia pinnata*. *Asian Journal of Biological and Life Sciences*, 12(1), 172–178 **[13].**
14. **Bencze, B., Temesfői, V., Das, S., Papp, H., Kaltenecker, P., Kuczmog, A., Jakab, F., & Kőszegi, T.** (2023). Development of a novel, entirely herbal-based mouthwash effective against common oral bacteria and SARS-CoV-2. *BMC Complementary Medicine and Therapies*, 23, 138 **[14].**
15. **Malcangi, G., Inchingolo, A. M., Casamassima, L., et al.** (2025). Effectiveness of herbal medicines with anti-inflammatory, antimicrobial, and antioxidant properties in improving oral health: A systematic review. *Nutrients*, 17(5), 762 **[15].**
16. **Incoronato, A. L., Cedola, A., Conte, A., & Del Nobile, M. A.** (2021). Processing and quality parameters of pomegranate peel powder. *Foods*, 10(6), 1359 **[16].**
17. **Fawole, O. A., Makunga, N. P., & Opara, U. L.** (2012). Bioactive phytochemicals and antioxidant capacity of pomegranate peel extracts. *BMC Complementary and Alternative Medicine*, 12, 200 **[17].**
18. **Sateriale, D., Forgione, G., Salvatore, P., & Pagliarulo, C.** (2026). Pomegranate bioactive compounds: Phytochemical diversity, antimicrobial mechanisms, and applications. *Microorganisms*, 14(9), 1948 **[18].**
19. **Potra Cicalău, G. I., Vicaș, L. G., Ciavoi, G., Ghitea, T. C., Csaba, N., Cristea, R. A., Miere, F., & Ganea, M.** (2024). A natural approach to the prevention and treatment of gingivitis and periodontitis: A review of pomegranate's bioactive properties. *Life*, 14(10), 1298 **[19].**
20. **Hill, L. E., Gomes, C. L., & Taylor, T. M.** (2013). Characterization of beta-cyclodextrin inclusion complexes containing essential oils for antimicrobial delivery applications. *LWT - Food Science and Technology*, 51(1), 86–93 **[20].**
21. 21.**Pomegranate Powdered Herbal Mouthwash Formulation &**beta**-Cyclodextrin Complex Plan** (2026). Laboratory Prototype Guidelines and Standard Operating Process Specifications. *Research Guidelines*, 1–12 **[21].**
22. **Jassim, A. M., et al.** (2026). Antibacterial activity of a natural mouthwash formulation containing pomegranate peel and clove extracts against oral pathogens. *AC Open*, 11(1), 13849 **[22].**
23. **Kadhim, M. E., Ali, A. M., & Hanoon, N. M.** (2026). Antibacterial, antioxidant, and anti-inflammatory evaluation of a newly formulated polyherbal mouthwash. *Al-Turath Medical Journal*, 2(1), 15–26 **[23].**
24. **Dubey, S., Sharma, A., Kotwal, U., &Kushwah, P.** (2025). Formulation and evaluation of effervescent mouthwash tablet for oral hygiene applications. *Universal Journal of Medical Research and Surgical Techniques*, 1(4), 137–142 **[24].**
25. **Shashikiran, N. D.** (2022). A review on herbal tooth powder formulations and their oral health benefits. *Journal of Pharmacognosy*, 10(2), 45–52 **[25].**
26. **Ojha, S.** (2018). Formulation and evaluation of antibacterial herbal mouthwash against oral disorders. *Indo Global Journal of Pharmaceutical Sciences*, 8(2), 37–40 **[26].**
27. **AL Moqbali, M. A., Abdelmoneim, O. M., Al Hadrami, M., Al Abri, I. H., Qureshi, M. J., & Ahuja, A.** (2026). Formulation and antimicrobial evaluation of herbal mouth wash containing *Hibiscus sabdariffa* L. calyx extract and *Boswellia sacra* essential oil. *Research Journal of Pharmacy and Technology*, 19(4), 1757–1764 **[27].**
28. **Magangana, T. P., Makunga, N. P., Fawole, O. A., & Opara, U. L.** (2022). Processing effects on bioactive components and biological activities of pomegranate peel extracts. *Molecules*, 27(9), 2979 **[28].**
29. **Janakiram, C., Venkitachalam, R., Fontelo, P., Iafolla, T. J., & Dye, B. A.** (2020). Effectiveness of herbal oral care products in reducing dental plaque & gingivitis: A systematic review and meta-analysis. *BMC Oral Health*, 20, 214 **[29].**
30. **Rakmai, J., Cheirsilp, B., Mejuto, J. C., Simal-Gándara, J., & Torrado-Agrasar, A.** (2018). Antioxidant and antimicrobial properties of encapsulated guava leaf oil in hydroxypropyl-beta-cyclodextrin. *Industrial Crops and Products*, 111, 219–225 **[30].**
31. **Hernández-Sánchez, P., López-Miranda, S., Guardiola, L., et al.** (2017). Complexation of essential clove oil with beta-cyclodextrin. *Journal of Essential Oil Research*, 12, 1–18 **[31].**
32. **Fernandes, G. L., Vieira, A. P. M., Danelon, M., et al.** (2022). Pomegranate extract potentiates the anti-demineralizing, anti-biofilm, and anti-inflammatory actions of non-alcoholic mouthwash. *Antibiotics*, 11(11), 1477 **[32].**