



Research Note

New microsatellite markers for bananas (*Musa* spp)

E.P. Amorim¹, P.H. Silva², C.F. Ferreira¹, V.B.O. Amorim¹, V.J. Santos³,
A.D. Vilarinhos¹, C.M.R. Santos⁴, M.T. Souza Júnior⁵ and R.N.G. Miller⁶

¹Embrapa Mandioca e Fruticultura, Cruz das Almas, BA, Brasil

²Universidade Federal de Lavras, Lavras, MG, Brasil

³Universidade Federal do Recôncavo da Bahia, Cruz das Almas, BA, Brasil

⁴Embrapa Recursos Genéticos e Biotecnologia, Brasília, DF, Brasil

⁵Embrapa Agroenergia, Brasília, DF, Brasil

⁶Departamento de Biologia Celular, Instituto de Ciências Biológicas,
Universidade de Brasília, Brasília, DF, Brasil

Corresponding author: E.P. Amorim

E-mail: edson@cnpmf.embrapa.br

Genet. Mol. Res. 11 (2): 1093-1098 (2012)

Received September 12, 2011

Accepted January 16, 2012

Published April 27, 2012

DOI <http://dx.doi.org/10.4238/2012.April.27.8>

ABSTRACT. Thirty-four microsatellite markers (SSRs) were identified in EST and BAC clones from *Musa acuminata burmannicoides* var. Calcutta 4 and validated in 22 *Musa* genotypes from the Banana Germplasm Bank of Embrapa-CNPMF, which includes wild and improved diploids. The number of alleles per locus ranged from 2 to 14. The markers were considered highly informative based on their polymorphism information content values; more than 50% were above 0.5. These SSRs will be useful for banana breeding programs, for studies of genetic diversity, germplasm characterization and selection, development of saturated genetic linkage maps, and marker assisted selection.

Key words: Genetic breeding; Microsatellites; Primer validation;
Musa spp