



Genome-wide identification, phylogenetic relationships, and expression analysis of the carotenoid cleavage oxygenase gene family in pepper

X.H. Zhang¹, H.Q. Liu¹, Q.W. Guo¹, C.F. Zheng², C.S. Li¹, X.M. Xiang¹,
D.F. Zhao¹, J. Liu³, J. Luo⁴, D.K. Zhao⁵, J.Q. Zheng⁶ and H.J. Wan⁷

¹Quzhou Academy of Agricultural Sciences, Quzhou, Zhejiang, China

²Zhejiang Mariculture Research Institute, Wenzhou, China

³Wulanchabu Academy of Agricultural Sciences, Wulanchabu, Inner Mongolia, China

⁴Institute of Digital Agriculture, Zhejiang Academy of Agricultural Sciences, Hangzhou, Zhejiang, China

⁵College of Agricultural, Yunnan University, Kunming, Yunnan, China

⁶Jiangsu Coastal Area Institute of Agricultural Sciences, Yancheng, Zhejiang, China

⁷State Key Laboratory Breeding Base for Zhejiang Sustainable Pest and Disease Control, Institute of Vegetables, Zhejiang Academy of Agricultural Sciences, Hangzhou, Zhejiang, China

Corresponding author: H.J. Wan

E-mail: wanhongjian@sina.com

Genet. Mol. Res. 15 (4): gmr.15048695

Received April 5, 2016

Accepted May 16, 2016

Published October 5, 2016

DOI <http://dx.doi.org/10.4238/gmr.15048695>

Copyright © 2016 The Authors. This is an open-access article distributed under the terms of the Creative Commons Attribution ShareAlike (CC BY-SA) 4.0 License.

ABSTRACT. Carotenoid cleavage oxygenases (*CCOs*) are a family of dioxygenases, which specifically catalyze the cleavage of conjugated

double bonds in carotenoids and apocarotenoids in plants. In this study, genome-wide analysis of *CCO* genes in pepper plants was performed using bioinformatic methods. At least 11 members of the *CCO* gene family were identified in the pepper genome. Phylogenetic analysis showed that pepper and tomato *CCO* genes could be divided into two groups (*CCDs* and *NCEDs*). The *CCD* group included five sub-groups (*CCD1*, *CCD4*, *CCD7*, *CCD8*, and *CCD-like*). These results indicate that there is a close genetic relationship between the two species. Sequence analysis using the online tool, Multiple Expectation Maximization for Motif Elicitation (MEME), showed that the *CCO* proteins comprise multiple conserved motifs, with 20 to 41 amino acids. In addition, multiple *cis*-acting elements in the promoter of *CCO* genes were identified using the online tool PlantCARE, and were found to be involved in light responsiveness, plant hormone regulation, and biotic and abiotic stresses, suggesting potential roles of these proteins under different conditions. RNA-seq analysis revealed that the *CCO* genes exhibit distinct patterns of expression in the roots, stems, leaves, and fruit. These findings suggest that the *CCO* genes have important roles in the vegetative and reproductive development of pepper plants.

Key words: Carotenoids; *CCO* genes; Bioinformatics; Pepper

INTRODUCTION

Carotenoids are a class of lipophilic compounds that contain C40 as their basic skeleton, and are comprised of several conjugated double bonds. To date, more than 700 types of C40 carotenoids have been identified (Britton et al., 2004). Carotenoids have important biological functions in organisms. Approximately 50 carotenoids are vitamin A precursors, which are needed for the human body to fight against cancer (Olson, 1989). Carotenoids are the main factors contributing to the yellow, orange, and red in color in some fruits (Nagal et al., 2012; Jabeen et al., 2013; Jarquín-Enríquez et al., 2013). In addition, some carotenoids participate in photosynthesis, and play an important role in light absorption and electron transport, and in the removal of triplet oxygen and superoxide anion species (Bartley and Scolnik, 1995; Tracewell et al., 2001; Woitsch and Römer, 2003). The results also showed that carotenoid cleavage products possess important biological functions. Some of the apocarotenoids derived from carotenoid cleavage are important determinants of flavor in agricultural products. In addition, some products of carotenoid-derived zeaxanthin aldehyde, which can be transformed into the plant hormone abscisic acid, can regulate stress, seed development, and other important functions (Winterhal and Schreier, 1995; Huang et al., 2009; Ilg et al., 2010; Liang et al., 2011; Heo et al., 2013; Sui et al., 2013). These cleavage products are mainly catalyzed by the carotenoid cleavage oxygenases (CCOs) (Bouvier et al., 2005; Heo et al., 2013).

CCOs are a class of dioxygenases, which specifically catalyze the cleavage of conjugated double bonds of carotenoids and apocarotenoids (Ilg et al., 2009; Walter et al., 2010). *CCOs* can be further classified as carotenoid cleavage dioxygenases (*CCDs*) and *cis*-epoxycarotenoid dioxygenases (*NCEDs*), based on their substrate informing an epoxy structure (Tan et al., 2003; Auldridge et al., 2006a). Recently, several studies have focused on

the identification and analysis of *CCO* genes of various plants (Simkin et al., 2004; Ohmiya et al., 2006; Sun et al., 2008; Adami et al., 2013; Liu et al., 2013). For example, the gene encoding the enzyme NCED (*Vp14*) was identified in maize (Tan et al., 1997; Woitsch and Römer, 2003). Homologs of this gene were subsequently discovered in other higher plants. In *Arabidopsis*, four of nine *CCO* genes are of the *CCD* type, and the remaining five have been identified as *NCED* genes (Schwartz et al., 2004; Auldridge et al., 2006b).

In the present study, 11 *CCO* genes were identified in the pepper genome. A comprehensive analysis of the *CCO* gene was performed, including sequence alignments, and determination of phylogenetic relationships and expression patterns. These results would aid in better understanding of the function and regulatory mechanisms of *CCO* genes in pepper.

MATERIAL AND METHODS

Identification of *CCO* genes in pepper plants

Pepper (*Capsicum annuum* L.) plant assembly and annotation V1.55 were downloaded from the PGP (Pepper Genome Platform) database (<http://passport.pepper.snu.ac.kr/?t=PGENOME>). A TBLASTp was performed using the protein coding sequence of the tomato *CCO* gene as the query against the pepper genome database. Subsequently, searches of candidate *CCO* genes in the pepper genome were repeated using BLASTp. The e-value used was 1e-5. Next, all candidate genes were evaluated for further verification using a Pfam database (<http://pfam.janelia.org/>), and SMART protein motif analyses (<http://smart.embl-heidelberg.de/>), in order to classify the *CCO* genes.

Alignment and phylogenetic analysis of *CCO* gene families

The amino acid sequences encoded by *CCO* genes in the pepper genome were aligned using Clustal X version 1.8, followed by manual adjustment, and were used to construct a phylogenetic tree using the Molecular Evolutionary Genetics Analysis software version 5.0 (MEGA 5.0) (Tamura et al., 2011). In addition, bootstrapping (1000 replicates) was performed to evaluate the degree of support for a particular grouping pattern in the phylogenetic tree of the *CCO* gene family. Missing sequence data were treated using pairwise deletions of the gaps. The branch lengths were assigned by utilizing pairwise calculations of the genetic distances.

Prediction of conserved motifs

To further investigate the diversity and structure of the *CCO* genes in pepper plants, their amino acid sequences were subjected to motif analyses using a Multiple Expectation Maximization for Motif Elicitation (MEME) (<http://meme.sdsc.edu/meme/website/intro.html>). Then, the optimal matching length of the parameter was set between 6 and 50, and all other parameters were set to the default values. Conservation of each motif among the *CCO* genes was determined with WebLogo version 2.8.2 (<http://weblogo.berkeley.edu/>) using the default settings.

Analysis of the *CCO* gene structure and its promoter

The exon and intron positions of pepper *CCO* genes were analyzed using the online

Gene Structure Display Server (GSDS: <http://gsds.cbi.pku.edu.cn>), with both coding and genomic sequences. In addition, the 5'-upstream domain (1500 bp) of each *CCO* gene was downloaded from the PGP (<http://peppergenome.snu.ac.kr/>). The promoter sequences were then used to scan for *cis*-elements in the PlantCARE database (<http://bioinformatics.psb.ugent.be/webtools/plantcare/html/>).

Expression analysis of *CCO* genes

To determine the expression profiles of pepper *CCO* genes in different tissues and in response to different stress conditions, the table reads per kb per million reads (RPKM) of the pepper genes was searched against RNA-seq data of each tissue (<http://www.nature.com/ng/journal/v46/n3/full/ng.2877.html>) utilizing the locus ID given in the PGP (<http://passport.pepper.snu.ac.kr/?t=PGENOME>). The data obtained were analyzed and grouped based on tissue specificity.

RESULTS

Genome-wide identification of pepper *CCO* genes

In this study, 11 *CCO* genes were identified in the pepper genome, and named *CaCCD1*, *CaCCD4a*, *CaCCD4b*, *CaCCD7*, *CaCCD8*, *CaCCD-like*, *CaCCD-like2*, *CaCCD-like3*, *CaNCED*, *CaNCED2*, and *CaNCED3*, based on homology with tomato and *Arabidopsis* genes (Table 1). Gene names, gene locus, chromosome locations, gene lengths, exon numbers, protein lengths, molecular weights, and isoelectric points are also shown in Table 1. *CCO* genes ranged in length from 954 bp (*CaNCED*) to 15,419 bp (*CaCCD1*). The number of amino acids ranged from 317 amino acids (*CaNCED*) to 646 amino acids (*CaCCD7*). The molecular weight varied from 34.96 kDa (*CaNCED*) to 72.72 kDa (*CaCCD7*), and the isoelectric point varied from 5.40 (*CaCCD-like2*) to 9.39 (*CaNCED*).

Table 1. *CaCCO* genes and properties of their deduced proteins.

Gene name	Description	Gene locus	Chromosome location	Gene length (bp)	Number of exons	Protein size (aa)	MW (kDa)	pI
<i>CaCCD1</i>	Carotenoid cleavage dioxygenase 1	CA01g20280	1	15,419	12	584	65.78	6.41
<i>CaCCD4a</i>	Carotenoid cleavage dioxygenase 4	CA01g28360	1	3937	4	475	52.87	6.86
<i>CaCCD4b</i>	Carotenoid cleavage dioxygenase 4	CA01g08910	1	4469	2	603	66.04	6.34
<i>CaCCD7</i>	Carotenoid cleavage dioxygenase 7	CA00g32540	0	6191	7	646	72.72	6.27
<i>CaCCD8</i>	Carotenoid cleavage dioxygenase 8	CA11g14280	11	2000	2	366	41.30	8.82
<i>CaCCD-like</i>	9-cis-epoxycarotenoid dioxygenase, putative	CA08g04710	8	3139	12	576	65.39	5.74
<i>CaCCD-like2</i>	9-cis-epoxycarotenoid dioxygenase, putative	CA11g20400	11	8262	12	575	65.85	5.40
<i>CaCCD-like3</i>	9-cis-epoxycarotenoid dioxygenase, putative	CA01g34880	1	5126	12	597	68.59	5.86
<i>CaNCED</i>	9-cis-epoxycarotenoid dioxygenase (Fragment)	CA07g16140	7	954	1	317	34.96	9.39
<i>CaNCED2</i>	9-cis-epoxycarotenoid dioxygenase 5	CA08g03620	8	1806	1	601	-	-
<i>CaNCED3</i>	9-cis-epoxycarotenoid dioxygenase, putative	CA05g17080	5	1740	1	579	64.67	8.25

Phylogenetic relationships of pepper *CCO* genes

Phylogenetic analysis showed that pepper and tomato *CCO* genes could be divided into two groups (*CCD* and *NCED* groups). The former contained five sub-groups (*CCD1*, *CCD4*, *CCD7*, *CCD8*, and *CCD-like*), and the latter contained three sub-groups (*NCED*, *NCED2*, and *NCED3*) (Figure 1). Among these, *SICCD1a*, *SICCD4b*, *SICCD7*, and *SICCD8* in tomato were orthologous to *CaCCD1a*, *CaCCD4b*, *CaCCD7*, and *CaCCD8* in pepper, respectively. Furthermore, *SINCED*, *SINCED2*, and *SINCED3* in tomato were orthologous to

CaNCED, *CaNCED2*, and *CaNCED3* in pepper, respectively. However, there were two copies of *CCD1* in the tomato genome (*SICCD1a* and *SICCD1b*) (Wei et al., 2016), while only *CCD1* (*CaCCD1a*) was observed in the pepper genome. In contrast, only one member of the *CCD-like* (*SICCD-like*) gene was found in tomato (Wei et al., 2016), and three homologous genes (*SICCD-like*, *SICCD-like2*, *SICCD-like3*) in pepper were very similar to the *CCD-like* genes in tomato. These results indicated that some gene loss or gain events may have occurred during the course of evolution between the pepper and tomato plants.

Genome-wide analysis of the *CCO* gene family has been performed in several plant species, including *Arabidopsis*, tomato, rice, maize, and sorghum, which were found to contain nine, ten, five, six, and nine members, respectively (Auldridge et al., 2006b; Rubio et al., 2008). Therefore, the *CCO* gene family in plants might encode a small family of proteins with only a few gene members.




Figure 1. Phylogenetic tree of carotenoid cleavage oxygenase (CCO) proteins from pepper and tomato was constructed with the MEGA5.0 software using neighbor-joining method.

Multiple-sequence alignment and conserved motif analysis of the CCO proteins

To determine whether there was high-sequence homology between the CCO proteins from pepper and tomato, the amino acid sequences of all *CCO* genes in the pepper and tomato were aligned using the ClustalX software program. Poor conservation was observed in these proteins (Figure S1). This result is consistent with the findings in other plant species (Auldridge et al., 2006a,b; Vallabhaneni et al., 2010; Walter et al., 2010; Lashbrooke et al., 2013; Wei et al., 2016).

Using the MEME online software, the 10 conserved motifs (Motif1-Motif10) were identified in all CCO proteins from pepper plants (Table 2 and Figure 2). Seven genes (*CaCCD1*,

CaCCD4b, *CaCCD-like*, *CaCCD-like2*, *CaCCD-like3*, *CaNCED2*, and *CaNCED3*) contained all of the 10 conserved motifs. *CaCCD4a* contained eight conserved motifs, but not Motif5 and Motif9. *CaCCD7* contained seven conserved motifs, but not Motif2, Motif5, and Motif8, and *CaNCED* contained five conserved motifs. *CaCCD8* only contained two conserved motifs (Motif5 and Motif6).

In addition, the length of these conserved motifs ranged from 20 to 41 amino acid residues (Motif8 and Motif6, respectively). Motif7 was comprised of 30 amino acid residues. Motif9, Motif10 and Motif5 contained 29 amino acid residues. Motif1 was comprised of 28 amino acid residues. Both Motif2 and Motif3 contained 27 amino acid residues, and Motif4 contained 21 amino acid residues (Table 2).

Table 2. Motifs of CCO proteins in pepper.

Motif	Width	Best possible match
1	28	GTGVANTNLFYHGGRYYAMAEDDMPYEI
2	27	RYYGDENSIKWFEVPPCCCFHLWNAWE
3	27	DEDDGWIIAYTHNENTWQSQVYIIDAK
4	21	EPVAIVKLPSPVYPYGFHGAFM
5	29	CPNGVYVRNGANPLFGPLAGHHWFDGDSM
6	41	MDRCSMCHDFAITERYIIIPDFQLTFCPQRMIRGGQPVIYD
7	30	GRWNFDGQWNQSMTAHPKIDPVTGELFAMG
8	20	TDQPRCQLQGNFAPVEECE
9	29	KWNASYCCRIVQTRFNQEKARGRPGFPK
10	29	WQWNMEFPMINEAYIGRKNKYVYAQIANP




Figure 2. Distribution of conserved motifs of CCO proteins in pepper.

Promoter analyses of *CCO* genes and exon-intron structure

To investigate the transcriptional activity of pepper *CCO* genes, the promoter regions of each *CCO* gene (approximately 1500-bp DNA upstream sequences) were obtained from the pepper genome sequences, and the putative *cis*-regulatory elements were further analyzed using the PlantCARE online tool (Table S1). A total of 88 *cis*-regulatory elements were identified, which could be divided into six categories according to their putative functions. The

first category included those involved in light responsiveness, such as ACE and G-Box. The second category contained eight types of *cis*-acting elements, which were associated with plant hormone responses, including: the TGA-element (auxin-responsive element); TCA-element (salicylic acid responsiveness); abscisic acid-responsive element (ABRE); CGTCA-motif, TGACG-motif (MeJA responsiveness); ethylene-responsive element (ERE); GARE-motif; and TGA-box. The third category responded to biotic and abiotic stresses, including: the TC-rich repeats (involved in defense and stress responsiveness); heat stress responsiveness (HSE); ARE (cis-acting-regulatory element essential for the anaerobic induction); Box-W1 (fungal elicitor-responsive element); MBS (MYB binding site involved in drought responsiveness); LTR (involved in low-temperature responsiveness); and WUN-motif (wound-responsive element). The fourth category was related to plant growth and development and included the following components: MSA-like, circadian, O2-site, CAT-box, CCGTCC-box, Skn-1_motif, GCN4_motif, AACA_motif, and MBSI. The fifth category involved elements associated with the activities of certain biological macromolecules, including 12 types of *cis*-elements. The final category included 19 types of *cis*-elements; however, their specific functions remain unclear. Therefore, it could be speculated that pepper *CCO* genes are involved in plant growth and development, and may also be induced in response to light and hormones, as well as biotic and abiotic stresses.

In addition, the exon-intron patterns of all pepper and tomato *CCO* genes were predicted using the GSDS online tool (Figure 3). The number of introns in pepper *CCO* genes ranged from 0 to 11. Four genes, *CaCCD1*, *CaCCD-like*, *CaCCD-like2*, and *CaCCD-like3*, contained 11 introns. No introns were observed in the *CaNCED*, *CaNCED2*, and *CaNCED3* genes. A similar pattern was found in tomato plants.




Figure 3. Intron-exon structure of *CCO* genes in pepper and tomato.

Expression analysis of *CCO* genes in different pepper tissues using RNA-seq

To determine the transcript levels of *CCO* genes in different pepper tissues, an RNA-seq transcriptome was selected for use in this study (Kim et al., 2014). A total of 24 samples were collected, including roots, stems, leaves, pericarp (PC), and placenta (PL) from the pungent cultivar CM334 and the non-pungent cultivar ECW30R plants, at 6-day post-anthesis (DPA), 16 DPA, 25 DPA, mature green (MG), breaker (B), 5-day post-breaker (B5), and 10-day post-breaker (B10). Figure 4 shows that most of the *CCO* genes were expressed in tissues selected from the pungent cultivar CM334, with the exception of *CaCCD1* and *CaCCD7*. Of the expressed genes, *CaCCD4* was expressed at high levels in the roots and at different stages of fruit development. High expression of *CaCCD8* was observed in the vegetative organs. Two genes, *CaCCD-like* and *CaNCED*, were expressed in all tissues. The remaining two genes, *CaCCD4b* and *CaNCED2*, showed tissue-specific expression.




Figure 4. Expression of *CCO* genes in different tissues of the pungent pepper, CM334, and non-pungent pepper, ECW. PC and B indicate pericarp and breaker stage with days post-anthesis, respectively.

The expression profiles of *CCO* genes were compared in placental tissue from the pungent cultivar CM334 with placental tissue from the non-pungent cultivar ECW (Figure 5). These results showed that three genes (*CaCCD1*, *CaCCD4b*, and *CaNCED2*) were not expressed in any of the tissues studied. In contrast, two genes, *CaCCD-like* and *CaNCED*, were expressed in all of the tissues sampled. *CaCCD8* was expressed at similar levels in the early stages of placental development in CM334 and ECW. Additionally, expression of *CaCCD7* was observed in the ECW-PL-B plant, but not in the CM334-PL-B plant. Similarly, *CaCCD1* was expressed at high levels in the ECW-PL-6DPA and ECW-PL-13DPA cultivars, but not in CM334-PL-6DPA and CM334-PL-16DPA.




Figure 5. Expression of *CCO* genes in placental tissue of the pungent pepper, CM334, and non-pungent pepper, ECW. PL and B indicate placenta and breaker stage with days post-anthesis, respectively.

DISCUSSION

The CCOs are a class of carotenoid cleavage oxygenases, which specifically catalyze the cleavage of conjugated double bonds in carotenoids and pro-carotenoids (Ilg et al., 2009; Walter et al., 2010). These proteins catalyze the cleavage of carotenoids, and help to determine the flavor quality of agricultural products, as well as the catalytic dehydration of the carotenoids, in order to produce a yellow aldehyde, and adjust the plant responses to stress (Espasandin et al., 2014). Therefore, it is beneficial to explore the biological characteristics of *CCO* genes.

In the present study, 11 members in the *CCO* gene family were identified in the pepper genome, while 10 members of the *CCO* gene family were identified in the tomato genome, which is suggestive of a gene duplication or deletion event following the differentiation of pepper and tomato plants. Three *CCD-like* genes (*CCD-like*, *CaCCD-like3*, and *CaCCD-like2*) in the *CCD-like* sub-group were identified in the pepper genome. However, only one *CaCCD-like* gene was identified in the tomato genome. The results of the present study suggest that duplication of *CCD-like* genes in the pepper genome occurred after the differentiation of pepper and tomato plants. However, only one *CCD1* gene (*CaCCD1*) was observed in the pepper genome, and two *CCD1* genes (*SICCD1a* and *SICCD1b*) were observed in the *CCD1* sub-group in the tomato genome. These results indicate that deletion events involving the *CCD1* gene may have occurred in the pepper genome.

In the present study, preferential expression of the *CaCCD4a* genes in roots and fruit was observed. This has previously been reported in other plant species (Sun et al., 2008; Ahrazem et al., 2010; Brandi et al., 2011; Adami et al., 2013). The *CaCCD4b* gene was expressed in a tissue-specific manner in the roots, which suggests a possible function in the growth and development of the roots. Additionally, the highest expression of *CaCCD8* was

observed in the leaves, roots, and stems, followed by the early stages of fruit development. Recently, similar patterns of expression have been observed for orthologous genes of *CaCCD8* in other plant species, which indicates that the putative functions of *CCD8* genes are conserved in different plant species (Zhang et al., 2009; Ledger et al., 2010; Pasare et al., 2013). In conclusion, this comprehensive analysis of the *CCO* gene family lays the foundation for investigation of their potential roles in the future.

Conflicts of interest

The authors declare no conflict of interest.

ACKNOWLEDGMENTS

Research supported by the New Cultivar Breeding Program of Zhejiang Province (grant #2012C12903-1-11), the State Key Laboratory Breeding Base for Zhejiang Sustainable Pest and Disease Control (#2010DS700124-KF1516), the Young Talent Training Program of Zhejiang Academy of Agricultural Sciences (#2015R23R08E09), the Scientific Special of Northern Jiangsu Province (#BN2015096), and the Zhejiang Provincial Natural Science Foundation (#LQ13C030002).

REFERENCES

- Adami M, De Franceschi P, Brandi F, Liverani A, et al. (2013). Identifying a carotenoid cleavage dioxygenase (CCD4) gene controlling yellow/white fruit flesh color of peach. *Plant Mol. Biol. Rep.* 31: 1166-1175. <http://dx.doi.org/10.1007/s11105-013-0628-6>
- Ahrazem O, Trapero A, Gómez MD, Rubio-Moraga A, et al. (2010). Genomic analysis and gene structure of the plant carotenoid dioxygenase 4 family: a deeper study in *Crocus sativus* and its allies. *Genomics* 96: 239-250. <http://dx.doi.org/10.1016/j.ygeno.2010.07.003>
- Auldridge ME, Block A, Vogel JT, Dabney-Smith C, et al. (2006a). Characterization of three members of the *Arabidopsis* carotenoid cleavage dioxygenase family demonstrates the divergent roles of this multifunctional enzyme family. *Plant J.* 45: 982-993. <http://dx.doi.org/10.1111/j.1365-313X.2006.02666.x>
- Auldridge ME, McCarty DR and Klee HJ (2006b). Plant carotenoid cleavage oxygenases and their apocarotenoid products. *Curr. Opin. Plant Biol.* 9: 315-321. <http://dx.doi.org/10.1016/j.pbi.2006.03.005>
- Bartley GE and Scolnik PA (1995). Plant carotenoids: pigments for photo protection, visual attraction, and human health. *Plant Cell* 7: 1027-1038. <http://dx.doi.org/10.1105/tpc.7.7.1027>
- Bouvier F, Isner JC, Dogbo O and Camara B (2005). Oxidative tailoring of carotenoids: a prospect towards novel functions in plants. *Trends Plant Sci.* 10: 187-194. <http://dx.doi.org/10.1016/j.tplants.2005.02.007>
- Brandi F, Bar E, Mourgues F, Horváth G, et al. (2011). Study of 'Redhaven' peach and its white-fleshed mutant suggests a key role of CCD4 carotenoid dioxygenase in carotenoid and norisoprenoid volatile metabolism. *BMC Plant Biol.* 11: 24. <http://dx.doi.org/10.1186/1471-2229-11-24>
- Britton G, Liaaen-Jensen S and Pfander HP (2004). Handbook of Carotenoids. Birkhäuser, Basel.
- Espasandin FD, Maiale SJ, Calzadilla P, Ruiz OA, et al. (2014). Transcriptional regulation of 9-cis-epoxycarotenoid dioxygenase (NCED) gene by putrescine accumulation positively modulates ABA synthesis and drought tolerance in *Lotus tenuis* plants. *Plant Physiol. Biochem.* 76: 29-35.
- Heo J, Kim SH and Lee PC (2013). New insight into the cleavage reaction of *Nostoc* sp. Strain PCC 7120 carotenoid cleavage dioxygenase in natural and nonnatural carotenoids. *Appl. Environ. Microbiol.* 79: 3336-3345. <http://dx.doi.org/10.1128/AEM.00071-13>
- Huang FC, Molnár P and Schwab W (2009). Cloning and functional characterization of carotenoid cleavage dioxygenase 4 genes. *J. Exp. Bot.* 60: 3011-3022. <http://dx.doi.org/10.1093/jxb/erp137>
- Ilg A, Beyer P and Al-Babili S (2009). Characterization of the rice carotenoid cleavage dioxygenase 1 reveals a novel route for geranyl biosynthesis. *FEBS J.* 276: 736-747. <http://dx.doi.org/10.1111/j.1742-4658.2008.06820.x>

- Ilg A, Yu Q, Schaub P, Beyer P, et al. (2010). Overexpression of the rice carotenoid cleavage dioxygenase 1 gene in Golden Rice endosperm suggests apocarotenoids as substrates in planta. *Planta* 232: 691-699. <http://dx.doi.org/10.1007/s00425-010-1205-y>
- Jabeen R, Bhat SV and Wani AA (2013). Functions and stability of lycopene: A review. *Indian J. Horticult.* 3: 57-63.
- Jarquín-Enríquez L, Mercado-Silva E, Maldonado J and Lopez-Baltazar J (2013). Lycopene content and color index of tomatoes are affected by the greenhouse cover. *Sci. Hortic.* 155: 43-48. <http://dx.doi.org/10.1016/j.scienta.2013.03.004>
- Kim S, Park M, Yeom SI, Kim YM, et al. (2014). Genome sequence of the hot pepper provides insights into the evolution of pungency in *Capsicum* species. *Nat. Genet.* 46: 270-278. <http://dx.doi.org/10.1038/ng.2877>
- Lashbrooke JG, Young PR, Dockrall SJ, Vasanth K, et al. (2013). Functional characterisation of three members of the *Vitis vinifera* L. carotenoid cleavage dioxygenase gene family. *BMC Plant Biol.* 13:156 <http://dx.doi.org/10.1186/1471-2229-13-156>
- Ledger SE, Janssen BJ, Karunairetnam S, Wang T, et al. (2010). Modified CAROTENOID CLEAVAGE DIOXYGENASE8 expression correlates with altered branching in kiwifruit (*Actinidia chinensis*). *New Phytol.* 188: 803-813. <http://dx.doi.org/10.1111/j.1469-8137.2010.03394.x>
- Liang YS, Jeon YA, Lim SH, Kim JK, et al. (2011). Vascular-specific activity of the *Arabidopsis* carotenoid cleavage dioxygenase 7 gene promoter. *Plant Cell Rep.* 30: 973-980. <http://dx.doi.org/10.1007/s00299-010-0999-1>
- Liu J, Novero M, Charnikhova T, Ferrandino A, et al. (2013). CAROTENOID CLEAVAGE DIOXYGENASE 7 modulates plant growth, reproduction, senescence, and determinate nodulation in the model legume *Lotus japonicus*. *J. Exp. Bot.* 64: 1967-1981. <http://dx.doi.org/10.1093/jxb/ert056>
- Nagal S, Kaur C, Choudhary H, Singh J, et al. (2012). Lycopene content, antioxidant capacity and colour attributes of selected watermelon (*Citrullus lanatus* (Thunb.) Mansfeld) cultivars grown in India. *Int. J. Food Sci. Nutr.* 63: 996-1000. <http://dx.doi.org/10.3109/09637486.2012.694848>
- Ohmiya A, Kishimoto S, Aida R, Yoshioka S, et al. (2006). Carotenoid cleavage dioxygenase (CmCCD4a) contributes to white color formation in chrysanthemum petals. *Plant Physiol.* 142: 1193-1201. <http://dx.doi.org/10.1104/pp.106.087130>
- Olson JA (1989). Provitamin-A function of carotenoids: the conversion of β -carotenoid into vitamin-A. *J. Nutr.* 119: 94-95.
- Pasare SA, Ducreux LJ, Morris WL, Campbell R, et al. (2013). The role of the potato (*Solanum tuberosum*) CCD8 gene in stolon and tuber development. *New Phytol.* 198: 1108-1120. <http://dx.doi.org/10.1111/nph.12217>
- Rubio A, Rambla JL, Santaella M, Gómez MD, et al. (2008). Cytosolic and plastoglobule-targeted carotenoid dioxygenases from *Crocus sativus* are both involved in β -ionone release. *J. Biol. Chem.* 283: 24816-24825. <http://dx.doi.org/10.1074/jbc.M804000200>
- Schwartz SH, Qin X and Loewen MC (2004). The biochemical characterization of two carotenoid cleavage enzymes from *Arabidopsis* indicates that a carotenoid-derived compound inhibits lateral branching. *J. Biol. Chem.* 279: 46940-46945. <http://dx.doi.org/10.1074/jbc.M409004200>
- Simkin AJ, Schwartz SH, Auldridge M, Taylor MG, et al. (2004). The tomato carotenoid cleavage dioxygenase 1 genes contribute to the formation of the flavor volatiles β -ionone, pseudoionone, and geranylacetone. *Plant J.* 40: 882-892. <http://dx.doi.org/10.1111/j.1365-313X.2004.02263.x>
- Sui X, Kiser PD, von Lintig J and Palczewski K. (2013). Structural basis of carotenoid cleavage: from bacteria to mammals. *Arch. Biochem. Biophys.* 539: 203-213. <http://dx.doi.org/10.1016/j.abb.2013.06.012>
- Sun Z, Hans J, Walter MH, Matusova R, et al. (2008). Cloning and characterisation of a maize carotenoid cleavage dioxygenase (ZmCCD1) and its involvement in the biosynthesis of apocarotenoids with various roles in mutualistic and parasitic interactions. *Planta* 228: 789-801. <http://dx.doi.org/10.1007/s00425-008-0781-6>
- Tamura K, Peterson D, Peterson N, Stecher G, et al. (2011). MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol. Biol. Evol.* 28: 2731-2739. <http://dx.doi.org/10.1093/molbev/msr121>
- Tan BC, Schwartz SH, Zeevaert JA and McCarty DR (1997). Genetic control of abscisic acid biosynthesis in maize. *Proc. Natl. Acad. Sci. U. S. A.* 94: 12235-12240. <http://dx.doi.org/10.1073/pnas.94.22.12235>
- Tan BC, Joseph LM, Deng WT, Liu L, et al. (2003). Molecular characterization of the *Arabidopsis* 9-cis-epoxycarotenoid dioxygenase gene family. *Plant J.* 35: 44-56. <http://dx.doi.org/10.1046/j.1365-313X.2003.01786.x>
- Tracewell CA, Vrettos JS, Bautista JA, Frank HA, et al. (2001). Carotenoid photooxidation in photosystem II. *Arch. Biochem. Biophys.* 385: 61-69. <http://dx.doi.org/10.1006/abbi.2000.2150>
- Vallabhaneni R, Bradbury LM and Wurtzel ET (2010). The carotenoid dioxygenase gene family in maize, sorghum, and rice. *Arch. Biochem. Biophys.* 504: 104-111. <http://dx.doi.org/10.1016/j.abb.2010.07.019>
- Walter MH, Floss DS and Strack D (2010). Apocarotenoids: hormones, mycorrhizal metabolites and aroma volatiles. *Planta* 232: 1-17. <http://dx.doi.org/10.1007/s00425-010-1156-3>

- Wei YP, Wan HJ, Wu ZM, Wang RQ, et al. (2016). A comprehensive analysis of carotenoid cleavage dioxygenases genes in *Solanum lycopersicum*. *Plant Mol. Biol. Rep.* 34: 512-523. <http://dx.doi.org/10.1007/s11105-015-0943-1>
- Winterhal TP and Schreier P (1995). The generation of norisoprenoid volatiles in star fruit (*Averrhoa carambola* L.): A review. *Food Rev. Int.* 11: 237-254. <http://dx.doi.org/10.1080/87559129509541041>
- Woitsch S and Römer S (2003). Expression of xanthophyll biosynthetic genes during light-dependent chloroplast differentiation. *Plant Physiol.* 132: 1508-1517. <http://dx.doi.org/10.1104/pp.102.019364>
- Zhang M, Yuan B and Leng P (2009). Cloning of 9-cis-epoxycarotenoid dioxygenase (NCED) gene and the role of ABA on fruit ripening. *Plant Signal Behav.* 4: 460-463. <http://dx.doi.org/10.4161/psb.4.5.8542>

Supplementary material

Table S1. Information on 88 different cis-regulatory elements identified in 11 *CaCCO* genes, including their numbers and the function of the respective elements.

Figure S1. Multiple-sequence alignments of SICCO and CaCCO proteins.