

Methodology

Simplified preparation of a DNA ladder using PCR

T.-Y. Wang, L. Wang and F. Wang

Department of Biochemistry and Molecular Biology, Xinxiang Medical University, Henan, P.R. China

Corresponding author: T.-Y. Wang E-mail: wtianyuncn@126.com

Genet. Mol. Res. 10 (3): 1631-1635 (2011) Received November 29, 2010 Accepted February 10, 2011 Published August 5, 2011 DOI http://dx.doi.org/10.4238/vol10-3gmr1177

ABSTRACT. Serving as a DNA molecular weight standard, the DNA ladder has been widely used in molecular biology applications. We developed a simple method for the preparation of a DNA marker, which involves designing primers to amplify 100- to 1000-bp DNA fragments using lambda DNA as a template for polymerase chain reaction, followed by extraction with phenol/chloroform, precipitation with ethanol and mixing. Fragments of 100- to 1000-bp DNA were successfully amplified; the sequences showed 100% identity with lambda DNA. This prepared DNA marker displayed clear bands, indicating that it can be used for molecular studies.

Key words: DNA; Ladder; λ DNA; Polymerase chain reaction

©FUNPEC-RP www.funpecrp.com.br

Genetics and Molecular Research 10 (3): 1631-1635 (2011)