

<u>Technical Note</u>

## Isolation and characterization of polymorphic microsatellite loci in the hard-shelled mussel, *Mytilus coruscus* (Mytilidae)

T.-J. Xu, Y.-N. Sun, Y.-T. Yuan, Z. Liao and R.-X. Wang

Key Laboratory for Marine Living Resources and Molecular Engineering, College of Marine Science, Zhejiang Ocean University, Zhoushan, P.R. China

Corresponding author: R.-X. Wang E-mail: wangrixin1123@126.com

Genet. Mol. Res. 9 (3): 1388-1391 (2010) Received March 19, 2010 Accepted May 20, 2010 Published July 13, 2010 DOI 10.4238/vol9-3gmr856

**ABSTRACT.** *Mytilus coruscus* is one of the most important cultured species of marine shellfish in China. Using an expressed sequence tag-library and two microsatellite-enriched genomic libraries of *M. coruscus*, we isolated and characterized 12 polymorphic microsatellites in a test population; the number of alleles ranged from three to seven, and the observed and expected heterozygosities ranged from 0.0333 to 0.8571 and from 0.3452 to 0.8267, respectively. Four loci deviated from Hardy-Weinberg equilibrium. These polymorphic microsatellite loci can be employed in genetic diversity analysis and molecular marker-assisted breeding of *M. coruscus*.

Key words: Mytilus coruscus; Microsatellite; Molecular marker

©FUNPEC-RP www.funpecrp.com.br

Genetics and Molecular Research 9 (3): 1388-1391 (2010)